


a brief introduction to Cloudy 95

results, environment

G.J. Ferland

Department of Physics and Astronomy University of Kentucky, Lexington <u>http://www.pa.uky.edu/~gary/cloudy</u>



Use of this program is not restricted provided each use is acknowledged upon publication. The bibliographic reference to this version of **Cloudy** is Ferland, G.J., 2001, *Hazy*, *a Brief Introduction to Cloudy*, University of Kentucky Department of Physics and Astronomy Internal Report.

Portions of this document have been published, and are copyrighted by, The American Astronomical Society and The Royal Astronomical Society. The remainder of this document, and the code *Cloudy*, are copyrighted 1978-2001 by Gary J. Ferland.

Cloudy is an evolving code. Updates are made on a roughly quarterly basis, while major revisions occur roughly every three years. You should confirm that you have the most recent version of the code by checking the web site http://www.pa.uky.edu/~gary/cloudy or asking to be placed on the **Cloudy** mailing list.

January 29, 2001.

CLOUDY 95

G. J. Ferland Department of Physics and Astronomy University of Kentucky Lexington

Table of Contents

1 MACHINE ENVIRONMENT	. 401
1.1 Overview	. 401
1.2 Floating Point Environment	. 401
1.3 Cloudy via ftp or the Web	. 401
1.4 Cloudy in Lexington	. 401
1.4.1 Versions of the Code	. 401
1.4.2 Running a single model	. 402
2 CLOUDY AS A SUBROUTINE	
2.1 Overview	. 404
2.1.1 Create the Cloudy object files	. 404
2.1.2 Computing a grid of models	. 405
2.2 Writing a New Main Program	
2.2.1 The cddrive.h header file	. 405
2.2.2 A note on return conditions	
2.2.3 Initializing the code	. 406
2.2.4 cdMPI - telling the exit handler about MPI	
2.2.5 cdRead - entering Commands	
2.2.6 cdOutp - sending output to a file	
2.2.7 cdTalk - produce output??	
2.2.8 cdDrive - calling the Code	
2.2.9 cdNoExec - checking without Computing	
2.3 Checking Predictions	
2.3.1 cdLine - emission lines intensities	
2.3.2 Special arrays of emission lines	
2.3.3 cdEmis - emissivity of lines	
2.3.4 cdColm - the computed column densities	
2.3.5 cdIonFrac - the computed ionization fractions	
2.3.6 cdTemp - the computed mean temperature	. 410
2.3.7 cdGetCooling, cdGetHeating – the cooling and heating in the	
last zone	
2.3.8 cdGetTe - reading the temperature	
2.3.9 cdGetPres – pressure	, 411
2.3.10 cdTimescales – several timescales	
2.4 Other information	
2.5 Printing the Comments	, 411

	2.5.1 Were comments generated?	411
	2.5.2 Printing the comments.	412
	2.5.3 cdErrors(FILE *io) - printing a summary of any problems	
	2.5.4 cdPrintCommands(FILE *io) – print the command stack	
	2.5.5 cdGetLineList – get list of emission lines	
	2.5.6 cdGetFeIICont – bin FeII emission into wavelength cells	
	2.5.7 cdGetFeIIBands – enter list of FeII bands into emission line	
	stack	413
	2.5.8 cdSPEC – get continuous spectra	414
	2.6 Example Call as a Subroutine	
	2.7 Storing Grids of Calculations	
	2.8 Computing Grids of Calculations	
3 (OUTPUT	
-	3.1 Overview	
	3.2 Header Information	
	3.2.1 Initial Information	
	3.2.2 Properties of the Continuum	
	3.3 Zone Results	
	3.3.1 Line 1	
	3.3.2 [Optional] wind parameters	
	3.3.3 [Optional] radiation pressure	
	3.3.4 Line 1 - Hydrogen I	
	3.3.5 Line 2 - Hydrogen II	
	3.3.6 Line 3 - Helium	
	3.3.7 Line 4 - Atomic Helium	422
	3.3.8 Line 5 - Ionized Helium	422
	3.3.9 Optional Grains	422
	3.3.10 Pressure	422
	3.3.11 Molecules	423
	3.3.12 Li, Be, B	423
	3.3.13 Carbon	423
	3.3.14 Nitrogen	423
	3.3.15 Oxygen	423
	3.3.16 Fluorine, Neon	423
	3.3.17 Remaining Elements	423
	3.4 Calculation Stopped Because	423
	3.4.1 because of radiation pressure	
	3.4.2 because lowest EDEN reached	
	3.4.3 because low electron fraction	
	3.4.4 because wind veloc too small	424
	3.4.5 because code returned BUSTED	424
	3.4.6 because DRAD small - set DRMIN	
	3.4.7 because DR small rel to thick	425
	3.4.8 because carbon fully molecular	
	3.4.9 because negative mole abundan	425
	3.4.10 because optical depth reached	425

3.4.11 because outer radius reached	. 425
3.4.12 because column dens reached	. 425
3.4.13 because lowest Te reached	. 426
3.4.14 because highest Te reached	. 426
3.4.15 because NZONE reached	. 426
3.4.16 because line ratio reached	. 426
3.4.17 because internal error - DRAD	. 426
3.4.18 because initial conditions out of bounds	. 426
3.4.19 because reason not specified	. 426
3.5 Geometry	. 426
3.6 Warnings, Cautions, Surprises, and Notes	. 426
3.7 Optional Plot	. 427
3.8 Final Printout	. 428
3.8.1 Emission-Line Spectrum	. 428
3.8.2 Thermal balance	. 431
3.8.3 Column densities, et al.	. 432
3.8.4 Averaged Quantities	. 433
3.8.5 Grains	
3.8.6 Continuum optical depths	. 435
3.8.7 Mean ionization	
3.8.8 Continuum	. 436
4 OBSERVED QUANTITIES	. 437
4.1 Overview	
4.2 Incident and Diffuse Continua	. 437
4.3 Line Equivalent Widths	. 437
4.4 Emission Line Asymmetries	. 438
4.5 Line to Continuum Contrast	. 438
4.6 Surface Brightness	. 439
4.7 Flux to luminosity	. 439
4.8 Relative hydrogen line intensities	. 439
4.9 Line Intensities in a dusty open geometry	. 440
4.10 Continuum pumping contribution to line intensities	. 440
5 THE EMISSION LINES	. 442
5.1 Overview	
5.2 Line wavelengths	. 442
5.3 Miscellaneous Line Entries	
5.4 Emission line identifications	. 451
5.4.1 Optically thin recombination lines	. 451
5.4.2 Transferred heavy element lines	
5.4.3 Output produced for the transferred lines	
5.5 The main blocks of lines	
5.6 Atomic data sources	. 453
6 PROBLEMS	. 456
6.1 Overview	
6.2 Thermal stability and temperature convergence	

6.2.1 Types of thermal maps	456
6.2.2 No Temperature Convergence	458
6.2.3 Thermal Stability	
6.2.4 Thermal fronts	
6.2.5 Map Output	459
6.3 Floating Point Errors	460
6.4 Optical depth convergence problems	460
6.5 Negative Populations of H, He	460
6.6 I can't fix it if I don't know its broken	460
7 REVISIONS TO CLOUDY	462
7.1 Overview	462
7.2 Cloudy and Moore's Law	
7.3 Major Past Versions	
7.4 Version 94 versus 90	
7.4.1 Commands	465
7.4.2 Hydrogen	
7.4.3 grains	465
7.4.4 Other changes	
7.5 Version 90 versus 84	
7.5.1 Commands	465
7.5.2 Continuum Transport	465
7.5.3 Hydrogen	
7.5.4 The helium ion	
7.5.5 Heavy elements	466
7.5.6 Free-free, line heating and cooling	467
7.5.7 Excited state photoionization cross sections	
7.5.8 The O ⁺ photoionization cross section	
7.6 Version 84 versus 80	
7.6.1 Commands	468
7.6.2 Mg II λ2798	468
7.6.3 General Results	
7.7 Known Modes for Cloudy 95	468
7.8 Making a Revision	468
7.8.1 The code	
7.8.2 Printing Hazy	469
8 COMPARISON CALCULATIONS	
8.1 Overview	
8.2 Cool HII Region	471
8.3 Paris HII Region	
8.4 Blister HII Region	
8.5 Paris Planetary Nebula	
8.6 Paris NLR Model	
8.7 Lexington NLR Model	
8.8 The DQ Her Shell	
8.9 The Kwan and Krolik Standard Model	

8.10 Rees, Netzer, and Ferland, low density	485
8.11 Rees, Netzer, and Ferland, high density	
9 SAMPLE INPUT	487
10 REFERENCES	489
11 INDEX	497

List of Figures

A grid of model calculations	404
Reflected continuum	438
Reflected continuum	440
Ccooling function for low density photoionized gas	456
Cooling function for low density collisionally ionized gas	457
Temperature as a function of density for photoionized gas	458
Cloudy's size as a function of time.	462
O+ photoionization cross section	

List of Tables

Cool HII Region	471
Cool HII Region vs Cloudy	471
Paris meeting HII region	472
Paris HII region vs cloudy	473
Blister HII Region	
Blister HII Region vs Cloudy	
Paris meeting planetary nebula	477
Paris planetary vs Cloudy	478
Paris meeting NLR model	
Paris NLR model vs CLOUDY	
Lexington NLR Model	481
Lexington NLR vs Cloudy	482
DQ Her Shell	483
Kwan and Krolik Standard Model	484
Rees, Netzer, Ferland low density BLR	
Rees, Netzer, Ferland high density BLR	
Single model test cases	

1 MACHINE ENVIRONMENT

1.1 Overview

This section describes the machine environment needed to run *Cloudy*. The code is designed to run on IEEE 32-bit computers, and has been tested on a Sparc, SGI, Alpha, HP Exemplar, and PC. Calculations on all machines produce essentially identical results.

The **readme.htm** file that is included with the source contains the most recent information about compiling and setting up the code.

1.2 Floating Point Environment

The floating-point environment should be set to ignore floating-point underflow but crash on any other floating-point error. Floating-point underflow is an unavoidable consequence of the attenuation of radiation as a beam of light is extinguished by an absorbing medium; underflow error checking should be disabled.

Floating point overflow or division by zero *must never* occur, nor should library function domain errors (i.e., the log of a negative number). I would appreciate hearing about these errors. I can't fix it if I don't know it is broken. My email address is gary@cloud9.pa.uky.edu. Please send the input file and version number.

1.3 Cloudy via ftp or the Web

The home page for **Cloudy** is http://www.pa.uky.edu/~gary/cloudy. This is the preferred method of obtaining the code. This is the best place to retrieve the source since it is the only place certain to have the current version.

Cloudy is an evolving code. Bugs are fixed as soon as they are discovered, and major revisions occur from time to time. It is important to check the web page several times per year to see what changes and improvements have occurred. I also maintain a mailing list to announce changes to the code. To be placed on this list send a request to gary@pa.uky.edu.

I would like to keep track of what the code is being used for, if only to try to make sure that it is within it range of validity. I would appreciate receiving preprints of any work done using *Cloudy*.

The bibliographic reference to this version of **Cloudy**, which should be cited in any paper that uses it, is given on the inside cover of this document. The citation should include the subversion number, i.e., version 95.09d.

1.4 Cloudy in Lexington

1.4.1 Versions of the Code

Many versions of **Cloudy** live on the UNIX network in Lexington. In general there will two versions of the code - a fully tested (older) version, and the current development version.

Software development goes through several phases. The initial version of a code is often referred to as an "alpha" version, the nearly de-bugged version as a "beta" version, and the fully tested version as the "gold" code. At any one time the last gold version of the code will live in a directory with a name indicating the version number. The version of *Cloudy* is indicated by two parts - an integer indicating the version number, and a decimal part (perhaps with a letter following it) indicating the sub-version. Sub-versions to the gold code indicate fixes of bugs or stability problems, or major changes in the atomic data. The last gold version of *Cloudy* was 95. An example of the current gold version number might be something like 95.09d. The source and executable for this version will live in a sub-directory with the name gary/c95. The current development version of the code lives in a directory called gary/cloudy.

Within the sub-directories the architecture is the same. The Fortran source files have names that end in .for. The executable has a name that indicates which processor it is intended for. Executable for a Sparc will have the name c.sun4, while those for the alpha will have the name c.alpha.

1.4.2 Running a single model

Cloudy is often used to read in the parameters for a single model, and compute the result. The easiest way to do this is to create a small file that contains the input stream for that model. As a typical case consider a simple planetary nebula:

hden 4 radius 17 black body 100,000K, luminosity 38

Assume this is saved as the file **pn.in**. (Cloudy stops reading the input steam when it reaches a blank line, or the end of file.)

I created a shell script with the name **run** which is in my "bin" directory, which I include on my path. The shell script **run** consists of the following:

```
echo reading input file $1.in
case $# in
0) echo there must be an input file ;;
1) /homeb/uwc0/gary/cloudy/c.sun4<$1.in >$1.out
echo created output file $1.out ;;
2) /homeb/uwc0/gary/cloudy/c.sun4 < $1.in >$2.out
echo created output file $2.out ;;
esac
echo $p
exit
```

run normally works by reading in a file called file.in, and creating a file called
file.out.

If **run** is executed with no input parameters it will complain that at least one argument is needed and then stop. If there is one parameter it is treated as the name of the input and output files. So in the above example, typing

run pn

would read the input stream in **pn.in** and create an output file of results called **pn.out**. When two parameters occur the first is the name of the input stream and the second is the name of the output steam. The example

run pn test

would read the file **pn.in** and create the file **test.out**.

2 CLOUDY AS A SUBROUTINE

2.1 Overview


It is possible to use **Cloudy** as a subroutine of other, much larger, codes. When used this way a series of subroutine calls, described next, are used to initialize the code, specify the initial conditions, drive the code, and finally examine the predictions.

This document gives an overview of all the routines that are intended to be "public" (needed to be accessed by agents other than **Cloudy** itself). The definitions for all public routines are contained in the header file *cddrive.h*, which gives the best and current description of all these routines. That file is the definitive reference source for all of the material in this chapter.

A common strategy is to call the code to compute line intensities for a large matrix of parameters. The results of one such calculation is shown in Figure 1. Such grids can be computed in a few dozen hours on modern workstations, and offer far greater insight to physical effects of changing model parameters, than does a single model.

2.1.1 Create the Cloudy object files

The goal is to replace the existing *Cloudy* main program with one that you write.



[Call] 7306/[SII] 6720

Figure 1 The results of a large grid of model calculations are shown. The x-y plane show the logs of the hydrogen density and stellar temperature. The z axis is the indicated line ratio, for solar abundances.

The file *maincl.c* that is included in the distribution must be deleted so that the program you write will be loaded instead. The remaining routines are then compiled with a command like the following:

```
gcc -0 -c *.c
```

which will create a large number of object files. The following subsections outline how to write code for this new main program.

2.1.2 Computing a grid of models

Next compose a main program that drives **Cloudy** by calling a series of routines (described beginning on page 404 of this document).

If the main program is called *grid1.c*, it could be compiled and linked with the *Cloudy* object files as follows. I define a shell script named "compile" which contains the commands

```
echo Begin compilation of $1.c
gcc -O -c $1.c
echo Compilation of $1.c is complete
```

so that *grid1.c* could be compiled with the command

compile grid1

Next it is necessary to link *grid.o* (produced above) into *Cloudy* itself. This is done with a shell script I called **linkedit**, which consists of the following:

```
echo about to link $1.0 using cloudy object files
gcc -0 -0 $1.exe $1.0 `ls /homeb/uwc0/gary/cloudy/*.o` -lm
echo created executable $1.exe
```

The command

linkedit grid1

will produce the executable file grid1.exe. Executing it as

grid1.exe > grid1.out

will produce the output file grid1.out.

2.2 Writing a New Main Program

2.2.1 The cddrive.h header file

The file *cddrive.h* is included in the distribution and contains definitions of all public routines, the routines that a user would call to drive *Cloudy*. That file is the definitive reference for the material contained in this section. Comments within the file explain all routines and their parameters. Often I copy from the directory containing the majority of the *Cloudy* source to other directories where special main routines are developed. It is very important that all versions of *cddrive.h* be updated with the source is updated since the file changes with time.

2.2.2 A note on return conditions

Some of the following routines return a value to indicate success or failure. I try to follow the C and Unix conventions to indicate success with zero and trouble with a non-zero return. This rule is not always followed however and *cddrive.h* should be consulted to make sure the return conditions are understood.

2.2.3 Initializing the code

Many variables must be initialized at the beginning of the calculation. Calling subroutine *cdInit* does this.

cdInit();

Routine *cdInit* must be called every time a new calculation is to be performed, *before* calling any of the following subroutines, but after the results of previous calculations have been read.

2.2.4 cdMPI - telling the exit handler about MPI

If MPI is used for parallel processing then the code must clean up by calling *MPI_Finalize* at exit. The exit handler is defined in *cdInit* and is called *exithandler*. Routine *cdMPI* must be called after *cdInit* and before *cdDrive* for *MPI_Finalize* to be called correctly.

2.2.5 cdRead - entering Commands

Command lines are entered by successive calls to routine *cdRead*. The argument of *cdRead* is the line image, a null-terminated string. The commands consist of a series of 80 character free-format command lines. They must obey all the rules outlined in Part I.

In the examples below some commands are directly entered as strings and others created by writing variables through *sprintf* (a standard C function).

```
char chLine[132];/* this will hold the command lines we will generate*/
cdRead("title Gene's series of constant pressure models" );
hden = 5.4;
sprintf( chLine , "hden %5.2f ", hden );
cdRead(chLine );
cdRead(chLine );
sprintf( chLine , "coronal %5.2f ", temp );
cdRead(chLine );
```

cdRead("stop zone 1 ");

cdRead returns the number of commands that can still be entered before exceeding the size of the storage arrays. So this routine is an exception to the general rule that a zero return condition indicates success – here it means that no further commands can be entered. When it decrements to 0 (false) no more lines can be entered. The return value was ignored in the examples above.

It is not now possible to read in more than 1000 command lines because of limits to the size of the character arrays used to store them. This limit is stored as the variable *nkrd*. If more than 1000 lines are read in by calling *cdRead* then *cdRead* will stop after explaining why. It will be necessary to increase *nkrd* if more than 1000 command lines are needed.

2.2.6 cdOutp - sending output to a file

Cloudy normally writes its standard output on the system's **stdout**. This can be changed to another stream by calling routine *cdOutp*, which has a file handle to an open file as its single argument. By combining this redirection with the C **fopen** statement it is possible to have the standard output sent into any file.

```
#include "cddriv.h"
/* this defines a standard C file handle */
FILE *ioData ;
    /* open the file output.txt for writing */
    ioData = fopen("output.txt","w");
    /* ioData is equal to NULL if we failed to open the file */
    if( ioData==NULL )
    {
        exit(1);
    }
    /* send output to this file*/
    cdOutp( ioData ) ;
    -  - code goes here
    /* at end of calculation close the file */
    fclose(ioData);
```

2.2.7 cdTalk - produce output??

Cloudy normally speaks what's on its mind. However, it does have a quiet mode in which nothing at all is printed. This quiet mode is set by the logical argument to subroutine *cdTalk*.

```
/*set no output at all*/
cdTalk( FALSE )
/*have the code produce the normal printout*/
cdTalk( TRUE )
```

The default is for *Cloudy* to produce output, and *cdTalk* does not have to be called if this is desired. However, it does need to be called with the logical variable *false* if the quiet mode is desired.

2.2.8 cdDrive - calling the Code

The calculation is performed when routine *cdDrive* is called. *cdDrive* returns an int indicating whether the calculation was successful. The value 0 indicates a successful calculation. The following shows an example of its use.

```
int lgOK;
if( cdDrive() )
{
    printf("problems!\n");
    exit(1);
}
```

If problems occurred and the results cannot be trusted then the return value is non-zero. This will only be set if the calculation suffered a complete meltdown. Routine *cdNwcns* (page 411) can be called to find out about less serious problems.

2.2.9 cdNoExec - checking without Computing

If routine *cdNoExec* is called after *cdInit* but before *cdDrive* then only the initial parts of a calculation will be performed when routine *cdDrive* is called.

```
CdInit();
/*read in commands */
cdRead( . . .);
/*tell it not to execute */
cdNoNoex();
/*call the code */
lgBad = cdDrive();
```

When *cdDrive* is called the code will generate the incident continuum, set the initial density, and the chemical composition. It will then stop just before the initial search for the physical conditions in the first zone. All of the initial printout, summarizing properties of the composition and continuum, will be generated. This provides a quick way to check that a large grid of models will be specified correctly, without actually fully computing the grid.

2.3 Checking Predictions

2.3.1 cdLine - emission lines intensities

The predicted line intensities or luminosities for all lines with non-zero values are stored within a set of structures which also contain the line identifiers, a four character label and integer wavelength. These are normally printed at the end of the calculation. You obtain the line intensity by calling subroutine *cdLine*. The label and wavelength of the line are specified in the call, and the routine returns the relative intensity and the log of the absolute intensity or luminosity.

```
Include "cddefines.h"
*
if( cdLine( "totl" , 4861 , &relint , &absint ) <= 0)
    printf("did not find this line\n");</pre>
```

The first variable is the line label, the four-character null-terminated string (upper or lower case) as used by the code to identify the line. The second integer variable gives the wavelength of the line. Both of these must exactly match the label and wavelength used by *Cloudy* to identify the line (see the chapter "Lines" for a full description). The third variable (*relint* in the above example) is a pointer to the double precision relative intensity of the line (relative to the normalization line, usually H β , but set with the **normalize** command). The log of the intensity (erg cm² s⁻¹) or luminosity (erg s⁻¹) of the line is returned as the last pointer to a double precision variable (*absint* in the above example). If the intensity of the line is zero or the line was not found then this variable will be set to -37.

If *cdLine* finds the line it returns the index of the line within the stack of emission lines. It returns the negative of the total number of lines in the stack if the line is not found. This may occur if the line wavelength or label were mistyped. This is an exception to the normal, C-like, function return convention in which a normal return is zero and an abnormal return non-zero.

2.3.2 Special arrays of emission lines

The routine *cdGetLineList* provides a way to access a block of lines in an automatic manner. This provides a way to access sets of predictions for either storage or printout.

cdInit must be called first to initialize needed variables. Then *cdGetLineList* is called as described here. Finally, the actual grid of calculations begins with another call to *cdInit*. Finally the intensities of a set of lines is extracted by calls to *cdLine*. So the first call to *cdInit* is generally not followed by execution of the code when these routines are used.

When routine *cdGetLineList* is called a list of lines, from either one of two files included in the standard data distribution, or user-created special file, will be entered

into a pair of vectors. This list can then be used to call *cdLine* to obtain intensities of these lines. The argument to the routine is the name of the file containing the lines. If a null string is passed ("") then *BLRLineList.dat* is used. That file is included in the standard data distribution and contains a description of the required file format. A second file, *NLRLineList.dat*, is also included in the standard distribution and contains lines appropriate for a low-density gas.

If the file was read successfully then the routine returns the number of lines contained in the file. If problems occurred then a negative integer is returned.

When the routine has been called the code creates a pair of vectors, *cdGetchLabel[n]* and *cdGetnWL[n]*. The first is an array of four-character strings that are emission line labels. The second vector is a long integer array of wavelengths. This string and integer are the label and wavelength used to identify the lines.

The following shows an example of getting the lines from *BLRLineList.dat*, executing the code, then obtaining the predicted intensities by calling *cdLine*.

```
/* initialize the code */
cdInit();
/* get list of lines from standard data file */
if( (nLines=cdGetLineList("")) < 0 )</pre>
      /* this is the error exit - could not obtain the lines */
      exit(1);
}
/* now set up the actual call to the code */
cdInit();
/* missing commands here, then call the code */
cdDrive();
/* now print the emission lines that were in the file */
for( n=0; n<nLines; ++n )</pre>
      lgOK = cdLine( cdGetchLabel[n], cdGetnWL[n] , &relative , &absolute );
      if (lgOK <= 0)
       {
             fprintf(stderr,"did not find %4s%5li\n",cdGetchLabel[n],cdGetnWL[n]);
             fprintf(ioDATA,"\ndid not find line.\n");
       }
      grid.pred[n] = log10(MAX2(1e-30,relative));
}
```

2.3.3 cdEmis - emissivity of lines

A second routine, *cdEmis* functions much the same as *cdLine* but returns the local emissivity (erg cm⁻³ s⁻¹ for unit filling factor) of the line for the last computed zone. The return value is the index of the line with the line stack if it was found, and the negative of the number of lines in the stack if the line could not be found.

2.3.4 cdColm - the computed column densities

The predicted column densities can be accessed by calling the routine *cdColm*.

```
chLabel = "carb";
ion = 3;
cdColm("carb", ion , &colum);
if(cdColm("carb", 3 , &colum))
{
        printf(" could not find C+2 - check in put\n");
}
else
```

```
printf("The predicted C+2 column density is e\n", column );
```

The routine returns 0 if it found the species, and 1 if it could not. It returns the predicted column density (linear, cm⁻²) as the third argument. The first argument *chLabel* is a four character identifier which must agree with the first four characters (upper or lower case) of the name used to indicate the element in the printout. The integer variable *ion* is the spectroscopic designation of the level of ionization, i.e., 1 indicates C I, 3 indicates C III, etc. The return value indicates whether the species was found.

2.3.5 cdIonFrac - the computed ionization fractions

The predicted ionization fractions, averaged over radius or volume, can be accessed by calling the subroutine *cdIonFrac*.

```
if( cdIonFrac( "carb" , 2 , &frac , "radius",FALSE) )
{
    exit(1);
}
printf("The predicted ionization fraction over radius is is%g\n", frac);
```

The routine returns the predicted ionization fraction A_{ion}/A_{tot}. *chLabel* is a four character identifier which must agree with the first four characters (upper or lower case) used to indicate the element in the printout. The integer variable *ion* is the spectroscopic designation of the level of ionization, i.e., 1 indicates C I or the atom, 3 indicates the second ion C III, etc. *chWeight* is a six character variable which must be either "radius" or "volume" (either upper or lower case). This determines whether the ionization fraction returned is weighted over radius or volume. The last variable determine whether (TRUE) or not (FALSE) the electron density is included in the weighting. The function returns zero if the ion was found, and non-zero if an error occurred.

2.3.6 cdTemp - the computed mean temperature

Routine *cdTemp* returns the computed mean temperature. The first parameter is a 4 character plus null termination string giving the name of an element. The second is the ionization stage, with 1 the atom, 2 the first ion, etc. The third parameter will be the computed mean temperature. The last parameter is a 6 character plus null termination string, either "radius" or "volume" that says whether the temperature should be weighted with respect to radius or volume. The routine returns 0 if it finds the species, and 1 if it could not find the species.

```
if( cdTemp( "carb" , 2 , &temp , "radius") )
{
     exit(1);
}
printf("The mean temperature is%g\n", temp);
```

The label "21cm" for the element name has a special meaning. The temperature will be the mean of n_H/T , the harmonic mean weighted with atomic hydrogen density, over radius. This is the temperature measured by 21 cm observations.

{

}

2.3.7 cdGetCooling, cdGetHeating - the cooling and heating in the last zone

The total cooling and heating for the last computed zone is obtained by called the void functions *cdGetCooling*. and *cdGetHeating*. The function returns the value as a double precision variable. In equilibrium cooling and heating are equal.

2.3.8 cdGetTe - reading the temperature

The electron temperature for the last computed zone is obtained by called the void function *cdGetTe*. The function returns the predicted temperature as a double precision variable.

2.3.9 cdGetPres – pressure

The pressure for the last computed zone is obtained by called routine *cdGetPres*. This routine has three arguments, pointers to the total (gas plus radiation) pressure, the gas pressure, and the radiation pressure. All are double precision variables.

2.3.10 cdTimescales - several timescales.

This routine has three arguments, pointers to doubles that will return the thermal timescale, the hydrogen recombination timescale, and the H₂ formation timescale.

2.4 Other information

cdDate(string) The date when the current version of the code was released will be placed as a null terminated string into the string passed as an argument. The calling program must allocate enough room in the string – the length is specified in *cddrive.h.*

cdVersion(string) The code's version number will be placed as a null terminated string into the string passed as an argument. It is a string rather than a float since it usually ends with a letter of the alphabet. The calling program must allocate enough room in string – the length is specified in *cddrive.h.*

double cdExecTime(void) This returns the time that has elapsed since the previous call to *cdInit*.

2.5 Printing the Comments

After the calculation is complete, but before the emission lines are printed, the code generates a series of statements that indicate warnings, cautions, comments, and surprises. These should be examined to confirm that the calculation is probably all right. A series of subroutines allows the driving code to determine whether these comments were generated, what type they were, and to print then on an arbitrary IO unit.

2.5.1 Were comments generated?

A series of routines are provided to check on problems in the calculation.

Routine *cdNwcns* will return the number of warnings, cautions, surprises, notes, and temperature and pressure failures:

cdNwcns(&lgAbort , &nw , &nc , &nn , &ns , &nte , &npe , &nione, &neden)

where the first variable is a flag indicating whether the calculation aborted, *nw* is the number of warnings generated (if this number is non-zero, then the calculation has serious problems), *nc* is the number of cautions generated (these are less severe than

warnings, but are still a cause of concern), and *nn* and *ns* are the number of notes and surprises. The next two arguments are the number of temperature and pressure failures. The last two are the number of ionization and electron density failures. All of these integer values should add up to zero in a successful calculation.

If either of the first two variables are non-zero then the code returned with an indication of serious failure. An abort is far more serious than a warning since it indicates total meltdown. I would appreciate learning about these.

2.5.2 Printing the comments.

The comments may be printed on any file by calling the series of subroutines described here. In all cases the routines take as an argument a file handle, which must point to a file open for writing.

```
/* output the comments into a file, but first open it */
FILE *ioOUT;
If( (IoOUT = fopen( "comments.txt", "w") ) == NULL )
{
      printf("error creating comments.txt file\n");
      exit(1);
/*print the reason the calculation stopped, and geometry*/
 cdReasonGeo( ioOUT )
/*print the warnings*/
 cdWarnings(ioOUT)
/*next print the cautions*/
 cdCautions(ioOUT)
/*now print any surprising results*/
 cdSurprises(ioOUT)
/*now print the notes
 cdNotes(ioOUT)
```

cdReasonGeo (*FILE *io*). It is very important to understand why the calculation stopped. The first two lines after the last zone results give the reason the calculation stopped, and the type of geometry. This information will be printed on the file whose handle is the argument, when this routine is called.

cdWarnings(FILE *io) All warnings will be printed on the output file.

cdCautions (FILE *io) All cautions will be printed on the output file.

cdSurprises(FILE *io) All surprises (denoted by a "!") are printed on the output file.

cdNotes(*FILE *io*) The notes concerning the calculation will be printed on the output file.

2.5.3 cdErrors(FILE *io) - printing a summary of any problems

Routine *cdErrors*(*FILE *io*) will generate a summary of any problems that happened during a calculation. The argument is a file pointer to the output file where the results will be placed. If problems occurred in the calculation, such as temperature or pressure failures, warnings, or cautions, these will be printed following the title for the calculation.

2.5.4 cdPrintCommands(FILE *io) - print the command stack

The single argument is a file handle. The entire series of input commands will then be printed on the file. The commands are preceded and followed by lines that begin with "c =====" to easily identify.

2.5.5 cdGetLineList - get list of emission lines

This command can read in an arbitrary list of emission lines to print at the end of the calculation. When called with the null string it will read in a list of typical BLR lines from the file *BLRLineList.dat*, which is included in the data distribution. If the string argument is not null it will attempt to read in a file with that name instead. The file location is assumed to be on the path as set with the **path** command, or in *path.c*.

The global variables *cdGetchLabel* and *cdGetnWL* are filled with labels and wavelengths for the lines in the file. The function returns a long int, the number of lines in the list. A typical spectrum could be pulled from the results and printed by including the following code after the call to *cdDrive*:

```
CdInit();
nLines = cdGetLineList("");
cdDrive();
for( n=0; n<nLines; ++n )
{
    lgOK = cdLine( cdGetchLabel[n], cdGetnWL[n] , &relative , &absolute );
    if( lgOK<=0 )
    {
        fprintf(stderr,"did not find %4s%5li\n",cdGetchLabel[n],cdGetnWL[n]);
    }
    grid.pred[n] = log10(MAX2(1e-30,relative) );
}
```

2.5.6 cdGetFeIICont - bin FeII emission into wavelength cells

Emission from the model FeII atom is rich and complex and there are far too many lines for a simple analysis to succeed. This command rebins the FeII spectrum into a series of cells to simulate a full pseudo-continuum. There are three arguments, the lowest and highest wavelengths (in vacuum Ångstroms) and an integer third argument, and number of cells to divide it into.

2.5.7 cdGetFeIIBands - enter list of FeII bands into emission line stack

A number of FeII bands, pseudo-lines that are entered into the emission line stack, are created with this command. If the argument in the call is a void string ("") then the file *Fe2Bands.dat* is read by default. This file is included in the data distribution. If a file name appears between a pair of double quotes, as in "*myfile.dat*", it will be used instead.

The routine returns an integer, the number of bands that were read in. These bands are entered into the emission line stack with the label "fe2b" and wavelengths given by the global variable float *FeII_Bands*[n][0]. The FeII bands can be extracted from the predictions, after calling cdDrive, as follows:

```
/* first the FeII bands */
for( n=0; n<nFe2Bands; ++n )
{
    lgOK = cdLine( "Fe2b", (long)FeII_Bands[n][0] , &relative , &absolute );
    if( lgOK<= 0 )
    {
        fprintf(stderr,"did not find
%4s%51i\n","Fe2b", (long)FeII_Bands[n][0]);
        fprintf(ioDATA,"did not find
%4s%51i\n","Fe2b", (long)FeII_Bands[n][0]);
        }
        grid.pred[n+nLines] = log10(MAX2(1e-30,relative) );
}</pre>
```

2.5.8 cdSPEC - get continuous spectra

This routine provides an interface between **Cloudy** and Keith Arnaud's X-Ray spectral analysis program XSPEC. It is called after *cdDrive* has computed a model. Depending on which option it is called with, it will return the incident continuum, the attenuated incident continuum, the reflected continuum, diffuse continuous emission, outward direction diffuse continuous emission, reflected lines, outward lines. All are $4\pi v J_v$ and have units of erg cm⁻² s⁻¹.

All lines and continuum emitted by the cloud assume full coverage of the continuum source. Details are given in *cddrive.h.*

2.6 Example Call as a Subroutine

```
The following is an example of a very simple use of Cloudy as a subroutine.
/*main program that calls cloudy when used as a stand-alone program */
#include <stdio.h>
#include <stdlib.h>
#include <math.h>
#include <string.h>
#include " cddrive.h"
int main( void )
{
       int lgOK ;
       FILE *ioDATA ;
       ioDATA = fopen("d:\\projects\\cloudy\\run2\\run2a.out", "w");
       if( ioDATA == NULL )
               printf(" could not open xspec.out for writing.\n");
              exit(1);
       /* turn off buffering so we results as they happen */
       setbuf( ioDATA , NULL );
       /* initialize the code */
       cdInit();
       cdOutp(ioDATA);
       cdRead( "hden 10.5 ");
       cdRead( "agn 6.00 -1.40 -0.50 -1.0 ");
       cdRead( "phi(h) 23 ");
       cdRead( "stop column density 26.860889 ");
       cdRead( "*constant temper 727,000 ");
       cdRead( "stop zone 3 ");
       cdRead( "punch results \"test.txt\" noclobber ");
       cdRead( "*trace h-like lithium full ");
       cdRead( "no times ");
       cdRead( "punch line intensities column \"run2a.int\" ");
       cdRead( "background z=0 ");
       cdRead( "failures 3 no map ");
       /* actually call the code */
       lgOK = cdDrive();
       fclose( ioDATA);
```

2.7 Storing Grids of Calculations¹

This subsection describes a strategy for creating and analysing a large grid of models.

The **punch results last** command is described in Part I. The line intensities and column densities computed by the code will be written into a file if this

¹ This method is seldom used today and will be removed in a future version of the code.

command is entered. Any post-processing software can read this file, but reading this file is facilitated by routine *cdGett*, described next.

The output produced by the **punch results last** command can be read with routine *cdGett*. *cdGett* has one argument, a pointer to the file containing the results of the **punch results** command. Its return value indicates whether (true) or not (false) the end of file was encountered when the model results were read. If the logical variable is true then the end of file was encountered and no results were read. If it has not been encountered then there may be more models further down the input stream. The next model in the grid will be accessed by the next call to *cdGett*.

Once *cdGett* has been called the routines *cdLine*, *cdIonf*, and *cdColm* routines can be used to obtain emission line intensities, ionization fractions, or column densities from the stored file. A common block, *getpar*, is associated with *cdGett*. This common block contains information concerning the initial parameters of the current model, as specified in the input stream.

This method is seldom used today. The method described next is simpler.

2.8 Computing Grids of Calculations

One or more calculations can be computed within loops, and results extracted for saving. The following examples illustrate producing a series of models with increasing stellar temperature, in which the stellar temperature and the stellar temperature and [OIII]/H β ratio are written to a file.

In the following only one call to *cdLine* is made to get results for a single line. In practice all desired lines are extracted at this stage and stored in a format where it can be read by other software. A call to *cdGetLineList* (page 413 above) provides an easy way to obtain large numbers of lines whose labels are stored in a file. This is the preferred method of calling the code today.

```
/* very simple main program to call cloudy as a stand-alone program */
#include <stdio.h>
#include <stdlib.h>
#include <math.h>
#include <string.h>
/* this must point to cddrive.h */
#include "d:\projects\cloudy\current\source\cddrive.h"
/*int main( int argc, char *argv[] )*/
int main( void )
   /* this will hold images of the command lines */
  char chCard[200];
  double TStar , rel , absol;
  FILE *ioDATA ;
   /* open file for writing some results */
  if( (ioDATA = fopen("1DGrid.out", "w")) == NULL )
     printf(" could not open 1DGrid.out for writing.\n");
     exit(1);
   }
  TStar = 3e4;
  while( TStar < 5e4 )
      /* initialize the code */
     cdInit();
      /* redirect output to the file we opened above */
     cdOutp(ioDATA);
```

```
/* but also say we want to output by pasing 0, for FALSE */
   cdTalk( 0 );
   /* write variables into strings and send string as input file */
  cdRead( "hden 5 ");
   cdRead( "ionization parameter -2 ");
   cdRead( "stop zone 1 ");
   /* this is example of writing a variable into the string then passing
   * the string to cloudy */
   sprintf( chCard , "blackbody, T= %f" , TStar );
   cdRead( chCard );
   /* actually call the code */
   if( cdDrive() )
   {
     printf("Beware: Cloudy returned error condition, so exit with 1.\n");
   }
   /* now increment TStar by 5000K */
  TStar += 5000.;
   /\star get intensity of [OIII] relative to Hbeta - remember cdLine is different
   * from most cd routines since it returns element within line stack, 0 for failure */
   if( !cdLine( "o 3", 5007, &rel, &absol ) <=0 )
   {
     printf("could not find 5007\n");
     exit(1);
   }
   /* now print stellar temperature and 5007/Hbeta ratio */
   fprintf(ioDATA , "%.0f %.2f\n", TStar , rel );
/* things were ok, exit with 0 */
exit(0);
```

}

3 OUTPUT

3.1 Overview

This section defines the output produced by **Cloudy**. Each section begins with a sample of the output described, and then goes on to describe the meaning of the printout in greater detail. The output actually shown is from the Meudon (Pequignot 1986) and Lexington (Ferland et al 1995) meetings Planetary Nebula test case.

3.2 Header Information

Several lines of output echo the input commands and outline some properties of the initial continuum.

3.2.1 Initial Information

	C	loudy 90.00				
**********	*************	***96Apr01****	*********	***********	*******	
*					*	
* title "New" Paris m	*					
* c recompute "standa:	*					
* init file='c84.ini'	*					
* elements read	*					
* sphere					*	
* black body, T=150,0	00K radius = 10)			*	
* hden = 3.4771213					*	
* radius = 17					*	
* abund -1 C-3.523 N-	1. 0-3.222 ne-3	8.824 na=-10 m	g-4.523 al=-1	0	*	
* continue si-4.523 s	4.824 ar-10 ca	a=-10 fe-10 ni	=-10		*	
* plot continuum range	e .1				*	
* punch overview last	70				*	
* c parispn.in					*	
* c Sun					*	
*					*	
***********	*************	*******	*******	***********	*****	
		nemical Compos				
H :1.00E+00 He:1.00E-01 Li:2.04E-0						
Ne:1.50E-04 Na:1.00E-10 Mg:3.00E-0						
K :1.35E-07 Ca:1.00E-10 Sc:1.22E-0				Mn:3.42E-07	Fe:1.00E-10	Co:2.24E-09
	Ni:1.00E-10	Cu:1.87E-08	Zn:4.52E-08			

a1 ---- a0 00

This begins with the version number of *Cloudy*. Major revisions, which have noticeable effects on the emission-line spectrum or which reflect significant improvements in the physics, are denoted by integer increases in the version number, while minor changes increment the revision number by 0.01. In a static version of the code, small changes (usually minor bug fixes) are denoted by letters (i.e., .02a). The following line gives the date this version was created.

All of the input command lines, with the exception of those starting with a #, %, or *, are echoed before the calculation begins, and are saved to be reprinted after the calculation is completed.

The input information is followed by the chemical composition of the gas. The numbers are the number densities of the elements, relative to a hydrogen abundance of unity. Only the active elements are included (those turned off with the **elements off** command are not printed).

3.2.2 Properties of the Continuum

1568Cell Peak3.83E+00	Lo 1.00E-05=0.9108cm	Hi-Con:7.63E+01 Ryd	E(hi):7.35E+06Ryd	E(hi): 100.01 MeV
L(nu>lryd): 37.5396	Average nu:2.935E+00	L(X-ray): 31.4007	L(BalC): 36.1444	Q(Balmer C): 46.9767
Q(1.0-1.8): 47.1613	Q(1.8-4.0): 47.4535	Q(4.0-20): 47.0522	Q(20): 40.7275	Ion pht flx:4.312E+12
L(gam ray): 0.0000	Q(gam ray): 0.0000	L(Infred): 34.4845	Alf(ox): 0.0000	Total lumin: 37.5571
log L/Lsun: 3.9743	Abs bol mg: -5.1858	Abs V mag: 2.4170	Bol cor: -7.6028	nuFnu(Bbet): 34.5868
U(1.0):4.794E-02	U(4.0):9.977E-03	T(En-Den):3.354E+01	T(Comp):1.436E+05	nuJnu(912A):2.908E+01
Occ (FartB):2.354E-10	Occ(H n=6):8.468E-14	Occ(1RVd):1.342E-15	Coc(48):3.707E-17	Occ (Nu-bi):0.00E+00
Occ(FarIR):2.354E-10	Occ(H n=6):8.468E-14	Occ(1Ryd):1.342E-15	Occ(4R):3.707E-17	Occ (Nu-hi):0.000E+00
Tbr(FarIR):3.757E-10	Tbr(H n=6):3.703E-10	Tbr(1Ryd):2.120E-10	Tbr(4R):2.351E-11	Tbr (Nu-hi):0.000E+00

This section gives a synopsis of the incident continuum, evaluated at the illuminated face of the cloud. The first line gives the number of numerical frequency

cells in the continuum, followed by the energy (in Ryd) of the hydrogen-ionizing continuum² with the largest flux density per unit energy interval (f_v). Next are the energies of the low and high energy limit of the continuum, both in Ryd and cm or MeV. The last two numbers are the energies of the high energy limit of the present version of the code, in Ryd and keV.

The second line gives the log of the energy (erg s⁻¹ cm⁻² or erg s⁻¹, depending on whether a flux or luminosity was specified) in the hydrogen ionizing continuum (1 Ryd \leq hv < 100 MeV), and the average energy of the hydrogen ionizing continuum, in Ryd, weighted by photon number;

$$\langle hv \rangle = \frac{\int_{1\,Ryd}^{\infty} 4\,\pi\,J_{\nu}\,d\nu}{\int_{1\,Ryd}^{\infty} 4\,\pi\,J_{\nu}\,/\,hv\,d\nu} \quad .$$
(400)

The log of the energy in the X-ray continuum (20.6 Ryd \leq hv \leq 7676 Ryd), the log of the energy (erg s⁻¹ cm⁻² or erg s⁻¹), and the number of photons (cm⁻² s⁻¹ or s⁻¹) in the Balmer continuum (0.25 Ryd to 1.0 Ryd) is then printed.

The third line gives the log of the number of photons (cm⁻² s⁻¹ or s⁻¹) in four frequency bins (1.0 Ryd \leq hv < 1.807 Ryd, 1.807 Ryd \leq hv < 4.0 Ryd, 4.0 Ryd \leq hv < 20.6 Ryd, and 20.6 Ryd \leq hv < 7676 Ryd). The last number "Ion pht flx" is the flux of hydrogen ionizing photons;

$$\Phi(H) = \frac{Q(H)}{4\pi r^2} \quad \text{cm}^{-2} \text{ s}^{-1} \quad .$$
(401)

In this equation Q(H) is the total number of hydrogen-ionizing photons emitted by the central object (s⁻¹), and r is the separation between the center of the central object and the illuminated face of the cloud. Unlike the majority of the quantities printed in the header, Φ (H) (per unit area) is always printed, never Q(H) (into 4 π sr).

The fourth line of the header gives some information about the low and high energy portions of the incident continuum. The first number is the log of the luminosity in the gamma-ray (100 keV ~ to ~ 100 MeV) continuum. The second number on the line is the log of the number of photons over this energy range. The third number is the log of the luminosity in the continuum between 0.25 Ryd and the lowest energy considered, presently an energy of 1.001×10^{-5} Ryd. All of these entries are either per unit area, or radiated into 4π sr, depending on how the continuum was specified.

The next entry, "Alf(ox)", is the spectral index α_{ox} , defined as in Zamorani et al. (1981), except for the difference in sign convention. This is the spectral index which would describe the continuum between 2 keV (147 Ryd) and 2500Å (0.3645 Ryd) if the continuum could be described as a single power-law, that is,

² The printed number was incorrect in versions 80through 88.01, but had no other effects on computed results.

$$\frac{f_{\nu}(2 \text{ keV})}{f_{\nu}(2500 \text{ Å})} = \left(\frac{\nu_{2 \text{ keV}}}{\nu_{2500 \text{ Å}}}\right)^{\alpha} = 403.3^{\alpha} \quad .$$
(402)

The definition of α_{ox} used here is slightly different from that of Zamorani et al. since implicit negative signs are *never* used by **Cloudy**. Typical AGN have $\alpha_{ox} \sim -1.4$. If no X-rays are present then $\alpha_{ox} = 0$.

The last number on the line is the log of the total energy in the continuum between 1.001×10^{-8} Ryd and 100 MeV. It is given as either erg cm⁻² s⁻¹ or erg s⁻¹, depending on how the continuum was defined. If the continuum is specified per unit area, then this number is 4π times the integrated intensity of the incident continuum. If it is specified as the total luminosity radiated into 4π sr, then the quantity is the luminosity.

The next line is optional, depending on whether the continuum is specified as the total luminosity or photon number radiated into 4π sr, or as an incident surface flux. If the continuum is specified in absolute terms, i.e., the luminosity or photon number radiated into 4π sr, then this optional line is generated. The first quantity is the log of the total luminosity in the continuum, in solar units. The absolute bolometric magnitude, absolute V magnitude, and the bolometric correction, are then given, followed by the log of the continuum specific luminosity (vF_v) at H β (the units of vF_v (H β) are erg s⁻¹).

The next line begins with two ionization parameters. The first is the dimensionless ratio of ionizing photon to hydrogen densities, defined as

$$U \equiv \frac{\Phi(H)}{n_{H}c} \tag{403}$$

where $n_{\rm H}$ is the total hydrogen density. The second number is defined in a similar way, but the numerator is the number of photons with energies greater than 4 Ryd (i.e., helium-ionizing). The third number is the equivalent black body temperature corresponding to the energy density u at the illuminated face of the cloud, from the incident continuum and Stefan's radiation density constant a; $T_u \equiv (L/4\pi r^2 ac)^{1/4}$, and the next quantity is the Compton temperature of the incident radiation field³. The last number on the line is $4\pi v J_v(912 \text{ Å})$, the flux at 912Å (erg cm⁻² s⁻¹). In this equation J_v is the mean intensity of the incident continuum as defined by Mihalas (1978).

The next two lines give some of the incident continuum photon occupation numbers $\eta(v)$, defined as

³For a blackbody radiation field $T_{Compton}$ is roughly 4% lower than the blackbody color temperature T_{color} when the energy density temperature T_u is « T_{color} . Only when $T_u \equiv T_{color}$ does induced Compton heating cause $T_{Compton} \equiv T_{color}$. If $T_u > T_{color}$ then $T_{Compton} > T_{color}$ because of induced Compton heating. All of the relevant physics is included in the Compton temperature printed here.

$$\eta(\nu) = J_{\nu}(\nu) \left(\frac{2h\nu^3}{c^2}\right)^{-1} , \qquad (404)$$

and the incident continuum brightness temperature $T_b(v)$, (K), defined as

$$T_{b}(\nu) \equiv J_{\nu}(\nu) \left(\frac{2k\nu^{2}}{c^{2}}\right)^{-1} , \qquad (405)$$

for five energies. These energies correspond to the lowest frequency considered (presently an energy of 1.001×10^{-5} Ryd); the ionization potential of the n=6 level of hydrogen (1/36 Ryd); an energy of one Rydberg; four Rydbergs, and the high energy limit of the incident continuum (this depends on the continuum shape; the energy is given by the fifth number on the first line of the continuum output).

3.3 Zone Results

#### 1 Te:1	877E+04 F	Iden 3 0001	E+03 Ne+3	512E±03 R.	1 001E+17 1	R-R0.1 2511	2+14 dR+2 4	502E+14 NT	R· 2 Htot	·7 223E-17	T912 9 90	90E+07##
				1.000E+00							H col den	
											T(diffs):	
											Gam 1/tot	
He singlet												
								2.477E-17				1.5055-20
											Force Mul	1 7208.02
								9.512E-01			FOICE Mui	1.7206+03
				9.513E-01			0.0056-05	9.5126-01	3.9076-02	1.01/6-05		
							0 00000.00	H2O+/Otot	0 00000.00	OU. (Ot at 1	0.0000.00	
											02+/0tot:	0.0008.00
											0.000E+00	
								0.000E+00			0.000±+00	A:-12.580
								0.000E+00				
											0.000E+00	
Magnesium 0												
Aluminium 0												
											0.000E+00	
Phosphors 0												
											0.000E+00	
Chlorine 0												
											0.000E+00	
Potassium 0												
											0.000E+00	
											0.000E+00	
											0.000E+00	
											0.000E+00	
											0.000E+00	
											0.000E+00	
											0.000E+00	
											0.000E+00	
											0.000E+00	
											0.000E+00	
Zinc 0	0.000E+00	0.000E+00	1.369E-08	3.726E-06	2.355E-04	3.976E-02	5.138E-01	4.362E-01	9.973E-03	1.011E-05	0.000E+00	0.000E+00

The results of calculations for the first and last zones are always printed. Results for intermediate zones can be printed if desired (see the **print every** command). The following is a line-by-line description of the output produced for each printed zone.

3.3.1 Line 1

The line begins with a series of # characters, to make it easy to locate with an editor. The zone number is the first number, followed by the electron temperature of the zone ("Te"). A lower case u will appear before the "Te" if the temperature solution is possibly thermally unstable (i.e., the derivative of the net cooling with respect to temperature is negative. See the section in Part III on thermal stability problems). The total hydrogen ("Hden") and electron ("Ne") densities (cm⁻³) follow. The next number ("R") is the distance to the center of the zone, from the center of the center of the zone, ("R-R0", or r-r_o), and the thickness of the zone ("dR", or δ r), (all are in cm), follow. The inner edge of the zone is (*r*-*r_o*) - $\delta r/2$ from the illuminated face of the cloud. The line ends with a number indicating how many ionization iterations were needed for this zone to converge (NTR), followed by the total

heating⁴ ("Htot"; photoelectric and otherwise, erg cm⁻³ s⁻¹), and the optical depth between the *illuminated* face of the cloud and the *outer* edge of the zone at the Lyman limit (T912; the number is the *total absorption* optical depth at 912Å, and *not* the hydrogen Lyman limit optical depth).

3.3.2 [Optional] wind parameters

A line describing the velocity and acceleration of the zone is printed if the **wind** option is used. The numbers are the wind velocity at the outer edge of the current zone (km s⁻¹), inward gravitational acceleration (cm s⁻²), total outward radiative acceleration, and the fraction of this acceleration caused by the incident continuum, line driving, and the gradient of the radiation pressure.

3.3.3 [Optional] radiation pressure

If the ratio of line radiation to gas pressure, P(radiation)/P(gas), is greater than 5%, then a line describing the source of the radiation pressure is generated. The line begins with the label **P**(Lines) and continues with the fraction of the total radiation pressure produced that an emission line, the spectroscopic designation of the line, followed by its wavelength in Ångstroms. Up to twenty lines can be printed, although in most cases only Lyα and a few others dominate.

3.3.4 Line 1 - Hydrogen I

The line begins with the abundance of neutral and ionized hydrogen relative to all atomic-ionic hydrogen (i.e., the ratios $H^{o}/(H^{o}+H^{+})$ and $H^{+}/(H^{o}+H^{+})$ where H^{o} is the population in all bound levels of hydrogen. If **print departure coefficients** has been specified then departure coefficients are also printed on the following line. Neutral hydrogen H^{o} is defined to be the total population of atomic hydrogen in all explicitly computed bound levels. Next comes $H^{+o}/Hden$, the ratio of the density of hydrogen in atomic or ionic form (this is indicated by the label "H+o") to the total hydrogen density in all forms (including molecular).

The following five numbers are abundances of the negative hydrogen ion and several molecules (H⁻, H₂, H₂⁺, and HeH⁺) relative to the total hydrogen abundance. The total hydrogen density is usually referred to by the label *hden*, and is the sum H^o + H⁺ + H⁻ + 2H₂ + 2H₂⁺ + 3H₃⁺. Note that, with this definition of the hydrogen density a fully molecular gas will have $n(H_2)/n(H)=0.5$. These molecular abundances are also expressed as departure coefficients if this option is set with the **print departure coefficients** command. The last number on the line is the total hydrogen column density (cm⁻²).

3.3.5 Line 2 - Hydrogen II

The first two numbers are the populations of the H^o 2s and 2p levels relative to the ionized hydrogen density. The next four numbers are populations of levels 3 to 6, again relative to the ionized hydrogen density. All of these populations usually are relative to the ionized hydrogen density, but can also be printed as LTE departure

⁴CLOUDY defines heating as the energy input by the freed photoelectron, or hv - IP, where IP is the ionization potential of the atom or ion, and hv is the energy of the photon. See Osterbrock (1988) for more details.

coefficients if the **print departure coefficients** command is given. The excitation temperature T_{exc} of Ly α , defined as

$$\frac{n(2p) / g(2p)}{n(1s) / g(1s)} = \exp(-h\nu / kT_{exc})$$
(406)

is given. This is followed by the temperature corresponding to the energy density of the attenuated incident continuum ("T(contn)"), and the diffuse continua ("T(diffs)"). This includes all trapped lines and diffuse continuous emission.

3.3.6 Line 3 - Helium

The first three numbers are the total populations of the three ionization stages of helium, relative to the total helium abundance. The population of atomic helium is the sum of the total population in the triplets and singlets, including the population of all explicitly computed levels of each. These populations can also be expressed as departure coefficients if this option is set with the **print departure coefficients** command. The population of He 2³S, relative to the total helium abundance, follows. The Compton heating and cooling rates (both erg cm⁻³ s⁻¹) are next, followed by the gas filling factor. The last number is the fraction of the total hydrogen ionizations that are caused by photoionization from the ground state.

3.3.7 Line 4 - Atomic Helium

The first group are the level populations of the populations of the n=1 to 6 levels of the He^o singlets. Level two is actually resolved into 2s and 2p, but the total population of 2 is printed. The next group consists of populations of the 2s, 2p, and n=3s,p,d levels of the He^o triplets. Both sets of populations are relative to the total helium abundance. Departure coefficients are also printed if requested.

3.3.8 Line 5 - Ionized Helium

The populations of the 2s, 2p, and n=3 to 6 levels are indicated. There are relative to He⁺⁺; departure coefficients are also printed if requested. The ratio of radiation pressure to gas pressure follows.

3.3.9 Optional Grains

If grains are present, then lines giving some properties of the grain populations are printed. Each line gives the results of calculations for a specific type of grain. Normally, a type of graphite and silicate are included when grains are present. There will be one line of output for each grain species. Each line begins with the name of the grain, and an asterisk appears if the species is quantum heated. The remainder of the lines gives the equilibrium temperature of the grain, the potential in volts, the charge, the drift velocity, followed by the gas heating (erg cm⁻³ s⁻¹) due to grain photoionization, and the dimensionless fraction of the total gas heating due to grain photoionization. For quantum heated grains the temperature is the average weighted by T⁴.

3.3.10 Pressure

Some information concerning the pressure is printed. The gas equation of state includes thermal gas pressure, the radiation pressure due to trapped line emission, and the radiation pressure due to absorption of the incident continuum. The first

number is the gas pressure $n_{gas} T_{gas}$ (with units cm⁻³ K), followed by the total pressure (dynes cm⁻²), and followed by the gas pressure ($n_{gas} kT_{gas}$) in dynes cm⁻². The radiation pressure follows. The second to last number is the radiative acceleration (cm s⁻²) at the inner edge of this zone. The radiative acceleration is computed with all continuous scattering and absorption opacities included. The last number is a force multiplier, defined as in Tarter and McKee (1973), and is the ratio of total opacity to electron scattering opacity.

3.3.11 Molecules

A line giving relative abundances of some molecules is printed if there is a significant molecular fraction. All molecular abundances are relative to either the total carbon or total oxygen abundance (this is indicated in the label for each). In order, the molecules are CH, CH⁺, CO, CO⁺, H₂O, and OH.

3.3.12 Li, Be, B

Abundances of each stage of ionization relative to the total gas phase abundance of the element are printed across two lines.

3.3.13 Carbon

The abundances of the seven stages of ionization of carbon relative to the total carbon abundance begin the line. The relative abundance of H_2O^+ and OH^+ (relative to the total oxygen abundance) follows.

3.3.14 Nitrogen

The relative populations of the eight ionization stages of nitrogen are printed first. The relative abundance of O_2 and O_2^+ (relative to the total oxygen abundance) follows.

3.3.15 Oxygen

The oxygen ionization stages are followed by the extra heat added at this zone (erg cm⁻³ s⁻¹); due to cosmic rays, turbulence, etc, and the log of the effective hydrogen recombination coefficient (cm³ s⁻¹).

3.3.16 Fluorine, Neon

The fluorine and neon relative ionization balances are printed across the line.

3.3.17 Remaining Elements

There are too many ionization stages to print across the line. Although all stages with non-trivial abundances are computed, only the highest twelve stages of ionization are printed. The first number is an integer indicating how many stages are "off the page to the left". If the number is 2, then the first printed stage of ionization is twice ionized, i.e., Fe⁺².

3.4 Calculation Stopped Because ...

Calculation stopped because lowest Te reached. Iteration 1 of 1 The geometry is spherical. Non-collisional excitation of [OIII] 4363 reached 2.08% of the total. 1805: Cloud age was not set. I cannot check whether the time-steady assumption is ok. Derivative of net cooling negative and so possibly thermally unstable in 4 zones. Photoionization of He 2TriS reached 17.1% of the total rate out, 10.6% of that was Lya. Grains were not present but might survive in this environment (energy density temperature was 3.35E+01K) The ratio of radiation to gas pressure reached 1.658+01. Caused by Lyman alpha. Line radiation pressure capped by thermalization length.

A series of messages appear after the printout of the last zone.

The first will say why the calculation stopped. In a valid calculation the model will stop because one of the specified stopping criteria specified was met. If no other criteria are specified, then the calculation usually stops when the default lowest temperature of 4000 K is reached. If the code stops because of an unintended reason (i.e., internal errors, or the default limit to the number of zones) then a warning is printed saying that the calculation may have halted prematurely.

Only one stopping criterion message will be printed. The possible messages, and their interpretations, are:

3.4.1 ... because of radiation pressure

The default density law is for a constant density. If constant pressure is specified instead (with the **constant pressure** command), then **Cloudy** will try to keep the total pressure, particle and radiation, constant. The radiation pressure is small at the boundaries of the cloud, so the cloud will be unstable if the ratio of radiation to total pressure exceeds 0.5. The calculation stops, and this message is generated, if this occurs after the first iteration.

3.4.2 ... because lowest EDEN reached.

The calculation can be forced to stop when the electron density (**eden**) falls below a certain value, as set by the **stop eden** command. This can be used to stop the calculation at an ionization front. The default lowest electron density is negative, so this stopping criterion applies only when the command is entered.

3.4.3 ... because low electron fraction.

The calculation can be forced to stop when the ratio of electron to hydrogen densities falls below a certain value, as set by the **stop efrac** command. This can be used to stop the calculation at an ionization front when the hydrogen density there is not known (for instance, in a constant pressure model). The default lowest electron density is negative, so this stopping criterion applies only when the command is entered.

3.4.4 ... because wind veloc too small

The code can perform a wind calculation which includes the outward force due to radiation pressure and the inward force of gravity. The solution is only valid well above the sonic point. This message is printed if the gas is decelerated to below the sonic point.

3.4.5 ... because code returned BUSTED

The calculation stopped because something bad happened. The results are suspect. I would appreciate learning about this - please send the input script and version number.

3.4.6 ... because DRAD small - set DRMIN

The Strömgren radius of the H⁺ zone is estimated at the start of the calculation, and the smallest allowed zone thickness is then set as a very small fraction of this. The calculation will stop if the zone thickness falls below this smallest thickness. This can occur because of any of several logical errors within **Cloudy** (adaptive logic is used to continuously adjust the zone thickness), although it can rarely occur for physical reasons as well. The smallest thickness can be reset to any number with the **set drmin** command, but it should not be necessary to do this. I would appreciate learning about this - please send the input script and version number.

3.4.7 ... because DR small rel to thick.

The depth into the cloud is stored as the double precision variable *depth* and the zone thickness is stored as the double precision variable *drad*. If the zone size becomes too small relative to the depth (*drad/depth* < 10⁻¹⁴) then the depth variable will underflow such that *depth* + *drad* = *depth*. The calculation will stop in this case and give the above reason if this problem prevents the density from being properly evaluated. This is a fundamental numerical problem with no clear solution.

3.4.8 ... because carbon fully molecular.

For mixtures where oxygen is more abundant than carbon the atomic carbon abundance can become vanishingly small when carbon monoxide forms. The matrix inversion routine may have trouble determining the carbon balance under these conditions. As a precaution the current version of the code will stop if the ratio of carbon monoxide to total gas phase carbon exceeds 0.80, the value of the code variable *colimt*. This limit can be reset with the **set colimt** command.

3.4.9 ... because negative mole abundan.

The matrix inversion routine can predict negative abundances of some of the heavy element molecules when the gas becomes predominantly molecular. *Cloudy* is not now designed to handle this situation, but should be well protected against this happening. I would appreciate learning about this occurring- please send the input script and version number.

3.4.10 ... because optical depth reached.

The default value of the largest allowed continuous optical depth is unphysically large, and can be reset with the **stop optical depth** command. The command specifies both the optical depth, and the energy at which it is to be evaluated. All absorption opacity sources included in the calculation contribute to the computed optical depths. If the calculation stops because the largest continuum optical depth is reached, then this line is printed. This line is also printed if the **stop effective column density** command is used to stop the calculation, since this command is actually a form of the **stop optical depth** command.

3.4.11 ... because outer radius reached.

The default outer radius is unphysically large, but can be changed with the **radius** or **stop thickness** commands. If the calculation stops because the outer radius set by one of these commands is reached, then this line is printed.

3.4.12 ... because column dens reached.

The default values of the largest allowed neutral, ionized, and total hydrogen column densities are unphysically large. They can be reset with the commands stop column density, stop neutral column density, or stop ionized column density. This message will be printed if one of these criteria stops the calculation.

3.4.13 ... because lowest Te reached.

The default value of the lowest temperature allowed is 4000 K. This is reasonable when only optical emission lines are of interest. The limit can be changed with the **stop temperature** command. This message is printed if the calculation stops because the lowest temperature is reached.

3.4.14 ... because highest Te reached.

The default value of the highest temperature allowed is 10^{10} K. The limit can be changed with the **stop temperature exceeds** command. This message is printed if the calculation stops because the highest allowed temperature is exceeded.

3.4.15 ... because NZONE reached.

The default condition is for up to 600 zones to be computed. This can be reset with the **stop zone** command. This message is printed if the calculation stops because the limiting number of zones is reached. A warning will be printed at the end of the calculation if it stops because it hits the default limit to the number of zones allowed, presently 600, since this was probably not intended.

The default limit to the number of zones can be increased, while retaining the check that the default limit is not hit, by using the **set nend** command.

3.4.16 ... because line ratio reached.

It is possible to set a limit to the largest value of an emission-line intensity ratio with the **stop line** command. This message is printed if the calculation stops because the largest value of the ratio is reached.

3.4.17 ... because internal error - DRAD.

An internal logical error caused this message to be printed. Send the command lines, and the version number of **Cloudy** to me. My email address is gary@cloud9.pa.uky.edu.

3.4.18 ... because initial conditions out of bounds.

The temperature of the first zone was not within the temperature bounds of the code. This is probably due to the incident continuum not being set properly.

3.4.19 ... because reason not specified.

This is another internal error I would appreciate learning about.

3.5 Geometry


After saying why the calculation stopped, **Cloudy** will say whether the geometry is plane parallel ($\Delta r/r_o < 0.1$), a thick shell ($\Delta r/r_o < 3$), or spherical ($\Delta r/r_o \ge 3$), where r_o is the inner radius and Δr is the thickness of the cloud.

3.6 Warnings, Cautions, Surprises, and Notes

The next, optional, messages fall into four categories: warnings, which begin with W-; cautions, which begin with C-; surprising results, which begin with an explanation mark (!), and notes.

Cloudy checks that its range of validity was not exceeded in the calculation. Warnings are issued to indicate that the program has not treated an important process correctly. For instance, warnings occur if the temperature was high enough for the electrons to be relativistic, if the global heating - cooling balance is off by more than 20%, or if the code stopped for an unintended reason. I would like to hear about warnings, my e-mail address is gary@cloud9.pa.uky.edu. Cautions are less severe, and indicate that **Cloudy** is on thin ice. Examples are when the optical depths in excited states of hydrogen change during the last iteration. Surprises begin with "!" and indicate that, while the physical process has been treated correctly, the result is surprising. An example is when induced Compton heating is more that 5 percent of the total Compton heating. Notes indicate interesting features about the model, such as maser effects in lines or continua, or if the fine structure lines are optically thick. The messages are usually self explanatory.

3.7 Optional Plot



If any of the optional plots are requested with the **plot xxx** command then they will appear next. The quantities plotted are described in the section of HAZY where the plot command is defined.

3.8 Final Printout sphere sphere black body, T=150,000K radius = 10 hden = 3.4771213 radius = 17 abund -1 C-3.523 N-4. O-3.222 ne-3.824 na=-10 mg-4.523 al=-10 plot continuum range .1 punch overview last 70 is actione in Emission Line Spectrum. Constant Density Model. Closed geometry. Iteration 1 of 1. Luminosity (erg/s) emitted by shell with full coverage. general properties..... O 4 1397A -19.618 0.0012 N 3 265A -18.070 0.0417 0.0028 N 3 277A -18 136 0.0358 Al 4 115A -18.802 Al 4 157A -18.512 Al 4 110A -19.172 0.0077 general properties TOTL 4861 -16.690 1.0000 TOTL 1216 -14.856 68.2311 Inci 0 2.563******** TotH 0 -15.017 47.0916 TotC 0 -15.338 22.4813 BFH1 0 -15.382 20.2978 BFHe 0 -15.375 5.1816 UFEr 0 -15.975 5.1816 0 4 1397A -19.618 0 4 1407A -19.246 0 4 1405A -18.737 0 4 1401A -18.585 0 4 789A -15.572 rec 789 -16.972 0 5 630A -15.569 rec 630 -19.299 TOTL 1218 -18.965 0 5 1218 -19.326 0.0028 N 3 277A -18.136 0.0358 0.0151 0.0090 0.0390 0.0033 0.0090 0.0127 13.1211 0.5224 13.1903 0.0238 0.0018 0.0164 0.0011 0.0167 0.0024 0.0025 0.0164 0.0013 0.0124 0.0013 0 -16.544 0 -16.544 0 -19.609 0 -17.355 HEFC 1.3992 0.0023 0.0127 0.0013 0.0023 0.0030 0.3864 0.2595 0.00127 0.0048 0.0037 0.0043 HFF 0 0059 H FF 0 -16.544 1.3992 ComH 0 -19.609 0.0012 CT H 0 -17.355 0.2159 Ca B 1216 -15.292 24.9963 Q(H) 4861 0.154******** Q(H) 1216 1.688******* 0.0017 0.1269 0.0024 0.0105 0.0487 0.2857 0.0025 0.0025 0.0015 0.0176 H-like iso-seq..... CaBo 4861 -16.830 0.7239 CaBo 1640 -15.973 5.2094 0.2346 0.0017 0.0138 0 0013 4365 -19 034 0.0013 Si 3 407A -19.294

The final printout begins by reprinting the input commands. The box surrounding it gives both the version number of *Cloudy* (at the top) and the log of the ionization parameter (the ratio of ionizing photon to hydrogen densities) at the bottom.

The line following the box summarizes some properties of the model and output. The first part of the line indicates whether the energy in the emission lines is given as the luminosity radiated by a spherical shell covering Ω sr (erg s⁻¹; $\Omega / 4\pi$ is the covering factor) or the intensity produced by a unit area of gas (erg s⁻¹ cm⁻²). Which of the two choices is printed is determined by whether the luminosity of the continuum was specified as the luminosity radiated by the central object into 4π sr or the intensity (4π) of the incident continuum (erg cm⁻² s⁻¹) at the illuminated face of the cloud. If the model is spherical and the incident continuum specified per unit area, then the emergent emission-line spectrum will be per unit area in units of the inner radius r_o (that is, the total line luminosity radiated by a shell covering 4π sr will be the listed intensity $4\pi j$ times $4\pi r_o^2$). The second part of this line indicates the density structure of the model (i.e., wind, constant density, constant pressure, constant gas pressure, power-law density distribution, etc). The next section tells whether the geometry was open or closed. The last part indicates which iteration this is.

The computed emission-line spectrum follows. Emission lines are divided into two groups. The first includes the effects of grain scattering and absorption, and is indicated by the header Emergent Line Intensities. This first group is only printed if grains are present and the geometry is open (i.e., **sphere** not set). The intensities are the *total* intensities observed from the illuminated face, including both absorption and scattering by grains.

The second, larger, group of lines is always printed, is usually the intrinsic intensity of the lines, and does not include the reddening effects of internal grains due to the photon's passage out of the nebula. This second group usually gives the

total intrinsic intensity of the lines. Although reddening effects of internal (or external) dust are not taken into account, photon destruction by background opacity sources during the transfer process is. This predicted spectrum should be compared with the reddening-corrected observed spectrum.

The spectrum is sorted into four large groups of columns, with each large group sub-divided into four smaller sub-columns. Within each column lines are organized by common origin with a comment beginning the section. The first sub-column is either the spectroscopic designation of the ion producing the line or an indication of how the line is formed. The second sub-column is the line wavelength, with a 0 to indicate a continuum. The third sub-column is the log of the power in the line, in the units given in the header (4π sr or cm⁻²). The last sub-column is the intensity of the line relative to the reference line, usually H β , unless this is reset with the **normalize** command.

In the current version of the code the line wavelength is an integer. This is to allow a precise match with a wavelength specified during a search. An integer format has obvious limitations in specifying fractions. To overcome this, the format of the wavelength changes with wavelength, in an attempt to have several significant figures. For wavelengths longward of 100µm the wavelength is given in µm. For wavelengths longward of one micron the wavelength is given in tenths of microns. For wavelengths between 10Å and 100Å the wavelength is given in tenths of Å, and in hundredths of Å below this⁵. (The logic deciding what format should be printed is given in routine *iWavLen*.) The units of the wavelength are indicated by the letter immediately following the wavelength. These are currently either A or m indicating Ångstroms or microns. An integer may follow this letter. This is the number of places that should have occurred after the missing decimal point. For instance, the label **Ne 3 156m1** indicates that the line is [Ne III] 15.6 μ .

The following sections give overviews of the general treatment of line formation. The section beginning on page 442 of this document gives more details about the predictions, and should be consulted for precise definitions.

Blocks of lines. The emission lines are organized according to their origin with the main set of lines. Each section begins with a comment that specifies its origin.

General properties. The first two entries, TOTL 4861 and TOTL 1216, are the total intensities of H β and Ly α , as predicted by the multi-level H atom. These intensities are the results of calculations that include all collisional, radiative, and optical depth effects. The total energy in the incident continuum is Inci 0. photoelectric heating, due to photoionization of hydrogen, is given by TotH. BFHx is heating due to photoionization of excited state hydrogen. The entries BFHe and TotM are the heating due to helium and metal photoionization. GrGH and GrGC are the amounts of heating and cooling provided by grains. The entry Q(H) 4861 is the intensity of H β predicted from the total number of ionizing photons, Q(H), assuming that each hydrogen-ionizing photon produces one hydrogen atom recombination

⁵ Before version 94.01 of the code the wavelength was given in tenths of Å below 100Å. The **set oldwavelength** command will cause the code to continue to do this.

(see, for example, Osterbrock 1989). Q(H) 1216 indicates the Ly α intensity produced if each hydrogen ionizing photon results in one Ly α photon in the high density limit (i.e., no two-photon emission).

H-like iso-seq. The entries starting with Ca B are the Case B intensities computed from the actual model ionization and temperature structure, but assuming that H β emits with its case B emissivity. Next the predicted intensities of all lines of the hydrogenic iso-electronic sequence are given. The lines have labels that identify the species and stage of ionization, such as H 1, He 2, Li 3, C 6, etc. The entries with a wavelength of zero are the total intensities of the 2s-1s two photon emission.

Continua... These given intensities of various continua. These are either the total integrated continuum or the product vF_v at certain energies.

Molecules. H2 1 2 is the intensity of the H₂ lines near 2 μ m, and H2 d is the cooling due to collisional dissociation of H₂. H-FB and H-FF are the free-bound and free-free continua of the H⁻ ion. H2+ and HEH+ are the cooling due to formation of H₂⁺ and HeH⁺.

He iso-sequence Atoms and ions of the helium-like iso-electronic sequence are treated as multi-level atoms. Each species and stage of ionization are specified by labels like He 1, Li 2, C 5, etc. A wavelength of zero indicates the two-photon continuum.

level 1 lines A few notes on deciphering the heavy element lines follow. In cases where the notation is unclear a careful examination of subroutine *lines* (which enters the line fluxes into the arrays) or *coolr* (which actually calculates the line intensities) should clarify the meaning. In general, the line wavelengths are given in Å, although the infrared fine structure lines are an exception. Often these IR lines have their wavelength given in microns (for instance, [O III] λ 88 µm), but sometimes it is given in microns or tenths of microns (this is because the wavelengths are integers). Notes on specific iso-electronic sequences follow.

Li-sequence. Examples include C IV λ 1549, O VI λ 1034, Mg II λ 2798, etc. A three level atom, with full treatment of optical depths and collisional excitation, is used. The "TOTL" intensity is the sum of both lines in the doublet, and is followed by the individual intensities of each member.

Be-sequence. Examples include C III] λ 1909, O V] λ 1215, Si III] 1895, etc. A four level atom, solving for populations of the individual ³P_j states, is used. The first printed intensity is the total intensity of the multiplet (both j=0,1 decays), and this is followed by the intensities of individual lines. The intensity of the permitted ¹P_o - ¹S transition is also calculated. Optical depth and collisional effects on both the permitted and intercombination lines are included.

B-sequence. Examples include C II and O IV. The ground term is treated as a two level atom, with optical depth and collisional effects included. The ^{4}P - $^{2}P_{o}$ lines are also predicted with a two level atom. The intensity printed is the total intensity of the multiplet.

³P- ground term. Examples include [O III] and [O I]. The infrared fine structure lines are computed with full treatment of collisional and optical depth effects. A comment is printed at the end of the model if these lines mase or become optically thick. The populations of ¹D and ¹S are computed with a three-level atom. The intensity of the ¹D - ³P transition is only that of the individual line (i.e. 5007), not the doublet.

⁴S^o - ground term. Examples include [O II] and [S II]. They are treated as a five level atom. Intensities of all individual lines, as well as co-added multiplets, are given.

Recombination ... These are a set of recombination lines that are predicted assuming that they are optically thin.

Level 2 lines . .. These are resonance lines that use Opacity Project wavelengths, which are generally accurate to about 10%, and have g-bar collision strengths, which are not very accurate at all.

A list of emission lines with negative intensities may follow the main block of lines. These are lines which heat rather than cool the gas (heating is negative cooling). This is not a problem, but occurs if the line de-excitation rate exceeds the line excitation rate. The most common reason for this to occur is if the line is radiatively excited but collisionally de-excited.

3.8.2 Thermal balance

Cooling: 0 3 5007:0.245 0 3 4959:0.082 Heating: BFH1 0:0.720 BFHe 0:0.233 IONIZE PARMET: U(1-) -1.3193 U(4-): -2.0010 U(sp): -2.51 Q(ion): 43.458 L(ion): 33.712 Q(low): 49.69 P(low) 37.341 ENERGY BUDGET: Heat: 37.222 Coolg: 37.222 Error: 0.2% Rec Lin: 37.064 WorkF: 37.437 F-F H 21.885 RadBetaMax:1.65E+01

Cooling: This line indicates the fraction of the total cooling (defined here as in Osterbrock 1989; that is, the energy of the freed photoelectron) carried by the indicated emission lines. The designation of the line is given as in the emission-line spectrum, and this is followed by the ratio of the energy in the line to the total cooling. This is an important indication of the fundamental power-losses governing conditions in the model. The labels used are the same as those in the line array.

Heating: This line indicates the fraction of the total heating produced by various processes. The labels used are the same as those in the line array.

IONIZE PARMET The line begins with the log of the H "U(1-)" and He⁺ "U(4-)" ionization parameters defined in the header. The third number "U(sp)" is the log of a spherical ionization parameter often used in spherical geometries, such as H II regions or planetary nebulae. It is defined as

$$U_{sph} = \frac{Q(H)}{4\pi R_s^2 n_H c} \tag{407}$$

where R_s is the Strömgren radius, defined as the point where the hydrogen neutral fraction falls to $H^0/H_{tot} = 0.5$. If no ionization front is present, then U_{sph} is evaluated at the outer edge of the computed structure. The next two numbers are the log of the number of hydrogen ionizing photons (hv ≥ 1 Ryd) exiting the nebula "Q(ion)", and the log of the energy in this ionizing continuum "L(ion)". The last two numbers are the equivalent quantities, for non-ionizing (hv < 1 Ryd) radiation. These are either

per unit area or by a shell covering 4π sr. These have been corrected for the r⁻² dilution if per unit area, and so are directly comparable with the numbers given at the start of the calculation.

ENERGY BUDGET This line gives an indication of the energy budget of the nebula. The first number "Heat" is the log of the total heating (in ergs s⁻¹, but again either into 4π sr or cm⁻²). The second number "Coolg" is the log of the total cooling, in the same units. Cooling, as defined in Osterbrock (1989), is the total energy in collisionally excited lines and part of the recombination energy, but *does not* include recombination lines. The percentage error in the heating-cooling match "Error" follows. The next number "Rec Lin" is the log of the total luminosity in recombination lines. The number indicated by "WorkF" is an indication of the work function (that is, the log of the energy needed to remove bound electrons from the atom or ion) of the cloud. The work function and the total cooling do not add up to the total energy absorbed from the incident continuum because some recombination lines of helium and heavy elements contribute to both. The next number "F-F H" is the log of the amount of energy deposited by free-free heating, and the last number "RadBetaMax" is the largest value of the ratio of radiation to gas pressures which occurred in the calculation.

3.8.3 Column densities, et al.

Column density	H12:9.024E+20			H-: 1.445E+12			He H+:4.187E+12
	CH:0.000E+00	CH+:0.000E+00	OH:0.000E+00	OH+: 0.000E+00	O2: 0.000E+00	C2:0.000E+00	CO:0.000E+00
	CO+:0.000E+00	H2O:0.000E+00H2	2O+:0.000E+00	02+: 0.000E+00	C2+: 0.000E+00	H3+:0.000E+00	H3O+:0.000E+00
	CH2+:0.000E+00	CH2:0.000E+00 0	CH3:0.000E+00M	oH2O 0.000E+00			
Col(Heff):	6.286E+20	snd travl time	2.34E+11 sec	NeN+dl: 2.98E+24	Te-low:3.81E+03	Te-hi1.88E+04	
He/Ha:9.82E-02	= 0.98*true	Lthin:1.00E+30	iter/zn:2.016	Hlu/zn:6.56E+00			
<a>:0.00E+00	erdeFe0.0E+00	Tcompt8.11E+07	Tthr6.76E+09	<tden>: 1.33E+04</tden>	<dens>:7.11E-21</dens>	<mol>:6.47E-01</mol>	
Mean Jeans	l(cm)7.67E+19	M(sun)8.46E+05	smallest:	len(cm):4.10E+19	M(sun):1.29E+05	Alf(ox-tran):	0.0000

Column density This line lists the column densities (cm^{-2}) of some ions and molecules. The first number "H12" is the total hydrogen column density (both H^o and H⁺). The following two numbers are the column densities in H⁺ and H^o only. The last four numbers are column densities in four ion - molecules (H-, H₂, H₂⁺, and HeH⁺).

The next series of three lines give column densities in various molecules.

Col (Heff) The effective column density "Col(Heff)", as defined in the section on the **stop effective column density** command, is printed. This is followed by "snd travl time", the sound travel time across the nebula in seconds. Constant pressure is only valid if the cloud is static for times considerably longer than this. The third number "NeN+dl" is the emission measure, the integral over radius of the product $n_e n_p f(r) dr$, where f(r) is the filling factor. The last two numbers are the lowest "Te-low" and highest "Te-hi" electron temperatures found in the computed structure.

He/Ha This line gives some quantities deduced from the predicted emission-line spectrum. The first (He/Ha) number is the apparent helium abundance He/H, measured from the emission-line intensities using techniques similar to those described in Osterbrock (1989);

$$\left(\frac{\text{He}}{\text{H}}\right)_{apparent} = \frac{0.739 \times I(5876) + 0.078 \times I(4686)}{I(H\beta)} \quad .$$
(408)

The intensity of both H β and HeI λ 5876 are the total predicted intensities, and includes contributions from collisional excitation and radiative transfer effects. The intensity of HeII λ 4686 is taken from Case B results, which are better than those of the model atom at low densities. The second number (i.e., 1.07*true), is the ratio of this deduced abundance to the true abundance. This provides a simple way to check whether ionization correction factors, or other effects, would upset the measurement of the helium abundance of the model nebula. This is followed by the longest wavelength in centimeters "Lthin" at which the nebula is optically thin. Generally the largest FIR opacity source is brems, and the number will be 10³⁰ if the nebula is optically thin across the IR. The last two quantities are related to the average number of iterations needed to converge each zone.

<a> The mean radiative acceleration (cm s⁻²) is printed if the geometry is a wind model and zero otherwise. This is followed by some time scales. The first "erdeFe" is the time scale, in seconds, to photoerode Fe (Boyd and Ferland 1987; this number is 0s if the γ -ray flux is zero). The next two are the Compton equilibrium timescale "Tcompt", and the thermal cooling timescale "Tthr". Both are in seconds. The density (gm cm⁻³) weighted mean temperature "<Tden>", radius weighted mean density "<dens>" (gm cm⁻³), and mean molecular weight "<Mol>" follow.

Mean Jeans This line gives the mean Jeans length "l(cm)" (cm) and Jeans mass "M(sun)" (in solar units), followed by the smallest Jeans length "smallest len(cm)" and the smallest Jeans mass "M(sun)" which occurred in the calculation. The last quantity "Alf(ox-tran)" is the spectral index $\alpha_{ox'}$ defined as in the header, but for the transmitted continuum (attenuated incident continuum plus emitted continuum produced by the cloud).

H and He atoms ,This line gives the number of levels of the model hydrogen atom, the "topoff" level, above which the remainder of the recombination coefficient is added, the type of topping off used for this calculation, and the number of levels used for the helium singlets and ion. The last number on the line is the execution time in seconds.

3.8.4 Averaged Quantities

 Averaged Quantities

 Te
 Te (Ne)
 Te (NeHe)
 Te (NeHe)
 Te (NeO+)
 Te (NeO+)
 NH
 Ne (02+)
 Ne (Np)

 Radius:
 1.334E+04
 1.358E+04
 1.363E+04
 1.363E+04
 1.109E+04
 1.139E+04
 3.000E+03
 3.339E+03
 3.429E+03

 Volume:
 1.179E+04
 1.207E+04
 1.214E+04
 1.100E+04
 1.115E+04
 1.117E+04
 3.000E+03
 3.305E+03
 3.294E+03

 Peimbert T(0IIIr)1.15E+04 T(Bac)1.21E+04 T(Hth)1.14E+04 t2(Hstrc) 2.72E-02 T(03-BAC)1.15E+04 t2(03-BC)
 1.81E-03 t2(03str) 7.36E-03

This begins with several temperature and density averages, over either radius or volume. The volume averages are only printed if the **sphere** command is entered. The quantity which is printed is indicated at the top of each column. The averaged quantity is the first part of the label, and the weighting used is indicated by the quantity in parenthesis. For instance, **Te** (**NeO2+**) is the electron temperature averaged with respect to the product of the electron and O²⁺ densities.

Peimbert This series of quantities deal with temperature fluctuations (t^2 , Peimbert 1967). The code attempts to analyze the predicted emission line and continuum spectrum using the same steps that Manuel outlined in this paper. The code does not attempt to correct the predicted emission line intensities for collisional suppression or reddening, so this line is only printed if the density is below the density set with

the **set tsqden** command - the default is 10⁷ cm⁻³. This code does not attempt to deredden the spectrum: a caution is printed if grains are present.

The nature of temperature fluctuations is, in my option, the biggest open question in nebular astrophysics. Theory (*Cloudy* too) predicts that they should be very small, because of the steep dependence of the cooling function on the temperature, while some observations indicate a very large value of t^2 (see Liu et al. 1995, and Kingdon and Ferland 1995 for a discussion). If something is missing from our current understanding of the energy source of photoionized nebulae then the entire nebular abundance scale (for both the Milky Way and the extragalactic nebulae) is in error by as much as 0.5 dex.

Two fundamentally different t^2 s enter here - the "structural" t^2 and the observational t^2 . The structural value comes from the computed ionization and thermal structure of the nebula, while the observational value comes from an analysis of the predicted emission line spectrum following the methods outlined in Peimbert's 1967 paper.

The structural t^2 for the H⁺ ion is defined as

$$t^{2}(H^{+}) = \left\langle \left[\frac{T(r) - \langle T \rangle}{\langle T \rangle}\right]^{2} \right\rangle = \frac{\int \left[T(r) - \langle T \rangle\right]^{2} n_{e} n_{H^{+}} f(r) dV}{\langle T \rangle^{2} \int n_{e} n_{H^{+}} f(r) dV}$$
(409)

where <T> is the density-volume weighted mean temperature

$$\langle T \rangle = \frac{\int T(r) n_e n_{H^+} f(r) \, dV}{\int n_e n_{H^+} f(r) \, dV}.$$
(410)

This quantity is given in the averaged quantities block as the column "Te(NeNp)".

The observational t^2 - related quantities are the following: "T(OIIIr)" is the electron temperature indicated by the predicted [OIII] 5007/4363 ratio in the low density limit. This number is meaningless for densities near or above the critical density of these lines. "T(Bac)" is the hydrogen temperature resulting from the predicted Balmer jump and H β . "T(Hth)" is the same but for optically thin Balmer continuum and case B H β emission. "t2(Hstrc)" is the structural HII t². The entries "T(O3-BAC)" and t2(O3-BC)" are the mean temperature and t² resulting from the standard analysis of the [OIII] and HI spectra (Peimbert 1967). Finally "t2(O3str)" is the structural t^2 over the O²⁺ zone. Only the structural t²s are meaningful for high densities. This section was developed in association with Jim Kingdon, and Kingdon and Ferland (1995) provide more details.

3.8.5 Grains

Average grain Properties: Gra-Ori Sil-Ori <Tdust>: 1.540E+02 1.218E+02 <Vel D>: 3.635E+04 2.891E+04 <Pot D>: 1.337E+00 1.419E+00

The next lines give some information concerning grains if these were included in the calculation. These lines give the mean temperature, drift velocity, and potential, for all of the grain populations included in the calculation. An asterisk will appear to the right of the name of any species with quantum heating included. In this case the mean temperature is weighted by T⁴.

Contin Optical Depths: COMP: 6.80E-04 H-: 5.59E-05 R(1300): 2.14E-04 H2+ 1.50E-06 HeTri:6.09E-04 Pfa:3.40E-04 Pa:3.40E-04 Ba:3.65E-04 Hb:3.60E-04 La:1.29E-01 1r:4.995E+07 1.8:1.19E+07 4.:1.738E+06 Line Optical Depths: 10830: 1.23E+02 3889: 5.24E+00 5876: 3.29E-06 7065: 1.82E-06 2.06m 2.54E-03

3.8.6 Continuum optical depths

Contin Optical Depths The first two lines give the continuum optical depths at various energies. These are the total optical depths, including the correction for stimulated emission, and will be negative if maser action occurs. These include grain opacity if grains are present. The labels, and their interpretation, are as follows. COMP is Thomson scattering. H- is the negative hydrogen ion at maximum cross section. R(1300) is Rayleigh scattering at 1300Å, H₂⁺ is the molecular hydrogen ion. HeTri is the helium triplet at threshold. The next line gives total continuous optical depths at the energies of various hydrogen and helium ionization edges and lines. These are the Pfund α , Paschen α , Balmer α and β , Ly α , and the ionization edges of hydrogen, atomic helium, and the helium ion.

Heavy element line optical depths are printed also if the **print line optical depths** command is entered.

Old hydro optical depths: Old H Lines:	1 9.99E+07 2-1 9.96E+19	2 1.00E-20 3-2 3.34E-02	3 1.00E-20 4-3 3.33E-04	4 1.00E-20	5 1.00E-20	6 1.00E-20 7-6 2.56E-07	7 1.00E-20 8-7 2.56E-08
New hydro optical depths:	1 4.99E+07	2 1.46E-05	3 2.82E-05	4 6.80E-06	5 1.25E-12	6 3.05E-12	7 6.49E-12
New H Lines: Old He Is optical depths:	2-1 2.02E+06 1 2.38E+07	3-2 9.37E-03 2 1.00E-20	4-3 1.98E-09 3 1.00E-20	5-4 6.50E-10 4 1.00E-20	6-5-5.01E-09 5 1.00E-20	7-6-2.94E-08 6 1.00E-20	8-7-1.81E-07 7 1.00E-20
Old He Is Lines:		3-2 1.00E-20		5-4 1.00E-20	6-5 1.00E-20	7-6 1.00E-20	8-7 1.00E-20
New He Is optical depths:	1 1.19E+07	2 1.46E-05	3 2.83E-05	4 6.06E-06	5 1.41E-12	6 3.33E-12	7 6.93E-12
New He Is Lines: Old He II optical depths:	2-1 6.15E+04 1 3.41E+06	3-2 1.12E-03 2 9.75E+07	4-3 3.50E-11 3 1.00E-20	5-4-2.59E-11 4 1.00E-20	6-5-1.25E-10 5 1.00E-20	7-6 5.98E-09 6 1.00E-20	8-7 2.36E-08 7 1.00E-20
Old He II Lines:	2-1 9.96E+19	3-2 1.00E-20	4-3 1.00E-20	5-4 1.00E-20	6-5 1.00E-20	7-6 1.00E-20	8-7 1.00E-20
New HE II optical depths: New He II Lines:	1 1.70E+06 2-1 1.13E+06	2 4.88E+07 3-2 9.14E-05	3 1.89E-04 4-3 1.36E-11	4 1.39E-05 5-4-4.73E-11	5 2.16E-05 6-5-2.17E-10	6 2.75E-05 7-6 5.52E-09	7 2.72E-05 8-7 2.11E-08
New He II Lines:	2-1 1.13E+06	3-2 9.14E-05	4-3 1.36E-11	5-4-4.73E-11	v-s-∠.1/E-10	/-0 5.52E-09	8-/ 2.11E-08

Hydrogen and helium optical depths in continua and $n \rightarrow n-1\alpha$ transitions follow. The first two lines are the optical depths assumed at the start of the present iteration, and the second pair of lines gives the newly computed total optical depths. Negative optical depths indicate maser action. For each of the pairs of lines, the first line is the optical depth at thresholds of the first seven levels of hydrogen. The second line gives the optical depths in the first seven of the $n \rightarrow n-1\alpha$ transitions of hydrogen or helium.

3.8.7 Mean ionization

Hydrogen	-1.117 -0.035			L	og10 Mea	an Ionis	sation	(over v	olume)						
	-1.454 -0.163	-0 556			- 5										
	-3.801 -0.106		60												
	-4.065 -0.553														
	-5.005 -0.770														
	-3.439 -0.851														
	-1.448 -0.791														
	-1.150 -0.933				-3 971										
	-2.080 -1.428														
	-2.102 -1.352														
	-3.859 -0.950					-0 180									
	-3.156 -0.657					5.105									
	-4.336 -0.630														
	-5.022 -0.547														
	-3.631 -0.799														
	-3.730 -0.780					7 002									
	-1.561 -0.913														
	-1.634 -1.371						E 240								
	-2.619 -1.216														
	-4.301 -1.854														
	-4.229 -1.170														
	-4.559 -1.198														
	-3.087 -1.296														
	-4.914 -1.336														
	-3.265 -1.348 -3.810 -0.849														
	-3.991 -1.413							7 150							
	-3.628 -0.848														
	-4.106 -1.406														
Zinc	-4.254 -1.360														
ZINC	1 2	-0.980 -0.2	81 -1.560 5	-0.930	-0.810	-1.558	-3.630	10	11	12	13	14	15	16	17
	1 2	3 4	5	6	/	8	9	10	11	12	13	14	15	10	1/
Hydrogen	-1.450 -0.016			T.	ogio Mei	an Ionis	ation	lover r	adine)						
	-1.797 -0.352	-0.268		1	ogio nea	101112	Sacion	(Over I)	aurus/						
	-4.049 -0.262		96												
	-4.379 -0.801														
Boron	-5.337 -1.073														
	-3.754 -1.162														
	-1.797 -1.101														
wrerogen	1.757 -1.101	0.514 -0.4	2.5 0.770	1.240											

Oxygen	-1.495	-1.233	-0.362	-0.588	-0.715	-1.621	-3.317			
Fluorine	-2.435	-1.754	-0.311	-0.652	-0.601	-1.782	-4.937			
Neon	-2.457	-1.663	-0.308	-0.595	-0.666	-1.909	-4.941			
Sodium	-4.193	-1.236	-0.361	-0.540	-0.726	-1.534	-3.649	-8.348		
Magnesium	-3.468	-0.945	-0.363	-0.529	-0.892	-1.549	-3.862			
Aluminium	-4.652	-0.926	-1.072	-0.176	-0.935	-1.866	-4.142			
Silicon	-5.342	-0.835	-0.710	-0.647	-0.380	-1.800	-4.425			
Phosphorus	-3.932	-1.106	-0.713	-0.413	-0.747	-0.789	-3.753			
Sulphur	-4.040	-1.090	-0.587	-0.567	-0.758	-0.788	-1.293	-6.378		
Chlorine	-1.908	-1.216	-0.588	-0.559	-0.864	-0.705	-1.257	-2.561		
Argon	-1.986	-1.694	-0.479	-0.634	-1.071	-0.702	-0.929	-2.364	-4.560	
Potassium	-2.960	-1.548	-0.469	-0.627	-0.992	-0.714	-1.038	-2.070	-4.337	
Calcium	-4.623	-2.157	-0.583	-0.443	-1.054	-0.728	-1.060	-2.053	-3.943	
Scandium	-4.564	-1.502	-0.959	-0.300	-0.860	-0.782	-1.283	-2.758	-5.012	
Titanium	-4.901	-1.542	-1.268	-0.583	-0.366	-0.730	-1.418	-2.834	-5.325	
Vanadium	-3.427	-1.643	-1.349	-0.650	-0.519	-0.436	-1.415	-3.062	-5.594	
Chromium	-5.259	-1.684	-1.383	-0.660	-0.590	-0.540	-0.766	-2.734	-5.520	
Manganese	-3.608	-1.696	-1.366	-0.591	-0.701	-0.574	-0.702	-1.814	-4.853	
Iron	-4.143	-1.179	-1.174	-0.500	-1.207	-0.676	-0.617	-1.455	-3.271	
Cobalt	-4.339	-1.762	-1.440	-0.705	-0.782	-0.711	-0.497	-1.159	-2.992	-6.406
Nickel	-3.966	-1.182	-1.590	-0.493	-1.586	-0.724	-0.525	-1.140	-2.852	-6.109
Copper	-4.452	-1.754	-1.299	-0.467	-1.602	-0.839	-0.451	-1.181	-2.880	-6.154
Zinc	-4.602	-1.706	-1.258	-0.479	-1.573	-0.784	-0.499	-1.077	-3.021	-6.265

The two large blocks of output give the mean ionization, averaged over volume, and over radius. The numbers printed are the log of the mean ionization fraction in the various stages. The volume average ionization fraction for ion *i* of element *a* is given by

$$\left\langle \frac{n_a^i}{n_a} \right\rangle_{vol} = \frac{\int n_a^i f(r) dV}{\int n_a f(r) dV}.$$
(411)

and the radius average by

$$\left\langle \frac{n_a^i}{n_a} \right\rangle_{rad} = \frac{\int n_a^i f(r) \, dr}{\int n_a f(r) \, dr} \,. \tag{412}$$

3.8.8 Continuum

This is only printed if the **print continuum** command is included. Then the following tables, all related to the transmitted continuum, will the printed.

X-Ray Continuum. The next line gives the photon fluxes (cm⁻² s⁻¹) in various X-ray bands, if the continuum extends to X-ray energies. The units of the energy bands are keV. The numbers are the numbers of photons exiting the cloud, integrated over the energy bands. This is the net continuum, that is, the incident continuum, less attenuation, with diffuse re-emission from the cloud added on. This is only printed if the **print continuum** command is entered.

Normalized Continuum. This block is a set of ordered pairs giving the emergent Balmer continuum, relative to the continuum which entered the cloud. The first number of each pair is the frequency in Rydbergs. The second is the ratio of the emergent continuum to the incident continuum (i.e., that which went into the cloud). In the absence of optical depth or diffuse emission effects, this block will be equal to 1.000 throughout. This is only printed if the **print continuum** command is entered.

Emergent Continuum. This block gives ordered pairs of energy (in Rydbergs) and the emergent continuum. It is expressed as photon fluxes (phot Ryd⁻¹ cm⁻²) corrected for r⁻² dilution, so as to be directly comparable with the continuum which went into the cloud. This is only printed if the **print continuum** command is entered.

4 OBSERVED QUANTITIES

4.1 Overview

This section describes how to convert the quantities actually used or predicted by *Cloudy* into commonly observed ones.

4.2 Incident and Diffuse Continua

The emission line printout gives the intensity of the incident continuum (λF_{λ} or νF_{ν}) at 4860 and 1215 Å. These appear with the label **Inci** followed by the wavelength. The entire incident continuum can be obtained with the output of the **punch continuum** command.

The diffuse continuum, that emitted by the cloud, is not normally included in the line output. The **print diffuse continuum** command will add the total emitted continuum to the emission line list. These are in units λF_{λ} or vF_{v} at the indicated wavelengths and have the label **nFnu**. The entry with the label *nTnu* is the sum of the reflected plus attenuated incident continuum. The inward total emission and the reflected incident continua will be printed if this command appears together with the **print line inward** command. Two contributors to the inward emission are predicted. That labeled *InwT* is the total inwardly emitted continuum, and includes both diffuse emission and the back-scattered incident continuum. The component labeled *InwC* is the back-scattered incident continuum alone.

4.3 Line Equivalent Widths

The equivalent width of an emission or absorption line is defined as the number of Ångstroms of the continuum that is equivalent to the energy in the line. It can be defined as

$$W_{\lambda} = \int \frac{F_{\lambda}^{c} - F_{\lambda}^{l}}{F_{\lambda}^{c}} d\lambda \approx -\lambda \frac{F_{line}}{\lambda F_{\lambda}^{c}}$$
(413)

where the fluxes are in the interpolated continuum (F_{λ}^{c}) and the integrated line (F_{line}). By this convention the equivalent width of an emission line is negative.

The code predicts the integrated fluxes of all lines. It also predicts the product λF_{λ}^{c} for the incident continuum at a few wavelengths. These are given the label **Inci** and the wavelength where it is evaluated follows. The entry **Inci 4860** is the intensity of the incident continuum at a wavelength near H β . The units of this incident continuum are either erg cm⁻² s⁻¹ or erg s⁻¹ depending on whether the incident continuum was specified as a flux or luminosity. The fluxes of lines and these continuum points can be read from the output, or obtained by software calling the *cdLine* routine. The continuum flux at any wavelength can be obtained with the **punch continuum** command. If the line intensity is given by F_{line} and the continuum intensity λF_{λ}^{c} , then the equivalent width of a line relative to the continuum where λF_{λ}^{c} is specified will be given by the last term in equation 413.

A covering factor will complicate this slightly. (Covering factors are defined in the section *Definitions* in Part I of this document.) If luminosities are predicted then partial coverage of the source is taken into account with the **covering factor** command, and the luminosities are correct for this coverage. The ratio of line to continuum given in equation 413 will represent what is observed. If fluxes are specified instead then the line flux is given per unit area of cloud, no matter what covering factor is specified. In this second case the ratio in equation 413 must be scaled by the covering factor.

4.4 Emission Line Asymmetries

The inward fraction of the total emission of each line is always predicted by the code, but not normally printed out. Many lines are significantly inwardly beamed, and this can lead to emission line asymmetries if the envelope is expanding. The inward part of the lines will be printed if the **print line inward** command is entered. The effects of this line beaming is very geometry dependent.

4.5 Line to Continuum Contrast

The code has several **punch** commands that will produce ancillary files containing

the predicted line and continuum spectra. There is an ambiguity in how strong the lines should appear to be relative to the continuum. This is described in Part I of this document where the **punch continuum** and **set PunchLWidth** commands are introduced.

Figure 2 shows the continuum predicted with the **reflector.in** test case. The lower curve shows the total continuum that would






Figure 2 This is the continuum predicted by the input file reflector.in. The lower curve has been divided by two and shows the total spectrum produced by setting the line width to the speed of light. The upper curve shows the same thing, but with the line width set to 100 km/sec. reflector

be predicted if the *PunchLWidth* variable is set equal to the speed of light⁶. Here lines are added to the continuum such that the difference between vF_v at the line peak and vF_v for the underlying diffuse continuum is equal to the line flux. As a result the resulting line to continuum contrast is very small. The upper curve shows

 $^{^{6}}$ This was the default for version 90.00 through version 90.03. In C90.04 the default was changed to 1000 km/s. Before version 90 the line to continuum contrast depended on the cell width at the particular energy.

the same model but with the line contrast enhanced by entering the command **set PunchLWidth 100 km/sec**. The entire spectrum is shifted by a factor of two to make the two appear separated. The default line width is 1000 km s⁻¹.

The only effect of the **set PunchLWidth** command is to change the contrast in the punch output. The computed results and line intensities in other output are not affected. If the width is set to the speed of light then the intensities in the punch output will be correct but the line to continuum contrast too small. If the width is set to a small value the contrast is increased but the total intensity in the punch output will be greater than the actual emission. (Energy *will not* appear to have been conserved in this punch output).

4.6 Surface Brightness

Cloudy will normally predict $4\pi J$, the intensity radiated into 4π sr by a unit area of cloud, erg cm⁻² s⁻¹. Observations of resolved sources often measure the surface brightness, with units erg arcsec⁻² s⁻¹. Be careful! – some workers may report surface brightness with units erg arcsec⁻² s⁻¹ sr⁻¹. Remove the sr⁻¹ before continuing by multiplying by 4π .

To obtain the surface brightness we must divide the intensity $4\pi J$ predicted by **Cloudy** by the number of square seconds of arc in 4π sr. One radian is $180/\pi = 57.29578$ deg, so 1 sr is $(180/\pi)^2 = 3282.806$ deg², and there are 5.3464×10^{11} square arc seconds in 4π sr. The surface brightness (per square second of arc) is the intensity $4\pi J$ (per square centimeter) multiplied by the inverse of this, or 1.8704×10^{-12} cm² arcsec⁻².

4.7 Flux to luminosity

The luminosity is the flux of a line multiplied by the total area of the shell. For full coverage this is $4\pi r^2$ where r is the radius of the shell. If the shell only partially covers the continuum source then this should be multiplied by the covering factor.

4.8 Relative hydrogen line intensities

Very accurate ratios of Balmer or Paschen lines of hydrogen can be used to determine reddening. Ferguson and Ferland (1997) describe **Cloudy**'s hydrogen atom. It gives good results for levels below 10 in the code's default state, which uses a 15 level atom. The number of levels can be increased to \leq 50 using the **hydrogen levels** command, and this gives better results at the expense of more compute time. The larger atom should give results accurate to better than 5% for lines arising from below principal quantum number 10, and 10% accuracy for lines with upper levels between 10 and 15. The accuracy decreases for upper levels higher than 15 although the total recombination efficiency of the atom is computed to high precision.

For pure recombination lines you can easily do better than **Cloudy**. The code is limited by the size of the model hydrogen atom that can be computed on the fly. The definitive calculation for hydrogen recombination is that of Hummer and Storey (1987), who used a 1000 level atom with all l-states explicitly considered (that works out to something like a million levels!). Storey and Hummer (1995) placed a program on the web that will interpolate on their tables of case B hydrogen emission, for any temperature and density they computed. The best way to obtain a very high quality hydrogen recombination spectrum is to get the mean H⁺ temperature and the electron density (perhaps those predicted by *Cloudy*) and then use their interpolating code to provide the hydrogen spectrum for these conditions.

The Hummer and Storey (1987) calculation is for case B conditions, which assume that many processes are unimportant (see Ferguson and Ferland 1997). Neglected processes include collisional excitation from the ground or first excited states, induced processes where the incident continuum causes the atom to fluoresce, and line transfer in all non-Lyman lines. These assumptions are an excellent approximation for conventional nebulae, such as planetary nebular or HII regions. They are questionable for gas denser than 10⁶ cm⁻³ or when X-Rays are present. When any of these processes are important the hydrogen spectrum is far more model dependent and *Cloudy*'s results may be more realistic than the case B results.

4.9 Line Intensities in a dusty open geometry

Two sets of line intensities are printed if a dusty open geometry is computed. The second block of lines is the conventional set of intrinsic emission line intensities. When grains are present these intensities would need to be corrected for line of sight reddening to be compared with observations.

The first block of emission line intensities would be that emitted from the

illuminated face of a molecular cloud. The geometry is appropriate for the Orion Nebula, a blister HII region on the surface of Orion Molecular Cloud 1 (OMC1). An idealized geometry is shown in Figure 3. The code computes the fraction of the line emission that is directed towards the illuminated face. The remainder is emitted towards the neutral gas, which is assumed to have an infinite optical depth due to grains. The local albedo of the gasgrain mixture is computed, and the fraction reflected is passed back towards the illuminated face. The resulting intensities are roughly half what would be expected were the cloud emitting from both sides. Something like 10% of the line striking the molecular cloud will be reflected back to the observer.

So, for the illustrated blister the first block of lines gives what would be seen by an observer a large distance off to the left.

4.10 Continuum pumping




Figure 3 This shows that geometry assumed when computing the first block of lines in an open dusty geometry. The light area at left is the HII region, which is assumed to be a layer on the surface of an infinitely optically thick molecular cloud, the dark area on the right. Light can be emitted towards, and freely escape from, the illuminated face of the cloud. A fraction of the light emitted towards the molecular cloud is reflected back towards the illuminated face.

contribution to line intensities

Continuum pumping or fluorescence is included for all lines. The contribution is not usually printed, but will be if the **print line pump** command is entered. Whether this contribution actually adds to the observed line emission depends on the geometry. Continuum pumping increases the line emission if no related absorption occurs. This will be the case if the continuum source is either not observed or not covered by absorbing gas. If absorbing gas covers an observed continuum source then the situation is like the P Cygni problem, and pumping does not increase the total intensity of the line at all. The printed line intensity includes this contribution unless the **no induced processes** command is entered.

The output produced by the **punch continuum** commands does not include the pumped part of the line contribution. This is correct if the continuum source is included in the beam, but is not if only the gas is observed.

5 THE EMISSION LINES

5.1 Overview

The following two sections give a complete list of all emission lines predicted by *Cloudy*. Before version 90 of the code all lines were listed in sub-section 5.3, immediately following this section. The code is being modified to bring all lines into a common set of line transfer vectors. Lines which have been moved to this formalism are now listed in sub-section 5.4, beginning on page 451.

This is a complete list of the predicted quantities. Some continua, and various indications of contributors to lines and continua, are mixed in what follows. The previous section of the document describes how to convert these into some observed quantities. Not all are printed by default – the **print** commands described in Part I and also in section 5.4.3 starting on page 452 tell how to get more or fewer predictions.

5.2 Line wavelengths

In the current version of the code the line wavelength is an integer. This is to allow a precise match with a wavelength specified during a search. An integer format has obvious limitations in specifying fractions. To overcome this, the format of the wavelength changes with wavelength, in an attempt to have several significant figures.

Many wavelengths are too small or large to be expressed in Ångstroms. Many FIR lines would overflow the output format if their wavelengths were expressed in Ångstroms. For wavelengths longward of 100µm the wavelength is given in µm. For wavelengths one micron and 100µm the wavelength is given in tenths of microns. For wavelengths between 10Å and 100Å the wavelength is given in tenths of Å, and in hundredths of Å below this⁷. The [C II] λ 157 micron line would be C 2 157. The [O III] 88 micron line would be O 3 883. The logic deciding what format should be printed is given in routine *iWavLen*.

In the optical wavelengths are usually given in air. Continua are usually indicated by a wavelength of zero. Emission lines are broken into four large groups in the following sections. Level 2 lines are described on page 451 (section 5.4.2) below. These lines use Opacity Project wavelengths, which are generally good to about 10%.

5.3 Miscellaneous Line Entries

These are a series of entries which contain most of the optical forbidden lines, some continua, and identify various contributors to the main lines.

For this set of lines, the first column gives the four character label printed in the final array listing and the second column gives the wavelength of the line, using the conventions described above. The label in the first column is the one used to access the line using the *cdLine* routine described elsewhere.

⁷ Before version 94.01 of the code the wavelength was given in tenths of Å below 100Å. The **set oldwavelength** command will cause the code to continue to do this.

The third column character indicates whether the entry is a heat source (indicated by h), a coolant (c), a recombination line (r), or an intensity entered for information only (i). The last column gives a brief description of the meaning of the line prediction. More information about individual lines can usually be had by examining the comments associated with the actual calculation of the line strength (in subroutine *coolr*) or in subroutine *lines*, where the line intensity is entered into the storage array.

Label λ

Description

TOTL 4861 i these lines added to outlin in metdif - following must be false TOTL 1216 i total Ly-a from multi-level atom total luminosity in incident continuum Inci 0 i 0 i TotH total heating, all forms, information since individuals added later BFH1 0 h hydrogen photoionization heating, ground state only BFHx 0 h normally zero, positive if excited states are net heating 0 h heating due to induced lines absorption of continuum Line Heli 0 C net cooling due to collisional ionization of Heo 3He1 0 h this is the heating due to 3-body recombination He2i 0 C net cooling due to collisional ionization of He+ 0 h this is the heating due to 3-body recombination 3He2 BFHe 0 h total helium photoionization heating, all stages TotM 0 h total heavy element photoionization heating, all stages net free-free cooling, nearly cancels with cooling in lte HFFC 0 C HFFh 0 h net free-free heating, nearly cancels with cooling in lte H FF 0 i H brems (free-free) cooling 0 i FF H total free-free heating ComH 0 h Compton heating Н- Н 0 h H- heating H-HC 0 C induced H- cooling 0 h GrGH gas heating by grain photoionization gas cooling by grain collisions GrGC 0 C "extra" heat added to this zone, from HEXTRA command 0 h extH extC 0 C "extra" cooling added to this zone, from CEXTRA command pair 0 h heating due to pair production Ca B 4861 i H beta recombination, assuming case B Ca B 1216 i case b intensity of Ly-alpha, no two photon H-beta computed from Q(H) and specified covering factor O(H) 4861 i Q(H) 1216 i Ly-alpha from Q(H), high-dens lim, specified covering factor TOTL 1216 i these lines added to outlin in metdif - following must be false Inwd 1216 i inward part of Lya TOTL 0 r these lines added to outlin in metdif - following must be false 0 i Tnwd Clin 912 c total collisional cooling due to all hydrogen lines Hlin 912 h total collisional heating due to all hydrogen lines 2 NU 0 r 2-photon two photon emission from multi-level atom LA X 1216 i la contribution from suprathermal secondaries from ground H-CT 6563 i H-alpha produce by H- mutual neutralization Ind2 1216 i "Ly alpha" produced by induced two photon Pump 4861 r H-beta produced by continuum pumping in optically thin ld limit CION 0 C collision ionization cooling of hydrogen 3bHt 0 h this is the heating due to 3-body recombination Strk 1216 i Stark broadening contribution to line Strk 6563 i Stark broadening contribution to line Strk 4861 i Stark broadening contribution to line Strk 18751 i Stark broadening contribution to line Strk 40512 i Stark broadening contribution to line Dest 1216 i portion of line lost due to absorp by background opacity Dest 6563 i portion of line lost due to absorp by background opacity Dest 40516 i portion of line lost due to absorp by background opacity

Dest 4861 i portion of line lost due to absorp by background opacity Dest 18751 i portion of line lost due to absorp by background opacity Fe 2 1216 i Ly-alpha destroyed by overlap with FeII e sc 1216 i electron scattering escape contribution to line e sc 6563 i electron scattering escape contribution to line e sc 4861 i electron scattering escape contribution to line e sc 18751 i electron scattering escape contribution to line e sc 40512 i electron scattering escape contribution to line e-e+ 511 i 511keV annihilation line nFnu 0 i total continuum produced by cloud at selected energy points 0 i InwT reflected diffuse continuum 0 i reflected incident continuum (only incident) InwC Inci 4860 i incident continuum nu*f_nu at H-beta, at illuminated face of cloud Inci 1215 i incident continuum nu*f_nu near Ly-alpha, at illuminated face of cloud Ba C 0 i integrated Balmer continuum emission 0 i PA C Paschen continuum emission GraT 0 i total grain heating by all sources, lines, collisions, incident continuum GraT 0 i grain heating by incident continuum GraL 1216 i grain heating due to destruction of Ly alpha GraC 0 i grain heating due to collisions with gas GraD 0 i grain heating due to diffuse fields, may also have grain emission FF X 0 i part of H brems, in x-ray beyond 0.5KeV ComC 0 C total Compton cooling expansion cooling, only non-zero for wind Expn 0 C eeff 0 C electron - electron brems H FB 0 i H recombination cooling HFBC 0 C net free-bound cooling and heating HFBh 0 h Hind 0 C cooling due to induced rec of hydrogen cooling due to induced rec of fully ionized helium 3He2 0 C 0 C cyclotron cooling Cycn 1216 i collisionally excited La cooling Cool Heat 1216 i collisionally de-excited La heating Crst 960 i cooling due to n>2 Lyman lines Hrst 960 i heating due to n>2 Lyman lines Crst 4861 i cooling due to n>3 Balmer lines heating due to n>3 Balmer lines Hrst 4861 i 0 i cooling due to higher Paschen lines Crst 0 i heating due to higher Paschen lines Hrst H2 1 2 C H2 rotation lines from Lepp and Shull ApJ 270, 578. H2dC 0 C H2 dissociation by H atoms (not e) H2dH 0 h heating by H2 dissociation by Lyman continuum $\operatorname{neg}\xspace{1.5}\xspace{1.5}$ H ion free-bound emission H-FB 0 C H2+ 0 C H+ + H => H2+ + photon continuum cooling HEH+ 0 i HeH+ formation cooling COdh 0 h carbon monoxide co photodissociation CO C 0 C cooling due to coll of vib rot levels HeFF 0 C He brems emission HeFB He recombination cooling 0 C MeFB 0 C heavy element recombination cooling MeFF 0 C metal brems emission Toff 0 i total brems emission He3I 0 C He I triplet net collisional ionization cooling He3b 0 h He I triplet net 3-body heating HeIC 584 c He I 584, collisional excitation cooling He I 584, collisional de-excitation heating HeIH 584 h 504 i He I He I rec to ground escaping cloud these lines added to outlin in metdif - following must be false these lines added to outlin in metdif - following must be false 584 i esc esc 626 i Ca B 4471 i He I 4471 recombination only, fit to Brocklehurst '72 TOTL 5876 i these lines added to outlin in metdif - following must be false Inwd 5876 i inward part of 5876 Ca B 5876 r He I 5876 REC, simple fit to Brocklehurst Ca B 6678 i He I 6678 REC, simple fit to Brocklehurst

TOTL 10830 i these lines added to outlin in metdif - following must be false He I 10830 produced by radiative recombination reco 10830 i coll 10830 i collisionally excited 10830 estimated from Clegg 1987 (not model atom) Inwd 10830 i inward escaping HeI 10830 TOTL 3889 i these lines added to outlin in metdif - following must be false TOTL 7065 i these lines added to outlin in metdif - following must be false total collisional He I triplet line cooling, from n-level atom CCHE 0 C total collisional de-exec He I heating, from n-level atom ChHE 0 h Ca B 2 i case B He I 2.06 micron TOTL 2 i He I 2.06 micron from model atom, all physics in Ca B 5016 i Case B He I 5016 HeII 228 i He II Lyman continuum He2C 911 i He II Balmer continuum escaping from cloud TOTL 304 i these lines added to outlin in metdif - following must be false these lines added to outlin in metdif - following must be false TOTL 1640 i Inwd 1640 i inward part of HeII 1640 тотт, 1217 і these lines added to outlin in metdif - following must be false Inwd 1217 i He II 1216 inward fraction TOTL 1085 i these lines added to outlin in metdif - following must be false Inwd 1085 i He II 1085 inward fraction TOTL 4686 i these lines added to outlin in metdif - following must be false Inwd 4686 i inward beamed part of 4686 TOTL 3205 i these lines added to outlin in metdif - following must be false Inwd 3205 i HeII 3205 inward fraction Ca B 1640 i He II 1640, case B at low densities Ca B 4686 i He II 4686, case B cooling due to collisional ionization of heavy elements Mion 0 C Li3r 19 i these lines added to outlin in metdif - following must be false these lines added to outlin in metdif - following must be false 19 i Be4r 19 i Bo5r these lines added to outlin in metdif - following must be false REC 1656 i C 1 1656 recomb; n.b. coll deexcitation not in C IC 9850 C C 1 9850, coll excit C Ir 9850 i was a big mistake TOTL 9850 i total intensity, all processes, C I 9850 С 1 8727 с C 1 8727; equivalent to 4363 C 1 4621 c 1S - 3P Phot 2326 i photoproduction, Helfand and Trefftz 1335 i C 2 1335 recombination, REC C II 3134 c C 2 intercombination line with same upper state as 1335 C3 R 977 i dielectronic recombination contribution to C 3 977 P386 977 r C 3 977 pumped by continuum near 386A TOTL 1909 i C 3 1909 collision, both lines together C 3 1907 i C 3 1908 j-2 to ground C 3 1909 recombination from Storey C3 R 1909 i Phot 1909 i C 3 1909 following relax following inner shell photoionization Rec 1175 i dielectronic recombination contribution to C 3 1175 TOTL 1549 i total intensity of C 4 1549, all processes Inwd 1549 i inward part of C 4 DEST 1549 i part of line destroyed by photoionization of Balmer continuum C4 r 1549 i recombination C 4 1549 from CV these lines added to outlin in metdif - following must be false C 6r 34 i N 1 5200 i N 1 5200, both 5198, 5200, collisions and recombination Coll 5200 c N 1 5200, both 5198, 5200, collisions and recombination 5200 i REC recombination contributon to [NI] 5200 N 1 3466 c [N 1] 3466, 3 - 1 transition, whole multiplet N 1 10400 c [N 1] 10400 3 - 2 transition, whole multiplet N 2 6584 c N 2 6584 alone N 2 6548 c N 2 6548 alone REC 6584 i N 2 6584 alone, recombination contribution N 2 5755 i N 2 5755 total, collisions plus charge transfer Coll 5755 c N 2 5755 collisional contribution СТ 5755 c N 2 5755 charge transfer contribution 1085 i dielectronic recombination contribution to N 2 1085 Rec

N2cn 1 i continuum pumped N 2 6584 N2cn 5755 i continuum pumped N 2 5755 N3cn 4640 i continuum pumped "Bowen" N 3, optically thin excited line N3cn 4634 i continuum pumped "Bowen" N 3, optically thin excited line N3cn 4642 i continuum pumped "Bowen" N 3, optically thin excited line 990 i total N 3 990, both electron excitation and continuum pumping extr rec 990 i part of N 3 990 due to recombination N 3p 990 r N 3 989.8, continuum pumped TOTL 1486 i N 4] 1486, total intensity of both lines N 4 1485 i the N 4] slow transition by itself rec 765 i N 4 765 recombination, TOTL 1240 i N 5 1240, total emission, collisions plus pumping Inwd 1240 i inward part of N 5 N 7r 25 i these lines added to outlin in metdif - following must be false these lines added to outlin in metdif - following must be false Fl7r 19 i 0 1 6300 c total Oxygen I 6300, including line optical depth 0 1 6363 C total Oxygen I 6363, including line optical depth 0 1 5577 c auroral OI TOIC 0 C total collisional cooling due to 6-level OI atom TOTh 0 h total collisional heating due to 6-level OI atom 6lev 8446 i be moved to call PutLine 6lev 1304 i OI 1304 from six level atom 6lev 1039 i OI 1039 from six level atom 6lev 4368 i OI 4368 from six level atom 6lev 13 i OI 1.3 micron from six level atom 6lev 11 i OI 1.1 micron from six level atom 6lev 29 i OI 2.9 micron from six level atom 6lev 46 i OI 4.6 micron from six level atom TOTL 3727 c O II 3727, all lines of multiplet together TOTL 7325 c O II 7325, all lines of multiplet together IONZ 3727 i line produced by photoionization of Oo; already in TOTL IONZ 7325 i line produced by photoionization of Oo; already in TOTL O II 3729 i five level atom calculations; D5/2 - S3/2 O II 3726 i D3/2 - S3/2 transition 0 II 2471 c both 2P 1/2 and 3/2 to ground P1/2-D5/2 and P3/2-D5/2 together O II 7323 i 7332 i O TT P1/2-D3/2 and P3/2-D3/2 together TOTL 1665 i total intensity of OIII] 1665, all processes Phot 1665 i contribution to OIII 1665 due to inner shell (2s²) ionization Augr 1665 i contribution to OIII 1665 due to K-shell ionization 0 3 5007 c fac = c5007/(1.+1./2.887)0 3 4959 c O III 4959 alone, collisions, tot OIII is this times 4 LOST 5007 i O III 5007 lost through excit photo TOTL 4363 i O III 4363, sum of rec, coll, ct excitation Coll 4363 c O III 4363, collisions from five level atom Rec 4363 i O III 4363 recombination, coef from Burgess and Seaton 0 3 2321 c collisional excitation of 2321, 5-level atom C EX 4363 i charge exchange, Dalgarno+Sternberg ApJ Let 257, L87. C EX 5592 i charge exchange rate, D+S 835 i O III 834A, dielectronic recombination only rec InSh 1401 i inner shell photoionization, relaxation 789 i O IV 789A, dielectronic recombination only rec rec 630 i O V 630A, dielectronic recombination only TOTL 1218 i O V 1218], total intensity of both lines 0 5 1211 i the slow transition by itself 0 5 5112 i BS O V 5112, recombination TOTL 1035 i O VI 1035, total of pumping and collisional excitation Inwd 1035 i inward part of OVI line 0 8r 19 i recombination from fully stripped ion Ne III 3869, of 3968+3869 doublet Ne III 3968, of 3968+3869 doublet Ne 3 3869 c Ne 3 3968 C Ne 3 3343 c NeIII auroral line Ne 3 1815 c NeIII auroral line

Ne 4 2424 c Ne IV 2424, collisional excitation Ne IV N=3 lines, three level atom approx Ne 4 4720 c Ne 4 1602 c Ne IV N=3 lines, three level atom approx Ne 5 3426 c Ne V 3426 of 3426, 3346 doublet Ne V 3346 of 3426, 3346 doublet Ne 5 3346 c Ne 5 2976 c auroral line Ne 5 1575 c collisionally excited Ne 5 1141 c both components of 5S-3P 1146.1, 1137.0 doublet TOTI 895 i Ne VII 895, collisionally excited, both lines Ne 7 890 i Ne VII 890, single line TOTL 774 i Ne VIII 774, collisionally excited 774 i inward part of NeVIII 774 line Inwd 12 i these lines added to outlin in metdif - following must be false NeLr Na 5 1365 c [NaV] 1365, sum of 1365.1+1365.8; cs only guess Na 5 2067 c [NaV] 2067, sum of 2066.9+2068.4; cs only guess Na 5 4017 c [NaV] 4017, sum of 4010.9+4016.7+4022.7; cs only guess [Na VI] 2568.9 Na 6 2569 c Na 6 1357 c [Na VI] 1356.6 Na 6 2972 c [Na VI] 2971.9 Na 6 2872 c [Na VI] 2872.7 these lines added to outlin in metdif - following must be false NaLr 10 i TOTL 2798 i Mg II 2798 Inwd 2798 i inward part of Mg II 2798 Mg 6 1806 c MG VI TOTL 615 i Mg 10 614.9 bothof doublet, li seq 2s 2p MqLr 7 i these lines added to outlin in metdif - following must be false totl 2665 i total emission in Al II] 2669.7, 2660 doublet Al 2 2660 i emission in Al II] 2669 alone TOTL 1860 i Al III Inwd 1860 i inward part of AlIII line Al 6 2428 c Al 6 2601 c [Al VI] 2428.4 [Al VI] 2601.0 Al 6 1170 c [Al VI] 1169.86 Al 6 2125 c [Al VI] 2124.95 TOTL 556 i Al 11, Li seq 2s2p these lines added to outlin in metdif - following must be false AlLr 6 i diel 1260 i SI II 1260, rough guess of dielec contribution diel 1909 i dielectronic recombination SiII 1909 rec 1207 i Si III 1207, dielectronic recombination only TOTL 1888 i Si III] 1892+1883, total intensity of both lines Si 3 1883 i Si III] 1883 by itself PHOT 1895 i photoproduction by inner shell removal TOTL 1397 i Inwd 1397 i Si IV 1397, collisionally excited inward part of SiIV 1397 Si 7 2148 c SI VII, 2148, O III like, collisionally excited Si 7 2148 c Si 8 1446 c SI VIII 1446, OIII like, collisionally excited Si 9 1985 c SI IX 1985, 2150, collisionally excited Si 9 949 C collisionally excited Si 9 1815 c collisionally excited Si 9 691 C both components of 5S-3P doublet Si10 606 C SI 10 606A, actually group of 4 intercombination lines. Si11 583 c SI XI 582.9, collisionally excited TOTL 506 i 6 i these lines added to outlin in metdif - following must be false SiLr P15r 19 i these lines added to outlin in metdif - following must be false S 1R 1807 i this is to check whether photoexcit of S II is ever important S 2 6720 c S II 6731 + 6716 together S 2 4074 c S II 4070 +4078 together S 2 10330 c S II N=3 lines, all four lines together S II 6731 i individual line from five level atom S II 6716 i individual line from five level atom S II 4070 i individual line from five level atom S II 4078 i individual line from five level atom

S II 10323 i individual line from five level atom S II 10289 i individual line from five level atom S II 10373 i individual line from five level atom S II 10339 i individual line from five level atom S 3 9532 c [S III] 9532 alone [S III] 9069 alone S 3 9069 c S 3 6312 C [S III] 6312, transauroral temperature sensitive S 3 3722 c [S III] 3722, same upper level as 6312 TOTL 1198 i S V 1198] both lines together S 5 1188 i Be seq, weaker of the two transitions total S VI 933+944 TOTL 933 i S 9 1715 c S IX 1715, 1987, collisionally excited S 10 1213 C S X 1213, 1197, collisionally excited S 11 1826 c S XI 1615, 1826, collisionally excited group of four intercombination lines S 12 520 c S 13 488 c S XIII 488.4, 1909 like, collisionally excited TOTT 427 i S 14 506 li seq 2s2p S LR 5 i these lines added to outlin in metdif - following must be false S LR 5 i Cl 2 8579 c Chlorine II 8581, 9127 doublet Cl 2 9127 c Chlorine II 8581, 9127 doublet Cl 2 9127 c Cl 2 6164 c Chlorine II 6164 auroral line Cl 2 3676 c Chlorine II 3679 auroral line TOTL 5525 c Cl III 5519, 5539 doublet, both together TOTL 3350 c Cl III 3354, 3344 doublet, both together TOTL 8494 c Cl III 8504, 8436, 8552, 8483 multiplet, all together Cl 3 5538 i Cl III 5538 Cl 3 5518 i Cl III 5518 Cl 3 3354 i Cl III 3354 Cl 3 3344 i Cl III 3344 Cl 3 8504 i Cl III 8504 Cl 3 8436 i Cl III 8436 Cl 3 8552 i Cl III 8552 Cl 3 8483 i Cl III 8483 ClIV 8047 Cl 4 8047 c Cl 4 7532 c ClIV 7532 Cl 4 3119 c ClIV 3119 Cl 4 5324 c ClIV 5324 Cl 4 5324 c ClRr 4 i Cl 17 ly a recombination 3.7A from fully stripped ion Argon III 7135 Ar 3 7135 c 7751 c Argon III 7751 Ar 3 Argon III 5192 Ar 3 5192 c Ar 3 3109 c Argon III 3109 Ar 3 3005 c Argon III 3005 TOTL 4725 i Argon IV 4711 + 4740 together, 4740=90% TOTL 2860 i [AvIV] 2868, 2854 together TOTL 7250 i [AvIV] auroral lines, 7237, 7331, 7171, 7263 [Ar IV] 4740 Ar 4 4740 c Ar 4 4711 c [Ar IV] 4711 [Ar IV] 2868 Ar 4 2868 c Ar 4 2854 c [Ar IV] 2854 [Ar IV] 7263 Ar 4 7263 c Ar 4 7171 c [Ar IV] 7171 [Ar IV] 7331 Ar 4 7331 c Ar 4 7237 c [Ar IV] 7237 Ar 5 7005 c Argon V, 3P lines, 7005, collisionally excited Ar 5 6435 c Argon V, 3P lines, 6435, collisionally excited 6435 c Ar 5 Ar14 4413 c Ar XIV 4413, predicted lambda, not observed(??) Ar15 409 c collisionally excited these lines added to outlin in metdif - following must be false ArRr 4 i

K19r 4 i these lines added to outlin in metdif - following must be false Ca 2 3933 c coll excit calcium k+h Ca 2 8579 c infrared triplet Ca 2 7306 c forbidden lines, 7291+7324 together Phot 3933 i fraction H Ly-alpha destruction of excited levels Phot 7306 i fraction H Ly-alpha destruction of excited levels Ca2K 3934 i individual lines from five level atom Ca2H 3969 i individual lines from five level atom Ca2X 8498 i individual lines from five level atom Ca2Y 8542 i individual lines from five level atom Ca2Z 8662 i individual lines from five level atom CaF1 7291 i individual lines from five level atom CaF2 7324 i individual lines from five level atom Rec 3933 i reccombination contribution to CaII emission Ca 5 6087 c Ca V optical and uv lines, collisional excitation, 3-level atom Ca 5 5311 c Ca V optical and uv lines, collisional excitation, 3-level atom Ca 5 2414 c Ca V optical and uv lines, collisional excitation, 3-level atom Ca 5 3997 c Ca V optical and uv lines, collisional excitation, 3-level atom Ca 7 5620 c Ca VII optical and uv lines, collisional excitation, 3-level atom Ca 7 4941 c Ca VII optical and uv lines, collisional excitation, 3-level atom Ca 7 2112 c Ca VII optical and uv lines, collisional excitation, 3-level atom Ca 7 3688 c Ca VII optical and uv lines, collisional excitation, 3-level atom CaLr 3 i these lines added to outlin in metdif - following must be false 3 i these lines added to outlin in metdif - following must be false ScLr Sc 2 21 c Sc II 2.08 (1-3) Sc 2 41 c Sc II 4.1 mic (1-2) Sc 2 42 C Sc II 4.22 (2-3) Sc 3 3933 c Sc III 3936 Sc 6 5054 c Sc VI 5054 (1-2) Sc 6 3592 c Sc VI 3595 (2-3) Sc 6 2100 c Sc VI 2100 (1-3) TiLr 3 i these lines added to outlin in metdif - following must be false Ti III 1.21 micron, (actually multiplet) 2-1 transition from model atom Ti 3 12 C Ti 3 9594 c Ti III 9594, 3-1 transition, (actually multiplet) from model atom Ti 3 45 c Ti III 4.57 micron, 3-2 transition, (actually multiplet) from model atom these lines added to outlin in metdif - following must be false V Lr 3 i V 3 8823 c V 3 8507 c V TTT 8823 V III 8507 V 3 8507 c V IV 7741 1-3 V 4 7735 c V 4 V IV 9496 2-1 9489 c V IV 4.19 mic 3-2 V 4 42 C CrLr 3 i these lines added to outlin in metdif - following must be false [CrIII] multiplet blend at 5828A Cr 3 5828 c Cr 4 7267 c [CrIV] 2 - 1 multiplet blend at 7272 Cr 4 6801 c [CrIV] 3 - 1 multiplet blend at 6806 Cr 5 7979 c [CrV] 2 - 1 multiplet blend at 7985 [CrV] 3 - 1 multiplet blend at 6582 Cr 5 6577 c Cr 5 37 C [CrV] 3 - 2 multiplet blend at 3.75 microns 3 i these lines added to outlin in metdif - following must be false MnLr Fe 2 6200 i Fe 2 the 3-2 transition of Netzer's atom Fe 2 4300 i Fe 2 forbidden 2-1 transition from Netzer's atom Fe 2 2400 i Fe 2 UV3, 3-1 transition from Netzer's atom 0 C total of all UV+optical Fe 2 cooling Fe2c Fe2h 0 h Fe 2 1100 i 1 to 6 transition of Fred's Fe 2 atom Fe 2 1500 i 2 to 6 transition of Fred's Fe 2 atom Fe 2 11500 i 3 to 4 transition of Fred's Fe 2 atom Fe 2 2500 i 3 to 5 transition of Fred's Fe 2 atom 2300 i 4 to 6 transition of Fred's Fe 2 atom Fe 2 Fe 2 8900 i 5 to 6 transition of Fred's Fe 2 atom all cooling due to 16 level atom Fe 2 0 C Fe 2 166 i Fe 2 1.664 microns 8-13

```
Fe 2
       160 i
              Fe 2 1.599 microns 7-12
Fe 2
       153 i
              Fe 2 1.534 microns 6-11
Fe 2
       164 i
               Fe 2 1.644 microns 6-10
Fe 2
       128 i
              Fe 2 1.279 microns 12-4
       130 i
Fe 2
              Fe 2 1.295 microns 11-3
Fe 2
       133 i
              Fe 2 1.328 microns 11-4
       126 i
Fe 2
              Fe 2 1.257 microns 10-1
Fe 2
       132 i
               Fe 2 1.321 microns 10-2
Fe 2
       259 i
               Fe 2 25.988 microns 2-1
Fe 2
       353 i
              Fe 2 35.348 microns 3-2
Fe 2
       178 i
               Fe 2 17.936 microns 7-6, label is 178 to be unique
Fe 2
       245 i
               Fe 2 24.518 microns 8-7
Fe 2
       358 i
               Fe 2 35.776 microns 9-8
Fe 2
       181 i
              Fe 2 1.810 microns 10-7
Fe 2
       168 i
              Fe 2 1.677 microns 11-7
              Fe 2 1.800 microns 11-8
Fe 2
      180 i
Fe 2
       171 i
              Fe 2 1.712 microns 12-8
Fe 2
      179 i
              Fe 2 1.798 microns 12-9
Fe 2
       229 i
              Fe 2 22.902 microns 11-10
Fe 2
      347 i
              Fe 2 34.660 microns 12-11
Fe 2 8619 i
              Fe 2 8619A 14-06
Fe 2 8894 i
              Fe 2 8894A 15-07
      9229 i
Fe 2
              Fe 2 9229A 15-08
Fe 2
      9270 i
              Fe 2 9270A 16-09
                emission from lage FeII atom, integrated over band
Fe2b
         2 i
Fe 3
               sum of 3p and 3g states together
         0 C
              Fe 3 5270, predictions from garstang et al 78
Fe 3 5270 c
Fe 3 4658 c
              Fe 3 5270, predictions from garstang et al 78
Fe 4
         0 C
               total cooling due to 12-level Fe 4 atom
     3096 i
              Fe 4 3096.A, 4-1 and 5-1 transitions together
Fe 4
Fe 4 2836 i
               Fe 4 2835.7A, 6-1 transition, 4P5/2 - 6S5/2
      2829 i
               Fe 4 2829.4A, 7-1 transition, 4P3/2 - 6S5/2
Fe 4
Fe 4
     2567 i
              Fe 4 2567.6+ 2567.4. 11-1 and 12-1 transitions
Fe 4
       277 i
              Fe 4 2.774 microns 12-7 transition
Fe 4
       271 i
              Fe 4 2.714 microns 12-6 transition
Fe 4
       272 i
              Fe 4 2.716 microns 11-6 transition
Fe 4
       281 i
               Fe 4 2.806 microns 10-7 transition
Fe 4
       287 i
               Fe 4 2.865 microns 10-8 transition
Fe 4
       284 i
              Fe 4 2.836 microns 9-6 transition
Fe 5 3892 c
              Fe 5 3892+3839
Fe 6
         0 C
               all of 2G lines together first
    5177 c
              Fe 6 5177, approximate correct
Fe 6
               [Fe 7] 6087
      6087 c
Fe 7
Fe 7
      5722 c
               [Fe 7] 5722
Fe 7
      242 C
              Fe 9 242 j=1 slower decay
Fell
     2649 c
              Fe 11 2649 collisional excitation
Fe11
     1467 C
              Fe 11 1467 collisional excitation
Fe12
     1242 c
              Fe 12, 1242, 1349 together, collisional excitation
      2170 c
              Fe 12, 2170, 2406 together, collisional excitation
Fe12
     2568 c
               Fe12 2904, 2567, 3567, 3073 together, collisional excitation
Fe12
Fe14 5303 i
              Fe 14 optically thin in line 344
Coll
     5303 c
               contribution from collisional excitation
Pump
     5303 r
              continuum fluorescense
347
     5303 c66 error! put this in
      592 c Fe 19 from loulergue et al '85
Fe19
Fe19
     7082 C
              Fe 19 from loulergue et al '85
Fe19
     1118 c
              Fe 19 from loulergue et al '85
      1328 c
              Fe 19 from loulergue et al '85
Fe19
Fe22
       846 C
              Fe 22 845.6A
Fe23
       263 c
              Fe 23 1909-like 262.6
FeKa
         2 i
              total intensity of K-alpha line
         2 i
              recombination from fully stripped ion
FeLr
               total hot iron Ka; Auger "hot" iron, plus recom
TotH
         2 i
              Auger production of "cold" iron, less than or 17 times ionized
AugC
         2 i
```

```
CoLr
         1 i
                these lines added to outlin in metdif - following must be false
         1 i
               these lines added to outlin in metdif - following must be false
NiLr
CuLr
         1 i these lines added to outlin in metdif - following must be false
         1 i these lines added to outlin in metdif - following must be false
ZnLr
Stoy 0 i optional sum of certain emission lines, set with "print sum"
BaC 3646 i residual flux at head of Balmer continuum, nuFnu
cdif 3646 i
               residual flux in Balmer continuum, nuFnu
cout 3646 i residual flux in Balmer continuum, nuFnu
cref 3646 i residual flux in Balmer continuum, nuFnu
thin 3646 i residual flux in Balmer continuum, nuFnu
```

5.4 Emission line identifications

This section lists three groups of lines. The groups are discussed in the order they occur. The line list was automatically generated by the code with the **punch line data** command. All quantities were evaluated to 10^4 K. The description of the command in Part I of this document explains how to evaluate the quantities at other temperatures.

5.4.1 Optically thin recombination lines

The first group consists of all recombination lines of C, N., and O, with coefficients taken from Nussbaumer and Storey (1984) and Pequignot, Petitjean, and Boisson (1991). For this set, the spectroscopic designation is followed by the wavelength and the log of the recombination coefficient evaluated at 10⁴ K.

These are all predictions for optically thin pure recombination. These should be accurate for classical nebulae, such as planetary nebulae and HII regions. They will not be accurate for dense environments where optical depths and collisional effects come into play. There are several instances where more than one line of an ion will have the same wavelength due to the integer Ångstrom format used for wavelengths. The worst case is O V 4953, where three lines of the same multiplet have the same wavelength.

5.4.2 Transferred heavy element lines

The following group lists all the lines now included in the line transfer arrays, except for lines of H and He. These lines fall into two groups, referred to as level 1 and level 2 lines.

Level 1 lines have quantal collision strengths, experimental energies, and are treated including destruction by background opacity sources.

Level 2 lines have only g-bar collision strengths and so their intensities are far less accurate than level 1 lines. Radiative data for level 2 lines are taken from the Opacity Project, and *Cloudy* uses the line energies given there. As a result these lines have wavelengths that are uncertain by typically 10%.

Line labels These lines are all treated in a fairly automatic way, on a common basis. The ion is the first column of the table. This is in a uniform format, beginning with the two character element symbol and followed by an integer indicating the level of ionization. "C 2" is C⁺ or CII. This is followed by the integer wavelength label used to identify the line in the printout. The third column, with the label "WL", is the correct wavelength of the line, with units of either microns ("m") or Angstroms

("A"). The remaining columns give the statistical weights of the lower and upper levels, the product of the statistical weight and the oscillator strength, and then the Einstein A.

The last column is the electron collision strength, generally evaluated at 10⁴ K. Exceptions are lines whose collision strengths are only evaluated for temperatures far from 10⁴ K, for instance , a Fe XXV transition. Usually these collision strengths are for only the indicated transition, although in some cases (the Be sequence) the value is for the entire multiplet. This is discussed further in the section on the evaluation of the cooling function in HAZY II.

5.4.3 Output produced for the transferred lines

Because the lines have a common format within their storage vectors, the output has a common format too. Generally only the total intensity of the transition, the result of the solution of a multi-level atom with all processes included, is done. The approach used to compute the level populations is described in Part II of Hazy, and includes continuum pumping, destruction by background opacities, and trapping.

The total intensity of the transition is printed in a form like "C 2 1335", with the spectroscopic identification given by the first part, as found in the first column of the table, and the wavelength as indicated by the integer in the second column of the table.

In a few cases (for instance, the C $4 \lambda \lambda 1548$, 1551 doublet), a total intensity is also derived. In these cases the label "Totl" will appear together with an average wavelength (1549 in this case). These lines are all explicitly shown in section 5.3 on page 442 above.

It is possible to break out various contributors to the lines with options on the **print line** command, described in Part I of this document and in the following. These contributors are printed following the total intensity.

print line heating An emission line will heat rather than cool the gas if it is radiatively excited but collisionally de-excited. The print out will include an entry beginning with the label "Heat" if this printout is turned on.

print line collisions The collisional contribution to the lines will be printed, with the label "Coll".

print line pump The contribution to the total line, produced by continuum pumping, is printed with the label "Pump". What is observed? Whether or not this is a net emission process contributing to the observed line intensity depends on the geometry, whether or not continuum source is in the beam. At some velocities within the line profile this can be a net emission process, due to absorption at other velocities. If the continuum source is in the beam and gas covers it, this is not a *net* emission process, since photons are conserved.

print line inward The inwardly directed part of the total emission is printed with the label "Inwd". This can be greater than half of the line intensity if the line is optically thick.

print line optical depths At the end of the calculation the optical depths for all optically thick lines will be printed. This is not done by default since it is quite long.

5.5 The main blocks of lines

In previous versions of this document a large list of emission lines appeared here. This list is now far too large to include here. Rather, the list can be generated by executing the code with the command **punch line data ``filename.txt''** included. This will create a file that

includes the full set of lines that was presented here. Note that the lines that are output are only those that exist when the code is run. It is possible to make many of the model atoms and molecules as large or small as you like, and the actual lines that exist when the punch command is entered will be output.

5.6 Atomic data sources

Codes like **Cloudy** can only exist because of the large body of work done by the atomic and molecular physics community. This work will only continue to be supported if it is cited in the literature whenever it is used. The following is a partial list of citations for the atomic data used within the code.

Aggarwal, K.M. 1983, JPhB, 16, L59 Aggarwal, K.M. 1983, MNRAS, 202, 15P Aggarwal, K.M. 1984, ApJS, 54, 1 Aggarwal, K.M., 1985 Ast Ap 146, 149. Aggarwal, K.M., Callaway, J., Kingston, A.E., Unnikrishnan, K. 1992, ApJS, 80, 473 Allard, N., Artru, M.-C., Lanz, T., & Le Dournef, M. 1990, A&AS, 84, 563 Baluja, K.L. 1985, JPhB, 18, L413 Baluja, K.L., & Zeippen, C.J. 1988, J Phys B 21, 1455 Berrington, K.A. 1985, JPhB, 18, L395 Berrington, K.A. 1988, J.Phys.B, 21, 1083. Berrington, K.A. 1988, JPhB, 21, 1083 Berrington, K.A., Burke, P.G., Dufton, P.L., Kingston, A.E. 1985, ADNDT, 33, 195 Bhatia, A.K., & Doscheck, G.A. 1995 ADNDT, 60, 9 Bhatia, A.K., & Mason, H.E. 1986, A&A, 155, 413 Blum, R.D., & Pradhan, A.K. 1992, ApJS 80, 425 Brage, T., Froese-Fischer, C., Judge, P.G. 1995, ApJ 445, 457 Brage, T., Hibbert, A., Leckrone, D.S. 1997, ApJ, 478, 423 Brage, T., Judge, P.G., & Brekke, P., 1996, ApJ 464, 1030 Burke, V.M., Lennon, D.J., & Seaton, M.J. 1989, MNRAS, 236, 353 Butler, K., & Zeippen, C.J. 1984, A&A, 141, 274 Butler, K., & Zeippen, C.J. 1994, A&AS, 108, 1 Butler, S.E., & Dalgarno, A. 1980, ApJ 241, 838 Calamai, A.G., & Johnson, C.E., 1991, Phys Rev A, 44, 218 Calamai, A.G., Smith, P.L., Bergeson, S.D. 1993, ApJ, 415, L59 Callaway, J. 1994, ADNDT, 57, 9 Callaway, J., Unnikrishnan, K., & Oza, D.H. 1987, Phys Rev A, 36, 2576 Chandra, S. 1982, SoPh, 75, 133 Chandra, S. 1982, SoPh, 75, 133 Chidichimo, M.C. 1981, JPhB, 14, 4149

Cochrane, D.M., & McWhirter, R.W.P. 1983, PhyS, 28, 25

Dopita, M.A., Mason, D.J., Robb, W.D. 1976, ApJ, 207, 102 Doschek, G.A. 1986, ApJ, 300, 448 Dufton, P.L., & Kingston, A.E. 1984, JPhB, 17, 3321 Dufton, P.L., & Kingston, A.E. 1987, JPhB, 20, 3899 Dufton, P.L., & Kingston, A.E. 1989, MNRAS, 241, 209 Dufton, P.L., & Kingston, A.E. 1991, MNRAS, 248, 827 Dufton, P.L., & Kingston, A.E. 1994, ADNDT, 57, xx Dufton, P.L., Brown, P.J.F., Lennon, D.J., Lynas-Gray, A.E. 1986, MNRAS, 222, 713 Dufton, P.L., Doyle, J.G., Kingston, A.E. 1979, A&A, 78, 318 Dufton, P.L., Hibbert, A., Keenan, F.P., Kingston, A.E., & Dufton, P.L., Hibbert, A., Kingston, A.E., & Doschek, G.A. 1982, ApJ, 257, 338 Dumont, A.M., & Mathez, G. 1981, A&A, 102, 1 Fang, Z., & Kwong, H.S. 1997 ApJ 483, 527 Federman, S.R., & Shipsey, E.J. 1983, ApJ, 269, 791 Fleming, J., Bell, K.L, Hibbert, A., Vaeck, N., Godefroid, M.R. 1996, MNRAS, 279, 1289 Fleming, J., Brage, T., Bell, K.L., Vaeck, N., Flower, D.R. 1976, A&A, 56, 451 Froese Fischer, C. 1983, JPhB, 16, 157 Froese Fischer, C. 1995, ApJ 455, 758 Galavis, M.E., Mendoza, C., & Zeippen, C.J. 1995, A&AS, 111, 347 Garstang, R.H., 1957, Vistas in Astronomy, 1, 268 Garstang, R.H., Robb, W.D., Rountree, S.P. 1978, ApJ, 222, 384 Giles, K. 1981, MNRAS, 195, 63 Haves, M.A., 1986, J Phys B 19, 1853. Hibbert, A., Godefroid, M.R., Froese-Fischer, C. 1995, ApJ, 455, 758 Ho, Y.K., & Henry, R.J.W. 1984, ApJ, 282, 816 Hummer, D.E., & Storey, P.J. MNRAS, 224, 801 Johansson, S., Brage, T., Leckrone, D.S., Nave, G. & Wahlgren, G.M. 1995, ApJ 446, 361 Johnson, C.T., Burke, P.G., Kingston, A.E. 1987, JPhB, 20, 2553 Johnson, C.T., Kingston, A.E., Dufton, P.L. 1986, MNRAS, 220, 155 Kafatos, M., & Lynch, J.P. 1980, ApJS, 42, 611 Kaufman, V., & Sugar, J. 1986, J Phys Chem Ref Dat, 15, 321 Keenan, F.P. Berrington, K.A., Burke, P.G., Dufton, P.L., Keenan, F.P., & Norrington, P.H. 1987 A&A, 181, 370 Keenan, F.P., Harra, L.K., Aggarwal, K.M., Feibelman, W.A. 1992, ApJ, 385, 375 Kingston, A.E. 1986, PhyS 34, 216 Krueger, T.K., & Czyzak, S.J. 1970, Pro Roy Soc Lond, 318, 531 Kwong, V., Fang, Z., Gibbons, T.T., Parkinson, W.H., Smith, P.L. 1993, ApJ, 411, 431 Leep, D., & Gallagher, A. 1976, Phys Rev A, 13, 148 Lennon, D.J. & Burke, V.M. 1991, MNRAS 251, 628 Lennon, D.J. Burke, V.M. 1994, A&AS, 103, 273 Lennon, D.J., Dufton, P.L., Hibbert, A., Kingston, A.E. 1985, ApJ, 294, 200 Lepp, S., & Shull, J.M. 1983, ApJ, 270, 578 Mason, H. 1975, MNRAS 170, 651 McLaughlin, B.M., & Bell, K.L. 1993, ApJ, 408, 753 Mendoza, C. 1982, in Planetary Nebulae, IAU Symp No. 103, ed by D.R. Flower, (D. Reidel: Holland), 143Mendoza, C., & Zeippen, C.J. 1982, MNRAS, 199, 1025 Mendoza, C., & Zeippen, C.J. 1983, MNRAS, 202, 981 Mendoza, C., & Zeippen, C.J. 1987, MNRAS, 224, 7p Mendoza, C., & Zeippen, C.J., 1982, MNRAS, 198, 127 Merkelis, G., Vilkas, M.J., Gaigalas, G., & Kisielius, R. 1994, priviate communication Mohan, M., Hibbert, A., & Kingston, A.E. 1994, ApJ, 434, 389 Muhlethaler, H.P., & Nussbaumer, H. 1976, A&A 48, 109

- Nussbaumer, H. 1986, A&A , 155, 205
- Nussbaumer, H., & Rusca, C. 1979, A&A, 72, 129
- Nussbaumer, H., & Storey, P. J. 1981, AA, 99, 177
- Nussbaumer, H., & Storey, P.J. 1982, A&A, 113, 21
- Nussbaumer, H., & Storey, P.J. 1988, A&A, 200, L25
- Oliva, E., Pasquali, A., & Reconditi, M. 1996, A&A, 305, 210
- Pelan, J., & Berrington, K.A. 1995, A&A Suppl, 110, 209
- Ramsbottom, C.A., & Bell, K.L. 1997, ADNDT, 66, 65
- Ramsbottom, C.A., Bell, K.L., Stafford, R.P. 1996, ADNDT, 63, 57
- Ramsbottom, C.A., Bell., K.L., & Keenan, F.P., 1997, MNRAS 284, 754
- Ramsbottom, C.A., Berrington, K.A., Bell, K.L. 1995, ADNDT, 61, 105
- Ramsbottom, C.A., Berrington, K.A., Hibbert, A., Bell, K.L. 1994, PhyS, 50, 246
- Saha, H.P., & Trefftz, E. 1982, A&A, 116, 224
- Saha, H.P., & Trefftz, E. 1983, SoPh, 87, 233
- Sampson, D.H., & Zhang, H.L. 1998, ApJ, 335, 516
- Saraph, H.E. & Tully, J.A. 1994, A&AS, 107, 29
- Saraph, H.E. 1970, JPhB, 3, 952
- Saraph, H.E., & Storey, P.J. A&AS, 115, 151
- Saraph, H.E., Storey, P.J., & Tully, J.A. 1995, 5th International Colloquium on Atomic Spectra and Oscillator Strengths, ed. by W.-U L. Tchang-Brillet, J.-F. Wyart, C.J. Zeippen, (Meudon: Publications de l'Observaroire de Paris), p.110
- Sawey, P.M.J., and Berrington, K.A., 1993, ADNDT 55, 81
- Seaton, M.S. 1964, Plan Sp Sci 12, 55.
- Sigut, A., & Pradhan, A.K., 1994, J Phys B sub
- Stafford, R.P., Bell, K.L, Hibbert, A. & Wijesundera, W.P., MN 1994 268, 816,
- Stafford, R.P., Hibbert, A., Bell, K.L. 1993, MNRAS, 260, L11
- Stepney and Guilbert, MNRAS 204, 1269 (1983)
- Storey, P.J., Mason, H.E., Saraph, H.E., 1996, A&A, 309, 677
- Sugar, J., & Corliss, C 1985 J Phys Chem Ref Dat, vol 14
- Tayal, S.S. 1997, ApJ 481, 550
- Tayal, S.S., & Henry, R.J.W. 1986, ApJ, 302, 200
- Tayal, S.S., Burke, P.G., Kingston, A.E. 1985, JPhB, 18, 4321
- Tayal, S.S., Henry, R.J.W., Pradhan, A.K. 1987, ApJ, 319, 951
- Tielens, A.G.G., & Hollenbach, D. 1985, ApJ, 291, 722
- Wahlgren, G.M. 1995, ApJ 446, 361
- Wiese, W.L., Fuhr, J.R., Deters, T.M. 1996, J Phys Chem Ref Data, Monograph 7
- Wills, B.J., Wills, D., Netzer, H. 1985, ApJ, 288, 143
- Wills, D., & Netzer, H. 1979, ApJ, 233, 1
- Zeippen, C.J. 1982, MN 198 111
- Zeippen, C.J. 1990, A&A, 229, 248
- Zeippen, C.J., Le Bourlot, J., Butler, K. 1987, A&A, 188, 251
- Zhang, H.L., Graziani, M., Pradhan, A.K. 1994, A&A, 283, 319
- Zygelman, B. & Dalgarno, A. 1987, Phys.Rev.A, 35, 4085
- Zygelman, B., and Dalgarno, A. 1990, ApJ 365, 239

6 PROBLEMS

6.1 Overview

This section describes some of the errors that can cause **Cloudy** to stop. Floating point errors should never occur. Several other internal errors, which the code is designed to catch and then complain about, can occur. Finally, it is possible that the code will stop because of thermal stability problems.

The most important single thing to understand about any calculation is why it stopped, and whether this affects the predictions. This is discussed further in the section *Stopping Criteria* in Part I of this document.

6.2 Thermal stability and temperature convergence

This section describes thermal stability problems, how to identify them, and what to do about them.

6.2.1 Types of thermal maps

Three types of thermal maps, showing the heating or cooling of gas as a function of temperature, can be produced by **Cloudy**. Each is the answer to a different question.

Figure 4 shows the heating and cooling rates as a function of temperature for a photoionized gas in which the electron temperature was varied. This figure was produced by running the test case **map.in**, one of the standard test




Figure 4 A typical cooling function for low density photoionized gas. The cooling and heating rates (erg cm⁻³ s⁻¹) are shown. cooling

cases included in the code distribution. Both the gas density and the flux of ionizing photons was held constant, so only one temperature is meaningful. The **map.in** file uses the **punch map** command to determine heating and cooling rates at a variety of temperatures. This is exactly what the code does to determine the equilibrium temperature, so this plot can be useful to find out why the code ran into problems. This is why the command was introduced. Note that the only valid solution, and the only one with physical meaning, is the one where heating and cooling match – the others are simple bookkeeping exercises.

Gas in collisional equilibrium has a well-defined cooling rate that is only a function of temperature. The sample program **coolcurve.c** (included in the source distribution) does such a calculation, and Figure 5 shows it. Here the electron temperature is set by some physics external to the problem. Each temperature, and the entire ionization solution, is valid for each temperature, under this assumption. The unspecified heat source would have to provide a local heating rate that is equal to the calculated cooling rate for the solution to be time stead.

The third map is the type of thermal stability map shown by Krolik, McKee, and Tarter (1981) and plotted in Figure 6. The program that generated these results is given in the file kmt.c. Here the equilibrium temperature is determined self-consistently for gas over a wide range of densities, but for a single flux of ionizing photons (or equivalently, distance from the central object).





Figure 5 A typical cooling function for low density collisionally ionized gas. coolcurve

6.2.2 No Temperature Convergence

A temperature failure occurs when the heating-cooling balance is not within a certain tolerance, set by the **tolerance** command, after 20 tries. Normally **Cloudy** will punt after an excessive number of temperature failures occur. The limit to the number of failures is reset with the **failures** command, which sets the variable



limfal. The default value is 20. (When **Cloudy** stops because of excessive failures it first produces a map of heating-cooling space to give an indication of where the equilibrium temperature should have been.)

Temperature failures most often occur for temperatures in the range 1 to 4×10^3 K, and 10^5 to 10^6 K. These are ranges where the cooling function permits more that one thermal solution (see, for example, Williams 1967). Figure 4 shows a typical cooling function for gas in photoionization equilibrium.

A peak is reached at a temperature near 10^3 K. This can occur when the finestructure lines are major coolants. At lower temperatures their cooling rate goes up exponentially (as expected), until roughly 10^3 K, when their Boltzmann factors are near unity. Above this temperature their cooling rate is nearly proportional to the Coulomb focusing factor $T^{-1/2}$, and the cooling *decreases* until the temperature is high enough for optical forbidden lines to become important (at roughly 4000 K). A similar phenomenon occurs near the ~ 10^5 to 10^6 K peak in the cooling function.

When failures occur because more than one temperature solution is possible, the reported failures are a physical (not numerical) problem. *Cloudy* will try to deal with this problem by forcing the temperature to values below the peak in the cooling

function. Increasing the number of allowed failures (with the **failures** command) to prevent the code from stopping prematurely is permissible as long as the global energy balance is preserved. A warning will be issued at the end of the calculation if the heating-cooling balance is not preserved.

6.2.3 Thermal Stability

The thermal solution may be unstable when the temperature derivative of the net cooling function (cooling minus heating) is negative (Field 1965). Possibly unstable solutions are indicated by a "u" just before the equilibrium temperature in the zone printout. (The temperature derivative is for isochoric, not isobaric, conditions.) Comments are printed at the end of the calculation if possibly unstable thermal solutions are present in the calculation.

6.2.4 Thermal fronts

Just as an ionization front is a region where the level of ionization changes dramatically over a small scale, a thermal front occurs where the temperature changes dramatically over a small scale. This can be caused by a real physical change of state of the gas such as those that occur near the peaks in the cooling curve. In reality electron conduction may step in as a significant heating/cooling agent, although Cage & Seaton (19xx) argue that this is not important. This could become a problem in the current implementation since electron conduction is not included. There is no obvious solution to this type of jump since it is a physical effect. Jumps can also occur if the grid zoning becomes too coarse for changes in the physical conditions, although the code should be protected against this.

The code will generate a caution or comment if the electron temperature changes discontinuously from one zone to the next.

6.2.5 Map Output

If an excessive number of temperature failures occur (the default limit is 20) then the program stops and produces a map of the heating and cooling as a function of temperature for the last computed zone. The limit to the number of failures allowed before the code punts is reset with the **failures** command. The map is described

 here. The start of the output from the test case map is shown below.

 90.02x map of heating vs cooling. Cloudy punts, Te= 9.254E+03 HTOT= 9.123E-24 CTOT= 9.118E-24 nzone= 1 COOLNG array is 0 4 25 0.340 0 3 5007 0.182 0 3 88 0.075 H FB 0 0.057 S 4 10 0.048 0 3 51 0.042 S 3 9532 0.335 H ff 0 0.022 S 3 33 0.020 Ne 3 15 0.019 Hefb 0 0.015 N 3 57 0.015 Ne 3 3869 0.013 S 3 18 0.013 Ne 5 24 0.010 Ne 5 14 0.009 C 3 1910 0.008 Heff 0 0.007 Si 2 34 0.006 Fe 5 3892 0.006 0 2 3727 0.005 Line heating array follows Te Heat------> Cool-----> dH/dT dC/DT Ne NH HII Helium 1.0000E+01 3.4774E-22 1 1 0.635 4.66195E-26 H FB 0A 0.723 -8.19E-24 1.56E-27 9.1178E-01 1.0000E+00 -0.07 -0.40 -0.24 -1.75 1.0209Fu1 3.44233E-22 1 1 0.635 4.67510E-26 H FB 0A 0.717 -7.74E-24 1.74E-27 9.1491E-01 1.0000E+00 -0.07 -0.41 -0.23 -1.73

The output begins with a listing of the strongest coolants. Then the program steps through increasing temperatures and prints the heating, cooling, and ionization of the gas. From this information it should be possible to determine the temperature where the equilibrium thermal solution should have been. Each solution is completely self-consistent (except that heating and cooling do not balance). Both the local attenuated radiation field and collisional ionization contribute to the ionization

balance at each temperature. All processes contribute to the thermal balance, including collisional ionization. The map is at constant density.

The first column gives the temperature. Columns 2 and 6 give the volume heating and cooling. Both have units erg s⁻¹ cm⁻³. Columns 3 and 4 constitute an indication of the main heating source. Columns 7 and 8 give the label and wavelength of the strongest coolant. Columns 5 and 9 give the fraction of the total heating or cooling due to these agents. Columns 10 and 11 give the heating and cooling derivatives. Columns 12 and 13 give the electron and hydrogen densities (cm⁻³) and the remaining columns give the logs of the hydrogen and helium ionization fractions. The location of the probable thermal solution is indicated by a comment surrounded by dashed lines.

6.3 Floating Point Errors

The code should be compiled and linked with options enabled so that the code will crash on overflow or division by 0, but ignore underflow. *Floating point errors should never occur*. The logic within the code is designed to identify problems, and complain, but not fail. The logic is only as good as the tests they were designed to pass. It is inevitable that circumstances will occur for which the logic now in the code is not sufficient. It is possible that the code will fail when these circumstances occur. I would be grateful for reports of any such failures, since they inevitably identify shortcomings in the code, and lead to its improvement. My email address is gary@cloud9.pa.uky.edu.

6.4 Optical depth convergence problems

The code generally will not converge if it has not done so within ten or so iterations. Convergence problems most commonly occur when the specified column density or thickness is very near a prominent ionization front. In this case very small changes in the physical conditions results in large changes in the optical depths. The code will not have convergence problems if an optical depth is used as a stopping criterion instead.

6.5 Negative Populations of H, He

It is possible that the code will stop because negative level populations were predicted for atoms or ions of hydrogen and helium. This is not supposed to occur, but sometimes happens because of numerical instabilities in the matrix inversion routine. Please send me the input stream and version of *Cloudy*.

6.6 I can't fix it if I don't know its broken.

Machines are growing faster far more rapidly than people are getting smarter. Reliability in the face of complexity is the major challenge to the development of any large-scale computer code. There can be little doubt that **Cloudy** contains bugs.

The code is well tested in many simple limits, and behaves in the correct manner. Simulations of H II regions, planetary nebulae, and other simple objects, are in good agreement with predictions of other photoionization codes (Ferland et al. 1995). Bugs can be discovered by strange behavior in situations where the code has not been well-tested. The discovery of the existence of problems is itself a major challenge. If problems arise or the code crashes then it is likely that a problem has been isolated. I would appreciate learning about such problems since they identify shortcomings which usually lead to improvements in the code (or the documentation). My email address is gary@cloud9.pa.uky.edu.

7 REVISIONS TO Cloudy

7.1 Overview

This section outlines some of the major versions of *Cloudy*, and gives an indication of the direction development will take in the next few years. Its development began in August of 1978, at the Institute of Astronomy, Cambridge, and has been continued at The University of Kentucky, The Ohio State University, and during extended visits to the Joint Institute for Laboratory Astrophysics, the Royal Greenwich Observatory, Cerro Tolodo Interamerican Observatory, and the Canadian Institute for Theoretical Astrophysics. Figure 7 shows the evolution of the code, as indicated by its size as a function of time⁸.

7.2 Cloudy and Moore's Law

Moore's Law is due to Gordon Moore, one of the founders of Intel Corporation. He observed that modern CPU's become about twice as powerful every 18 months. This trend has held true for the past twenty years, shows no sign of failing, and seems to be associated with the control of complexity.

By this standard the growth of **Cloudy** has been conservative, in that it is growing slower and complex on the same timescale. As an example, the Meudon 1985 Meeting planetary nebula test (**parispn.in** in the distributed test cases) has always




Figure 7 The size of the code, indicated by the number of lines of executable Fortran, as a function of time. size

⁸ Before mid-1995 the size was the total number of lines in the distributed source. After 1995 the size only includes the number of lines excluding block datas. The number of lines of block data

taken about one minute to compute.

7.3 Major Past Versions

• 67 August 1987. Hydrogen atom goes to LTE in limit of large electron densities. Many small bugs uncovered as result of careful comparison with Netzer's ION.

- 68 Cambridge, Fall 1987. Development work in progress.
- 69 December 1987. Hydrogen atom goes to LTE in limit of large photon densities. Ferland and Rees (1988).
- 70 September 1988. He II Lyα transfer improved. Improved form of escape probabilities with explicit damping constants. H⁻ and improved free-free heating. Many high excitation metal lines transferred. (Rees, Netzer, and Ferland 1989; Ferland and Persson 1989).
- 71 December 1988. Photon array rewritten, now Compton exchange problem is exact for black bodies with temperatures between 2.7 K and 10^{10} K. He II Ly α radiation pressure included.
- 72 January 1989. Static version, minor bug fixes.
- 73 1989. Major rewrite of helium treatment. Dust changed to two populations, scattering and absorption included. Default radius and thickness increased by ten orders of magnitude. He⁺ goes to LTE. Development work on getting helium to go to LTE. IR power law for default AGN continuum now v^{+2.5} below 100 micron break. No *anumm* array, all continuum one array. table star. *coolr* broken up. Helium to LTE for high electron density.
- 74 1990 January. Hydrogen double precision, many bug fixes. 10 tables among the continua. Kurucz (1989) atmospheres. Improved dust treatment, including photoionization and charge.
- 75 1990, JILA visit. Cosmic abundances changed to Grevesse and Anders. Major bug in constant pressure for HII regions, PNs, etc; did not affect BLR. Calculation of $vf_v(H\beta)$ was incorrect. Molecules at low temperatures. HDEN now $n(H^o) + n(H^+) + n(H^-) + 2n(H_2) + 2n(H_2^+)$. Improved Rayleigh scattering treatment. Dielectronic recombination for sulfur (guess). Many changes in dust; Orion paper (Baldwin et al. 1990). Subordinate lines changed to Hummer's K2 function.
- 76 1990, static version from end of JILA visit.
- 77 through Nov. 1990. optimize option added using Bob Carswell's code. Gaunt factor for brems input spectrum. Reflected continuum predicted. Frequency partition adjusted. X-Ray optical depth now at 0.5 keV. Read in table of points from previous calculation. *opsav* deleted, now single pointer *opsv* for all opacities. Numerical array to 100 MeV. Hummer Lyα escape destruction prob. *tautot* arrays now both in and outward directions. Bound

became very large, thanks to Dima and Katya Verner, so that the total size is roughly 2.5 times the indicated size.

Compton included for all ionization levels. Mean ionization arrays rewritten to make sense. Bug in wind velocity fixed, result now exact. C, O outward diffuse fields changed to OTS.

- 78 through May 91. Continuum escape probability formal H-only opacity. H recombination, cooling over wide range of temp. X-Ray optical depth back to 1 keV. **abundances no dust** no longer changes abundances of depleted elements. OI-Ly β treatment is now six-level atom. Default table AGN changed. Continuum normalization rewritten. Milne relation for diffuse fields of H, all He. Fe K α divided into hot and cold. Beams paper (Ferland, Peterson, Horne, et al. 1992) Many high ionization lines included as OTS and outward ionization sources.
- 79 Summer 91. H molecules completed, C, N, O molecules included. Continuum binning changed for Ca, Fe L-shell ionization potentials. Extensive testing.
- 80 July 91, static version .05. Version 80.06 fixed many small problems discovered by several people. New collision strength for [NeV] put in. This static version ended with 80.09, in January of 1993.
- 81 Late 1991. New collision strengths for [NeV] IR lines (10x larger). Also for [CII], [NIII], and [OIV]. Some are 2x larger. Opacity arrays totally rewritten with eye to Opacity Project data. NIII paper (Ferland 1992).
- 82 Early 1992. X-Ray opacity arrays rewritten. Improved pressure convergence. HJBAR now function. Error in cooling due to collisional ionization of H, He. Sodium and nickel added. /TAU/ array broken up. All heavy element opacities converted to table look-up. Revised collision strengths for NIII lines, improved treatment of atom. NLR abundances deleted, ISM put in their place. Summary comments now driven by subroutine. Cap on 911 OTS field. Note on [FeXI] maser (Ferland 1993). Luminosity command separated into luminosity and intensity commands.
- 83 Autumn 1992, Cambridge and CTIO visits. Hydrogen molecule network completed, Ferland, Fabian, and Johnstone (1993). Collision strengths for fine structure lines changed to Blum and Pradhan 1992, Hollenbach and McKee 89; these changed temperatures for cold ISM by factors of 2. Heavy element molecule network as in Hollenbach and McKee 1989. Code works in fully molecular limit. Kevin Volk's stars (Atlas 91, and Werner models). More accurate treatment of secondary ionization after Voit visit. [OI] lines each include escape prob. Opacities, destruction probabilities, evaluated within all loops, code far more stable, but roughly three times slower. Transferred HeI 2.06 line correctly, after Shields papers.
- 84 1993 Feb 13, Static version following CTIO visit.
- 85 1994, Revisions following Lexington meeting. Outward only now default continuum transport.
- 86 1995, All of first thirty elements are now in code. Photoionization database changed to Dima Verner's *phfit*.

- 87 1995 summer, Map now converges electron densities. Negative populations of OI and FeII atoms solved. Dima's 6k lines included in cooling and radiative acceleration. Total rewrite of *nextdr* logic. Ionization predictor corrector logic completely rewritten. Dima's *phfit* now fitted to all Opacity Project data.
- 88 1995 Fall-winter. Kirk's extensive grids of BLR models run. March 1996 visit to Tel Aviv to compare results with Hagai. Kirk carefully went over atomic data base. Iron recombination changed to Arnaud and Raymond (1992). Default iron abundance down 33%. Entire line data base revised.
- 89 1995 winter-spring. These were mainly beta versions with no major changes, but many fixes to problems.
- 90 1996 June 17, static version, extensive year of debugging. Ferland et al. (C90) paper.
- 91 1977, preparation of C version.
- 92 1998, sabbatical at CITA, revision and debugging of C version.
- 94 1999 December 24, gold version of C code.

7.4 Version 94 versus 90

7.4.1 Commands

The **hydrogen** and **feii** commands have been combined into the **atom** command, which has many options.

7.4.2 Hydrogen

The full hydrogenic isoelectronic sequence is now treated with a single model atom and code base. The model atom can have up to 400 levels.

7.4.3 grains

Parameters used in the treatment of the old-style grains have been revised as per the Weingartner & Draine (2000) paper. The **pgrains** command makes it possible to resolve the grain size distribution function, solving for grain properties and emission as a function of their size. This is described in van Hoof et al. (2000).

7.4.4 Other changes

The code is now ANSI 1989 C, making it especially gcc and Linux friendly.

7.5 Version 90 versus 84

7.5.1 Commands

The **abundances** command now needs 29 numbers by default. A new command "init" allows a commonly used set of commands to be saved as a single file and used by a variety of scripts.

7.5.2 Continuum Transport

Versions 86 and before used a modified version of on-the-spot approximation (OTS) for the Lyman continua of hydrogen and helium. This method was numerically stable and gave results in excellent agreement with Van Blerkom and Hummer (1967). This has been changed to outward-only to obtain better agreement with predictions of Pat Harrington's and Bob Rubin's codes (Ferland et al. 1995). The

OTS code is still in place and will be used if the **diffuse ots** command is entered, but outward-only is the default. The two methods result in temperatures at the illuminated face which can differ by as much as several thousand degrees, but the resulting spectra are surprisingly similar.

7.5.3 Hydrogen

The model hydrogen atom has been generalized to an arbitrary multi-level atom (Ferguson and Ferland 1996). The **hydrogen levels** command is used to specify the number of levels to be used. The collision strengths have been changed to Vriens and Smeets (1980) for levels higher than 3, and Callaway (1994) for collisions with 1,2 and 3.

Predicted infrared line intensities are now correct for all densities and temperatures greater than 10³ K. Versions before 89 used a well l-mixed hydrogen atom, and its predictions were not correct for some infrared lines at low densities.

The routine that computes the free-free gaunt factors has been extended to include the full range the code can handle.

7.5.4 The helium ion

The helium ionization balance at low photon and particle densities, and at high particle densities, has always been exact, and this continues to be the case. There was a problem in the helium ion for high radiation densities, in versions 87 and before. The code used three pseudo levels to represent the levels between 7 and 1000, for H, He, and He⁺. This seemed to work well for the atoms for the cases of high densities, but testing has shown that it did not represent the physics of the high radiation density limit well. The problem is that the pseudo-levels had very large statistical weights, they represented line energies in the far infrared, and had A's appropriate for lower levels. As a result they had very large induced rates when the photon occupation numbers were large, and this affected populations of lower levels. As a result the atom became too ionized - as much as a factor of two for He⁺. The following test illustrates this problem:

```
title helium ionization in high photon density limit
print departure coef
set dr 0
stop zone 1
constant temper 4
hden 11.000
phi(h) 20.750 range 1
stop thickness 11.7
table agn
```

In versions 88 and later no pseudo levels are used for any hydrogen or helium atom or ion.

7.5.5 Heavy elements

The atomic data base, the organization of aspects of the code dealing with storing heavy element ionization, and all the associated routines, have been totally rewritten. The lightest 30 elements are now included. Photoionization data are from Verner et al. (1996), recombination data partially from Verner and Ferland (1996), and roughly 10⁴ lines of the heavy elements have been added (Verner, Verner, and Ferland 1996).

The number of resonance lines has increased by more than an order of magnitude. All resonance lines listed by Verner, Verner, and Ferland (1996) are included. As a

result of these many additional lines the cooling function tends to be larger and smoother.

All lines are now fully transferred, and include pumping by the attenuated incident continuum as a general excitation mechanism. Pumping can be a significant contributor to the formation of weak high excitation lines.

The default solar mixture has been changed to Grevesse and Noel (1993). The biggest change is in the iron abundance. Previous versions had used a higher photospheric abundance. The current version is the 1993 suggested meteoritic abundance.

7.5.6 Free-free, line heating and cooling

These are counted in a different but equivalent manner. Now the *difference* between cooling and heating is used, since this is more numerically stable at high radiation densities. This difference has no physical affect on the predictions, but the printed contributors to the total heating and cooling do appear different.

7.5.7 Excited state photoionization cross sections

OP data are now used. For the excited state of Mg⁺ this is nearly ten times smaller than old screened hydrogenic values. This affects the intensity of Mg II λ 2798 in some BLR calculations.

7.5.8 The O⁺ photoionization cross section

The Reilman and Manson photoionization cross sections, used before version 87,

show a jump in the photoionization cross section at the 2s - 2p edge, and low values above that threshold extending up to the valence electron threshold. Opacity Project cross sections are used in the current version of the code, and these do not show the 2s edge (the OP calculations find that the 2s and 2p electrons are highly correlated). The cross section remains large up to the valence threshold. The difference approaches a factor of two, and this affects high ionization parameter clouds since O⁺ is the dominant opacity for some energies.

Verner, et al. (1996) comment on all other cases where the photoionization cross sections have changed. There are generally atoms and first ions where Opacity Project data are now available.




Figure 8 This figure shows the O^+ photoionization cross sections now used, compared to the Reilman and Manson values, used in version 84 and before. o2photo

7.6 Version 84 versus 80

7.6.1 Commands

In previous versions of the code both luminosity (quantity radiated into 4π sr) and intensity (a surface flux) were specified with the same command. The code decided which was intended by checking the resulting ionization parameter. This method never failed to the best of my knowledge, but, as the code grows more capable of considering ever more extreme cases, there might eventually come a time when it made the wrong decision. All luminosity-intensity commands have now been split up. For instance, the Q(H) command is now two commands, Q(H) (for number of photons radiated into 4π sr) and ϕ (H), for the surface flux. These commands are all discussed in Part I of HAZY.

7.6.2 Mg II 2798

The biggest difference between the two versions is in the predicted intensity of Mg II λ 2798. The intensity of this line is now a factor of two stronger in many models. The new version uses L-shell photoabsorption cross sections from Reilman and Manson (1979), and the older version used inner shell cross sections extrapolated from the table in Weisheit (1974). The cross sections differ by a factor of nearly 3, in the sense that Mg now tends to be more neutral, and Mg II stronger. As a result of the increased cooling by λ 2798 other lines formed in the same region tend to be weaker.

7.6.3 General Results

The following may affect certain specific models, but did not result in changes in any of the "standard" test cases.

The treatment of line-continuum fluorescence in the optically thin limit has been much improved, following Ferland (1992). This can affect hydrogen line emission in clouds that are optically thin in the Lyman continuum.

The treatment of molecules has been vastly improved. The code now goes to the fully molecular (H_2 and CO) limit, and reproduces the Tielens and Hollenbach (1985a, b) PDR results for Orion.

The elements Na and Ni have been added.

7.7 Known Modes for Cloudy 95

- Time dependent model turned off.
- Molecular abundances can become negative for very large CO fractions.
- 2s level of hydrogen not exactly unity in radiation-dominated LTE limit. Induced two-photon emission.

7.8 Making a Revision

7.8.1 *The code*

- Compile code with array bounds checking. Run all test cases.
- Run all test cases on Alpha, HP, Sparc, SGI, and PC.

- Confirm *fabden* stops immediately.
- Verify **aaaa.c** has correct version.
- Summarize changes in c95rev.htm in the web site. Use comments in version.c. This is the primary documentation for changes to the code.
- Create the test case files for each of the platforms. Each lives in a separate subdirectory off the hazy directory. Create a compressed tar file with the name of the platform included. Place this in the ftp site with the script **cp.ftp**.
- Tar all input files into single file with name tests.tar. The command is tar -cvf tests.tar *.in *.for. Compress and copy to ftp.

7.8.2 Printing Hazy

- Compile the line list from lines.c and bring these into HAZY. To do this, go to the programs subdirectory off the hazy main directory, and run do.all to create formatted output file call lines.out for HAZY. Edit the links in HAZY lines.doc to update this.
- Create the list of transferred lines. Compile the line data list for the transferred lines by running the input file punchline.in in the Cloudy main directory. This will create a punch file named LineData.txt. This is the file that will be linked into HAZY. Edit the links in HAZY's file lines.doc to update the list of transferred lines.
- **Update the list of subroutine names.** Do this by running do.all in the include directory. The list of routine names in is the file routines.txt, and this is automatically generated by do.all. This was probably done to update the source, as described in the previous section. Edit links in HAZY routines.doc to update this.
- *Update all comparison tables.* These are in the section starting on page 470. Go over these to confirm that line predictions are OK.
- *Update all test case input scripts.* These are listed in the section beginning on page 487 of this document. Edit the links to manually update them.
- Confirm that all cross-linked variables are ok. Change labels in HAZY headinfo.doc.
- *Summarize changes to the code*. These are listed in **version.for** and should go into the past major revisions in this section of HAZY.

8 COMPARISON CALCULATIONS

8.1 Overview

This section presents comparisons between the current predictions of the code, and results from other independent calculations. The "other" calculations are from the compendium resulting from the Lexington meeting on model nebulae (Ferland et al. 1995).

The scatter among the calculations, as well as the changes that have occurred in the predictions made by **Cloudy**, are in some sense an indication of the stability and reliability of these types of extreme non-LTE calculations. The largest discrepancies between current predictions made by **Cloudy** and the other models from the Meudon meeting (which were computed in 1985) are due to changes which have occurred in the atomic data base between 1985 and the present. In general, the strongest lines are in very good agreement (as they must because of energy conservation) while weak lines (which are very sensitive to changes in the computed temperature and ionization structure) scatter by nearly a factor of two.

8.2 Cool HII Region

This is an HII region ionized by a very cool star. It is one of the Lexington Meeting test cases and is computed with the input script **coolhii.in** in the code's test suite. This is the simplest model since helium is predominantly neutral. The entry *L(total)* comes from the "Stoy" printed entry.

		Mean	Ferland	Harrington	Netzer	Pequignot	Rubin
L(Hβ)	E36	4.90	4.98	4.93	4.85	4.83	4.93
[NII]	6584+	0.87	0.91	0.82	0.97	0.82	0.84
[OII]	3727+	1.21	1.16	1.22	1.32	1.14	1.21
[Nell]	12.8 µ	0.30	0.35	0.29	0.29	0.29	0.29
[SII]	6720+	0.57	0.64	0.55	0.61	0.52	0.52
[SIII]	18.7 µ	0.31	0.27	0.36	0.17	0.37	0.37
[SIII]	34 μ	0.51	0.47	0.60	0.27	0.61	0.62
[SIII]	9532+	0.57	0.48	0.55	0.64	0.60	0.56
L(total)	E36	21.3	21.3	21.7	20.7	21.0	21.8
T(in)		6860	6952	6749	6980	6870	6747
T(H+)		6767	6740	6742	6950	6660	6742
<he+>/<h+></h+></he+>		0.047	0.041	0.044	0.068	0.048	0.034
R(out)	E18	8.96	8.93	8.94	9.00	8.93	9.00

Table 1 HII Region Ionized by Cool Star

Table 2 Cool HII Region vs Cloudy

			0		
		Mean	STD	90.05	94.00
L(Hβ)	E36	4.90	0.06	4.91	5.03
[NII]	6584	0.65	0.05	0.64	0.63
[OII]	3727+	1.21	0.07	1.08	1.04
[Nell]	12.8 μ	0.30	0.03	0.26	0.26
[SII]	6720+	0.57	0.05	0.45	0.44
[SIII]	18.7 μ	0.31	0.09	0.43	0.45
[SIII]	34 µ	0.51	0.15	0.61	0.64
[SIII]	9532	0.43	0.05	0.42	0.42
L(total)	E36	21.29	0.45	21.0	21.7
T(in)		6860	110	7261	7188
T(H+)		6767	108	6600	6530
<he+>/<h+></h+></he+>		0.047	0.01	0.048	0.043
R(out)	E18	8.96	0.04	8.81	8.92

8.3 Paris HII Region

This compares current predictions of the code with those of other participants at the Meudon meeting on photoionization calculations for the case of a simple spherical HII region. The input used to generate this model HII region is contained in the sample input file *parishii.in*.

	8 8							
		Meu	Lex	Ferland	Harrington	Netzer	Pequignot	Rubin
L(Hβ)	E37	2.06	2.04	2.06	2.04	2.02	2.02	2.05
Hel	5876	0.116	0.111	0.109	0.119	0.101	0.116	
CII	2326+	0.17	0.17	0.19	0.17	0.16	0.14	0.18
CIII]	1909+	0.051	0.07	0.059	0.059	0.078	0.065	0.076
[NII]	122 µ		0.031	0.033			0.036	0.031
[NII]	6584+	0.73	0.80	0.88	0.74	0.87	0.78	0.73
[NIII]	57 μ	0.30	0.28	0.27	0.29	0.26	0.30	0.30
[OII]	3727+	2.01	2.20	2.19	2.14	2.3	2.11	2.26
[OIII]	51.8 μ	1.10	1.06	1.04	1.11	0.99	1.08	1.08
[OIII]	88.4 µ	1.20	1.20	1.07	1.28	1.16	1.25	1.26
[OIII]	5007+	2.03	2.09	1.93	1.96	2.29	2.17	2.10
[Nell]	12.8 µ	0.21	0.21	0.23	0.19	0.22	0.20	0.20
[Nelll]	15.5 µ	0.44	0.41	0.43	0.43	0.37	0.42	0.42
[Nelll]	3869+	0.096	0.091	0.103	0.086	0.100	0.079	0.087
[SII]	6720+	0.14	0.18	0.23	0.16	0.22	0.17	0.13
[SIII]	18.7 µ	0.55	0.53	0.48	0.56	0.5	0.55	0.58
[SIII]	34 μ	0.93	0.87	0.82	0.89	0.81	0.88	0.94
[SIII]	9532+	1.25	1.31	1.27	1.23	1.48	1.27	1.30
[SIV]	10.5 µ	0.39	0.38	0.37	0.42	0.36	0.41	0.33
Sum	Sum	11.59	11.87	11.58	11.70	12.17	11.90	12.01
L(total)	E37	24.1	24.4	24.1	24.1	24.8	24.3	24.6
T(in)		7378	7655	7815	7741	7670	7650	7399
T(H+)		7992	8052	8064	8047	8000	8060	8087
He+>/ <h+< td=""><td>></td><td></td><td>0.76</td><td>0.71</td><td>0.77</td><td>0.76</td><td>0.75</td><td>0.83</td></h+<>	>		0.76	0.71	0.77	0.76	0.75	0.83
R(out)	E18	1.45	1.46	1.46	1.46	1.47	1.46	1.46

Table 3 Paris Meeting HII Region

		Table 4	Paris H	III Kegi	on vs C	loudy		
		Mean	STD	76.03	80.06	84.15	90.05	94.00
L(Hβ)	E37	2.04	0.02	2.06	2.04	2.01	2.05	2.02
Hel	5876	0.11	0.01	0.12	0.12	0.12	0.11	0.117
CII]	2326+	0.17	0.02	0.24	0.17	0.17	0.15	0.145
CIII]	1909+	0.07	0.01	0.06	0.11	0.09	0.063	0.060
[NII]	122 µ	0.03	0.00			0.03	0.028	0.028
[NII]	6584	0.60	0.05	0.58	0.60	0.58	0.58	0.57
[NIII]	57 μ	0.28	0.02			0.29	0.289	0.297
[OII]	3727+	2.20	0.08	2.29	2.43	2.31	2.02	1.996
[OIII]	51.8 µ	1.06	0.05	1.15	1.09	1.11	1.22	1.24
[OIII]	88.4 μ	1.20	0.09			1.06	1.11	1.13
[OIII]	5007	1.57	0.11	1.83	1.62	1.50	1.66	1.62
[Nell]	12.8 µ	0.21	0.02	0.22	0.23	0.23	0.18	0.18
[NeIII]	15.5 μ	0.41	0.02	0.47	0.45	0.45	0.28	0.29
[NeIII]	3869	0.0689	0.008	0.098	0.090	0.082	0.063	0.061
[SII]	6720+	0.18	0.04	0.27	0.29	0129	0.126	0.12
[SIII]	18.7 μ	0.53	0.04	0.54	0.53	0.52	0.61	0.61
[SIII]	34 μ	0.87	0.05			0.88	0.85	0.86
[SIII]	9532	0.98	0.07	1.04	1.04	1.05	0.90	0.89
[SIV]	10.5 µ	0.38	0.04	0.09	0.11	0.36	0.53	0.53
Sum	Sum	11.87	0.24	10.0	9.9	11.8	11.9	11.8
L(total)	E37	24.44	0.31	20.9	20.4	23.9	24.5	24.3
T(in)		7655	157			6547	7822	7746
T(H+)		8052	32			7530	7970	7920
<he+>/<</he+>		0.76	0.04			0.75	0.72	0.74
R(out)	E18	1.46	0.00			1.45	1.45	1.46

Table 4 Paris HII Region vs Cloudy

8.4 Blister Hll Region

This is one of the Lexington Meeting test cases, and is meant to be similar to inner regions of the Orion Nebula and is called **blister.in** in the test suite.

		Mean	Ferland	Harrington	Netzer	Pequignot	Rubin
Ι(Ηβ)		4.69	4.59	4.81	4.69	4.67	4.70
Hel	5876	0.12	0.13	0.11	0.12	0.12	
CII	2326+	0.16	0.14	0.20	0.10	0.15	0.23
CII	1335+	0.15	0.17	0.14	0.13	0.16	
CIII]	1909+	0.18	0.22	0.17	0.18	0.15	0.20
[NII]	6584+	0.81	0.58	0.94	0.74	0.90	0.87
[NIII]	57 μ	.033	.035	.033	.033	.032	.034
[OII]	7330+	0.12	0.10	0.13	0.09	0.12	0.14
[OII]	3727+	0.86	0.73	0.98	0.69	0.86	1.04
[OIII]	51.8 μ	0.29	0.31	0.29	0.28	0.28	0.28
[0111]	5007+	4.18	4.74	3.90	4.40	3.90	3.96
[Nell]	12.8 µ	0.34	0.32	0.33	0.35	0.33	0.35
[Nell]	15.5 μ	1.06	1.24	1.07	0.96	1.04	1.00
[NeIII]	3869+	0.34	0.48	0.32	0.35	0.26	0.29
[SIII]	18.7 μ	0.33	0.31	0.34	0.31	0.33	0.35
[SIII]	9532+	1.46	1.41	1.46	1.51	1.42	1.53
[SIV]	10.5 µ	0.51	0.54	0.52	0.51	0.53	0.43
Sum	Sum	10.78	11.32	10.80	10.63	10.46	10.71
l(total)		51.3	52.6	52.4	50.4	49.4	50.3
T(in)		7911	8206	7582	8200	8200	7366
T(H+)		8303	8324	8351	8310	8200	8328
<he+>/<h+></h+></he+>		0.85	0.94	0.78	0.93	0.79	0.84
ΔR	E17	2.99	2.88	3.08	2.93	2.98	3.09

Table 5 Blister HII Region

	Table 6	Blister Hl	I Region	vs Cloudy	
		Mean	STD	90.05	94.00
Ι(Ηβ)		4.69	0.08	4.71	4.64
Hel	5876	0.12	0.01	0.129	0.132
CII]	2326+	0.16	0.05	0.133	0.124
CII	1335+	0.15	0.02	0.179	0.171
CIII]	1909+	0.18	0.03	0.222	0.234
[NII]	6584	0.54	0.09	0.468	0.417
[NIII]	57 μ	0.03	0.00	0.037	0.038
[OII]	7330+	0.12	0.02	0.115	0.113
[OII]	3727+	0.86	0.15	0.737	0.713
[OIII]	51.8 μ	0.29	0.01	0.303	0.309
[OIII]	5007	3.13	0.28	3.74	3.84
[Nell]	12.8 µ	0.34	0.01	0.140	0.141
[Nelll]	15.5 μ	1.06	0.11	0.458	0.462
[Nelll]	3869	0.34	0.08	0.157	0.159
[SIII]	18.7 μ	0.255	0.015	0.301	0.304
[SIII]	9532	1.09	0.037	0.950	.955
[SIV]	10.5 μ	0.51	0.04	0.68	0.68
Sum	Sum	10.78	0.32	10.6	10.7
l(total)		51.29	1.39	50.9	50.2
T(in)		7910	406	8447	8370
T(H+)		8302	59	8325	8330
<he+></he+>	/ <h+></h+>	0.85	0.07	0.895	0.90
ΔR	E17	2.99	0.09	2.99	2.94

Table 6 Blister HII Region vs Cloudy

8.5 Paris Planetary Nebula

This compares current predictions of the code with those of other participants at the Meudon (1985) and Lexington (1993) meetings on photoionization calculations, for the case of ionization by a very hot black body. The input used to generate this model planetary nebula is shown in the sample input section and is called **parispn.in** in the test suite. The model results are very sensitive to the detailed transfer of HeII Ly α ; this line is the dominant heat source across the He⁺⁺ region of the model nebula. The parameters were chosen to be roughly similar to NGC 7027, a very well studied object.

	Tab	ole 7 Par	is Meeting	g Planeta	ary Nebula	a a	
Line		Meudon	Lexington	Ferland	Harrington	Netzer	Pequignot
L(Hβ)	E35	2.60	2.68	2.63	2.68	2.73	2.68
Hell (35)	erg/s	0.87	0.88	0.83	0.88	0.94	0.85
He I	5876	0.11	0.11	0.11	0.10	0.10	0.11
He II	4686	0.33	0.33	0.32	0.33	0.35	0.32
C II]	2326+	0.38	0.33	0.33	0.43	0.27	0.30
C III]	1909+	1.70	1.77	1.82	1.66	1.72	1.87
C IV	1549+	1.64	2.33	2.44	2.05	2.66	2.18
[N II]	6584+	1.44	1.49	1.59	1.45	1.47	1.44
N III]	1749+	0.11	0.12	0.13	0.13	0.11	0.13
[NIII]	57 μ		0.13	0.12	0.13	0.13	0.13
N IV]	1487+	0.12	0.19	0.20	0.15	0.21	0.19
NV	1240+	0.09	0.17	0.18	0.12	0.23	0.15
[O I]	6300+	0.15	0.14	0.15	0.12	0.14	0.14
[O II]	3727+	2.23	2.25	2.23	2.27	2.31	2.18
[O III]	5007+	20.9	20.76	21.1	21.4	19.4	21.1
[O III]	4363	0.16	0.15	0.16	0.16	0.14	0.16
[O III]	52 μ	1.43	1.43	1.42	1.44	1.40	1.46
[O IV]	26 μ	3.62	3.67	3.52	3.98	3.32	3.86
O IV]	1403+	0.13	0.26	0.20	0.23	0.26	0.33
0 V]	1218+	0.09	0.20	0.20	0.11	0.29	0.19
[Ne III]	15.5 μ	2.51	2.78	2.75	2.76	2.80	2.81
[Ne III]	3869+	2.59	2.69	3.33	2.27	2.74	2.44
Ne IV]	2423+	0.56	0.78	0.72	0.74	0.91	0.74
[Ne V]	3426+	0.73	0.67	0.74	0.60	0.73	0.61
[Ne V]	24.2 μ	1.67	0.87	0.94	0.76	0.81	0.99
Mg II	2798+	1.48	1.58	2.33	1.60	1.22	1.17
[Mg IV]	4.5 μ	0.09	0.12	0.12	0.13		0.12
[Si II]	34.8 μ	0.13	0.19	0.16	0.26	0.19	0.17
Si II]	2335+	0.11	0.16	0.15		0.16	0.16
Si III]	1892+	0.20	0.41	0.39	0.32	0.46	0.45
Si IV	1397+	0.15	0.18	0.20	0.15	0.21	0.17
[S II]	6720+	0.39	0.35	0.21	0.45	0.33	0.43
[S III]	18.7 μ	0.49	0.48	0.48	0.49	0.46	0.49
[S III]	9532+	2.09	1.96	2.04	1.89	2.05	1.87
[S IV]	10.5 µ	1.92	1.98	1.92	2.21	1.81	1.98
L(total)	E35	129	137	139	136	135	136
T(in)	E4		1.81	1.83	1.78	1.84	1.78
T(H+)	E4		1.25	1.22	1.21	1.35	1.21
<he+>/<h+></h+></he+>			0.72	0.74	0.74	0.71	0.71
R(out)	E17		4.06	4.04	4.04	4.07	4.07

Table 7 Paris Meeting Planetary Nebula

		r	Table 8	Paris P	lanetary	vs Cloud	dy		
Line		Lexington	STD	74.23	76.03	80.06	84.15	90.05	94.00
L(Hβ)	E35	2.68	0.04	2.57	2.66	2.52	2.34	2.62	2.55
Hell (35)	erg/s	0.88	0.05				0.83	0.83	0.866
He I	5876	0.11	0.01	0.11	0.11	0.11	0.11	0.111	0.108
He II	4686	0.33	0.01	0.29	0.32	0.35	0.36	0.319	0.343
C II]	2326+	0.33	0.07	0.36	0.35	0.37	0.35	0.286	0.289
C III]	1909+	1.77	0.09	1.57	1.48	1.72	1.72	1.88	1.83
CIV	1549+	2.33	0.27	2.24	2.76	2.48	2.19	2.65	2.63
[N II]	6584	1.12	0.05	1.06	1.05	1.08	1.11	0.978	1.00
N III]	1749+	0.12	0.01	0.10	0.08	0.10	0.11	0.123	0.118
[NIII]	57 μ	0.13	0.00				0.12	0.128	0.125
N IV]	1487+	0.19	0.03	0.16	0.12	0.11	0.15	0.242	0.238
NV	1240+	0.17	0.05	0.14	0.09	0.06	0.09	0.175	0.18
[O I]	6300	0.105	0.007	0.11	0.11	0.12	0.12	0.116	0.118
[0 II]	3727+	2.25	0.06	2.24	2.19	2.35	2.40	2.22	2.32
[0 III]	5007	15.6	0.68	15.9	15.8	15.3	15.6	17.0	16.3
[O III]	4363	0.15	0.01	0.14	0.13	0.15	0.16	0.174	0.166
[O III]	52 μ	1.43	0.03	1.40	1.35	1.37	1.35	1.32	1.28
[O IV]	26 µ	3.67	0.30			3.42	3.65	3.42	3.59
O IV]	1403+	0.26	0.05	0.19	0.22	0.11	0.15	0.248	0.25
0 V]	1218+	0.20	0.07	0.17	0.11	0.07	0.11	0.200	0.20
[Ne III]	15.5 μ	2.78	0.03	2.77	2.70	2.67	2.71	1.90	1.87
[Ne III]	3869	2.01	0.35	2.41	2.25	2.43	2.49	2.15	2.09
Ne IV]	2423+	0.78	0.09	0.62	0.51	0.51	0.63	0.823	0.827
[Ne V]	3426	0.50	0.06	0.48	0.40	0,40	0.48	0.589	0.614
[Ne V]	24.2 µ	0.87	0.11	0.24	0.25	1.01	1.04	1.03	1.09
Mg II	2798+	1.58	0.54	0.83	1.82	1.96	2.33	2.26	2.27
[Mg IV]	4.5 μ	0.12	0.00	0.12	0.13	0.14	0.13	0.118	0.12
[Si II]	34.8 µ	0.19	0.04	0.16	0.16	0.16	0.17	0.157	0.16
Si II]	2335+	0.16	0.01	0.15	0.14	0.16	0.18	0.150	0.15
Si III]	1892+	0.41	0.07	0.32	0.42	0.42	0.42	0.526	0.49
Si IV	1397+	0.18	0.03	0.17	0.24	0.22	0.15	0.235	0.223
[S II]	6720+	0.35	0.11	0.38	0.68	0.66	0.36	0.354	0.354
[S III]	18.7 μ	0.48	0.02	0.58	0.71	0.67	0.47	0.467	0.472
[S III]	9532	1.47	0.075	1.27	1.58	1.55	1.48	1.34	1.35
[S IV]	10.5 μ	1.98	0.17	1.64	1.32	1.53	1.78	2.20	2.04
L(total)	E35	137	1.60	117	124	128	121	135	
T(in)	E4	1.81	0.03				1.49	1.828	1.80
T(H+)	E4	1.25	0.07				1.28	1.22	1.22
<he+>/<h+< td=""><td></td><td>0.72</td><td>0.02</td><td></td><td></td><td></td><td>0.72</td><td>0.74</td><td>0.72</td></h+<></he+>		0.72	0.02				0.72	0.74	0.72
R(out)	E17	4.06	0.02				3.90	4.03	3.99

8.6 Paris NLR Model

This compares current predictions of the code with those of other participants at the Meudon meeting on photoionization calculations, for a model similar to the NLR of active nuclei. Results for other codes are from the 1985 Meudon meeting. The input stream is the file *parisnlr.in* in the test suite.

Line		Netzer	Pequignot	Binette	Kraemer	Mean
Нβ	erg/s/cm ²	0.129	0.134	0.124	0.12	0.127±0.006
Нβ	4861	1.00	1.00	1.00	1.00	1.00
Lyα	1216	35.3	33.1	-	24.0	30.8±6.0
He I	5876	0.095	0.098	0.092	0.090	0.094 ± 0.004
He II	4686	0.36	0.32	0.38	0.37	0.358 ± 0.026
C II]	2326	0.96	0.77	1.70	1.06	1.12±0.40
CII	1335	0.14	0.14	0.20	0.08	0.14±0.05
C III]	1909	4.59	4.99	6.50	4.91	5.25±0.85
CIV	1549	7.03	7.20	5.30	7.20	6.68±0.93
[N I]	5200	0.31	0.33	0.82	0.37	0.46±0.24
[N II]	6548	2.68	1.52	1.77	1.63	1.90±0.53
N III]	1749	0.40	0.40	0.43	0.48	0.428±0.038
N IV]	1487	0.45	0.43	0.51	0.48	0.468 ± 0.035
NV	1240	0.32	0.30	0.32	0.28	0.305±0.019
[O I]	63.2 µm	-	0.62	0.14	0.10	0.29±0.29
[O I]	6300	1.32	0.90	1.62	1.04	1.22±0.32
[O II]	7325	0.11	0.094	0.16	0.10	0.116±0.03
[O II]	3727	3.4	2.62	4.41	2.73	3.29±0.82
[O III]	52µm	2.5	2.54	2.31	2.65	2.50±0.14
[O III]	5007	27.36	27.36	23.28	27.76	26.44±2.11
[O III]	4363	0.42	0.41	0.44	0.44	0.428±0.015
O III]	1663	0.97	0.95	0.92	1.01	0.963±0.038
[O IV]	25.9µm	5.69	5.19	5.49	-	5.46 ± 0.25
O IV]	1403	0.53	0.44	0.51	0.66	0.534 ± 0.092
O V]	1218	0.33	0.32	0.45	0.24	0.335±0.086
O VI	1035	0.17	0.17	0.22	0.10	0.165 ± 0.049
[Ne II]	12.8µm	0.28	0.18	0.48	0.13	0.268±0.155
[Ne III]	15.5µm	2.8	2.62	1.83	1.25	2.13±0.72
[Ne III]	3869	2.70	2.59	2.27	1.67	2.31±0.46
Ne IV]	2423	0.82	0.79	1.03	1.12	0.94±0.16
[Ne V]	24.2µm	3.54	2.64	3.54	-	3.24±0.52
[Ne V]	3426	1.17	1.02	1.13	1.05	1.095±0.066
Mg II	2798	1.58	1.43	1.51	1.10	1.40 ± 0.21
Si II	34.8µm	1.73	0.97	0.51	-	1.07±0.62
Si II	2335	0.21	0.17	0.09	-	0.16±0.06
Si III]	1892	0.15	0.19	0.69	0.14	0.29±0.26
Si IV	1397	0.21	0.14	0.02	0.13	0.13±0.08
SII	6720	1.00	0.62	1.29	0.37	0.82 ± 0.41
SII	4070	0.07	0.04	0.078	0.03	0.055 ± 0.023
S III	18.7µm	0.75	0.49	0.68	0.65	0.64±0.11
S III	9532	2.25	1.38	1.73	1.62	1.74 ± 0.37
S IV	10.5µm	1.39	0.73	0.94	1.57	1.16±0.39

Line	10010-10-1	aris NLR Mo Mean	84.15	90.05	94.00
Нβ	erg/s/cm ²	0.127±0.006	0.133	0.136	0.131
Ηβ	4861 1216	1.00	1.00	1.00	1.00
Lyα Lia I		30.8±6.0	32.3	32.3	33.8
He I	5876	0.094±0.004	0.104	0.103	0.100
He II	4686	0.358±0.026	0.351	0.34	0.330
C II]	2326	1.12 ± 0.40	0.766	0.652	0.693
CII	1335	0.14 ± 0.05	0.126	0.141	0.142
C III]	1909	5.25±0.85	5.02	4.64	4.54
C IV	1549	6.68±0.93	8.42	7.47	7.15
[N I]	5200	0.46±0.24	0.14	0.144	0.151
[N II]	6548	1.90±0.53	2.32	2.36	2.50
N III]	1749	0.428 ± 0.038	0.45	0.388	0.376
N IV]	1487	0.468±0.035	0.553	0.544	0.518
N V	1240	0.305±0.019	0.391	0.302	0.272
[O I]	63.2 μm	0.29±0.29	0.36	0.439	0.464
[O I]	6300	1.22±0.32	1.02	1.02	1.08
[0 II]	7325	0.116±0.03	0.111	0.106	0.117
[O II]	3727	3.29±0.82	2.99	2.69	2.97
[O III]	52µm	2.50±0.14	2.34	2.23	2.20
[O III]	5007	26.44±2.11	25.3	26.05	25.4
[O III]	4363	0.428±0.015	0.45	0.441	0.428
O III]	1663	0.963±0.038	1.05	1.03	1.00
[O IV]	25.9µm	5.46±0.25	5.76	5.91	5.79
O IV]	1403	0.534±0.092	0.54	0.517	0.490
OV]	1218	0.335±0.086	0.453	0.292	0.264
O VI	1035	0.165±0.049	0.222	0.142	0.120
[Ne II]	12.8µm	0.268±0.155	0.220	0.168	0.172
[Ne III]	15.5µm	2.13±0.72	3.13	2.12	2.12
[Ne III]	3869	2.31±0.46	3.99	3.22	3.19
Ne IV]	2423	0.94±0.16	1.19	1.16	1.12
[Ne V]	24.2µm	3.24±0.52	2.74	2.69	2.69
[Ne V]	3426	1.095±0.066	1.45	1.25	1.20
Mg II	2798	1.40±0.21	1.76	1.66	1.62
Si II	34.8μm	1.07±0.62	1.12	1.03	1.07
Si II	2335	0.16±0.06	0.218	0.191	0.201
	1892				
Si III] Si IV		0.29±0.26	0.401	0.501	0.474
	1397 6720	0.13±0.08	0.140 1.50	0.215	0.188
SII	6720 4070	0.82±0.41		0.988	0.958
S II S III	4070 18.7μm	0.055 ± 0.023	0.1004 0.625	0.095	0 71 E
	•	0.64±0.11		0.685	0.715
SIII	9532	1.74±0.37	1.92	1.67	1.74
S IV	10.5µm	1.16±0.39	1.73	1.44	1.32

Table 10 Paris NLR Model vs CLOUDY

8.7 Lexington NLR Model

This is the NLR model computed at the 1994 Lexington meeting, and is called **nlr.in** in the test suites.

		Lexington	Binette	Ferland	Netzer	Pequignot	Viegas
I(Hβ)	E0	1.36	1.33	1.31	1.37	1.43	1.34
Lyα	1216	33.70	38.3	32.1	32.4	31.5	34.2
Hel	5876	0.12	0.11	0.13	0.12	0.13	0.13
Hell	4686	0.24	0.25	0.25	0.25	0.23	0.24
Hell	1640	1.62	1.60	1.74	1.53	1.56	1.67
CIII]	1909+	2.90	2.90	2.99	2.87	2.83	2.90
CIV	1549+	3.35	2.70	3.85	3.69	3.17	3.36
[NII]	6584+	2.55	1.40	3.20	3.10	2.67	2.40
NIII]	1749+	0.23	0.24	0.24	0.22	0.22	0.22
NIV]	1487+	0.21	0.20	0.23	0.22	0.21	0.21
[OI]	6300+	1.65	2.20	1.61	1.67	1.31	1.46
[OI]	63 µ	1.12	0.25	1.13		1.44	1.64
[01]	3727+	1.42	1.60	1.44	1.58	1.30	1.20
OIII]	1663+	0.56	0.35	0.63	0.61	0.57	0.63
[OIII]	5007+	33.54	31.4	34.5	33.0	32.8	36.0
[OIII]	4363	0.32	0.30	0.34	0.31	0.30	0.33
OIV	1403+	0.36	0.49	0.30	0.36	0.42	0.25
[Nelll]	15.5 μ	1.89	1.50	2.01	1.94	2.05	1.95
[Nelll]	3869+	2.13	1.90	2.51	2.16	1.72	2.34
[Ne IV]	2423+	0.44	0.52	0.42	0.47	0.41	0.38
[NeV]	3426+	0.52	0.59	0.55	0.53	0.44	0.50
Mgll	2798+	1.84	3.50	1.72	1.23	1.12	1.61
[Sill]	34.8 µ	0.90	1.00	0.96	1.07	0.96	0.52
[SII]	6720+	1.29	2.40	1.01	0.93	0.99	1.10
[SIII]	9532+	1.91	1.60	2.15	2.06	1.67	2.08
[SIII]	18.7 μ	0.49	0.36	0.61	0.57	0.52	0.37
[SIV]	10.5 µ	1.02	0.86	1.24	0.82	0.94	1.22
l(total)	E0	130	131	128	128	131	133
T(in)	E4	1.70	1.71	1.70	1.72	1.68	1.68
T(H+)	E4	1.18		1.24	1.06	1.20	1.23

Table 11 Lexington NLR Model

	Table 12 Lexington NLR vs Cloudy				
		Mean	STD	90.05	94.01
Ι(Ηβ)	E0	1.36	0.05	1.38	1.33
Lyα	1216	33.70	2.76	30.0	31.6
Hel	5876	0.12	0.01	0.128	0.12
Hell	4686	0.24	0.01	0.24	0.24
Hell	1640	1.62	0.09	1.76	1.68
CIII]	1909+	2.90	0.06	2.62	2.55
CIV	1549+	3.35	0.45	3.99	3.78
[NII]	6584+	2.55	0.72	3.02	3.19
NIII]	1749+	0.23	0.01	0.20	0.20
NIV]	1487+	0.21	0.01	0.286	0.27
[OI]	6300+	1.65	0.34	1.63	1.73
[OI]	63 µ	1.12	0.61	0.77	0.79
[OII]	3727+	1.42	0.17	1.33	1.45
OIII]	1663+	0.56	0.12	0.62	0.60
[0]]]	5007+	33.54	1.76	34.4	33.4
[OIII]	4363	0.32	0.02	0.327	0.32
OIV	1403+	0.36	0.09	0.35	0.38
[Nelll]	15.5 μ	1.89	0.22	1.43	1.44
[Nelll]	3869+	2.13	0.32	1.96	1.98
[Ne IV]	2423+	0.44	0.06	0.47	0.45
[NeV]	3426+	0.52	0.06	0.60	0.60
MgII	2798+	1.84	0.96	1.48	1.48
[Sill]	34.8 µ	0.90	0.22	0.87	0.90
[SII]	6720+	1.29	0.63	0.91	0.91
[SIII]	9532+	1.91	0.26	2.02	2.04
[SIII]	18.7 μ	0.49	0.12	0.68	0.72
[SIV]	10.5 μ	1.02	0.20	1.14	1.06
l(total)	•	130	1.86	125	125
`T(in) ´	E4	1.70	0.02	1.7	1.69
T(H+)	E4	1.18	0.08	1.24	1.23

Table 12 Lexington NLR vs Cloudy

8.8 The DQ Her Shell

This is more or less the model of the DQ Her nebula proposed by Ferland et al. (1984). The input stream for this model is called **dqher.in** in the test suite. The big difference between C90 and previous versions is in the intensity of H β predicted. The code no longer assumes case B when the temperature is too low to do the matrix solution. The nebula is optically thin to many Lyman lines and these escape, robbing

Table 13 The DQ Her Shell					
Line		80.09	84.15	90.05	94.00
Ηβ (+30)	4861	1.65	1.62	0.879	1.07
Totl	4861			0.649	0.883
Case B	4861			1.00	1.00
Lyα	1216	40.9	20.7	20.6	27.9
He I	5876	0.786	0.315	0.246	0.274
He II	4686	0.166	0.085	0.0877	0.052
CII	158µ	0.777	0.62	0.092	0.102
CII	1335		0.062	0.0195	0.021
[N I]	5200	0.144		0.078	0.087
[N II]	122µ	7.85	3.43	3.22	3.58
[N III]	57μ	5.30	3.78	9.20	10.2
[O II]	3727	1.16	0.304	0.294	0.325
[O III]	88µ	12.1	6.36	4.95	5.45
[O III]	52μ	12.4	6.60	5.51	6.06
[Si II]	35μ	0.586	0.79	0.83	0.92
[S III]	34µ	0.114	0.167	0.230	0.258
[Fe II]		0.121	0.187	0.137	0.151
<te></te>	Κ		643	862	853

Table 13 The DO Her Shell

flux from H β .

8.9 The Kwan and Krolik Standard Model

Table 14 gives the spectrum of the Kwan and Krolik (1981) standard model, called **kk.in** in the test suite.

Table 14. The Kwan and Krolik Standard Model.					
Line	KK81	80	84.15	90.05	94.00
Lyα 1216	5.512+7 ⁹	5.78	5.59	6.03	5.98
Lyα 1216	100	100	100	100	100
Ηβ 4861	10.3	6.37	6.05	5.85	5.55
Ηα 6563	42.8	19.9	19.56	24.4	17.2
BaC	47.0	38.8	39.5	59.6	58.5
PaC	30.7	20.2	20.0	37.4	34.1
2s-1s	3.3	2.35	3.19	2.70	2.95
free-free	29.0	25.3	23.3	40.5	39.2
He I 10830	4.4	2.86	2.84	3.30	2.96
He I 5876	0.9	0.697	0.845	0.68	0.64
He II 4686	-	0.365	0.316	0.29	0.27
He II 1640	2.5	3.17	2.77	2.49	2.29
C II] 2326	3.3	2.70	0.686	3.17	3.70
C II 1335	-	0.661	0.611	0.68	0.71
C III 977	5.1	7.53	6.52	5.58	5.47
C III] 1909	13.0	20.7	15.9	13.5	13.2
C IV 1549	67.0	95.7	76.3	70.2	68.1
N III] 1750	1.5	4.26	4.95	4.23	4.17
N III 990	-	0.324	1.35	1.40	1.35
N IV] 1486	5.8	4.88	5.24	5.27	5.13
N V 1240	8.4	3.61	4.95	3.78	3.72
O I 1304	6.9	4.01	3.93	5.13	8.21
O III] 1663	9.5	18.4	18.3	18.6	18.3
O IV] 1402	5.2	4.81	5.84	6.20	6.04
O V] 1218	6.7	3.07	4.63	3.23	3.15
O VI 1034	15.0	2.85	4.71	2.92	2.85
Mg II 2798	18.0	17.6	9.71	30.0	33.8
Si III] 1892	-	14.6	15.7	11.6	12.2
Si IV 1397	5.5	14.2	9.77	8.54	7.60
Fe II	23.9	9.58	11.3	13.6	15.5
Fe Ka		2.11	1.74	1.78	1.81

⁹Line intensity in erg s⁻¹ cm⁻². The entries which follow are relative to a scale where Ly α =100.

8.10 Rees, Netzer, and Ferland, low density

This is the lower density model listed in Table 1 of Rees et al. (1989). It is **rnfa.in** in the sample input streams.

Table 15 Rees, Netzer, and Ferland, low density case.				
	1989	90.05	94.00	lon
log n = 10				
Lyα	1.000	100	100	1.000
Ηβ	0.030	2.76	3.18	0.026
Ηα	0.197	13.0	12.1	0.180
Ρα	0.022	1.45	1.43	0.020
Ba C	0.127	13.3	12.1	0.125
Pa C	0.052	5.01	4.54	0.025
ff	0.082	8.38	8.24	0.089
Hel 5876	0.007	0.656	0.584	0.008
Hell 4686	0.005	0.39	0.392	0.004
Hell 1640	0.039	3.44	3.38	0.034
CII 1335	0.009	1.04	1.00	0.008
CII] 2326	0.014	1.05	0.96	0.011
CIII 977	0.030	2.59	2.52	0.035
CIII] 1909	0.106	9.91	9.66	0.103
CIV 1549	0.424	44.8	42.9	0.453
NIII] 1750	0.012	1.31	1.26	0.014
NIV] 1486	0.009	1.19	1.14	0.009
NV 1240	0.002	0.235	0.22	0.003
OI 1304	0.033	0.91	0.915	0.013
OI 8446	0.005	0.0579	0.064	0.005
OIII] 1663	0.055	6.17	5.92	0.060
OIV] 1402	0.011	1.41	1.35	0.017
Mg II 2798	0.076	19.2	18.4	0.160
Silll 1207	0.012	1.04	1.05	0.013
Silll] 1892	0.085	8.56	8.51	0.090
SilV 1397	0.048	4.47	4.10	0.044
Fell	0	1.79	2.77	0.052
Κα	0	0.098	0.102	0.001
sum	2.492	254.2169	248.787	2.602
Lyα	4.85E+07	4.54+07	4.42+7	4.55+07
l(total)	6.03E+07	1.15+8	1.09+8	6.02+07

8.11 Rees, Netzer, and Ferland, high density

This is the higher density model listed in Table 1 of Rees et al. (1989). It is **rnfb.in** in the sample input streams.

Table 16 Rees, Netzer, and Ferland, high density case.				
	1989	90.05	94.00	lon
	log n = 12			
Lyα	1.000	100	100	1.000
Hβ	0.063	1.99	1.66	0.054
· Hα	0.175	5.87	3.00	0.134
Ρα	0.009	0.45	0.22	0.008
Ba C	2.938	230	234	2.680
Pa C	1.313	104	104	0.590
ff	1.563	114	120	1.300
Hel 5876	0.038	3.13	3.42	0.032
Hell 4686	0.030	2.78	2.76	0.017
Hell 1640	0.266	17.5	18.3	0.140
CII 1335	0.082	4.67	4.84	0.099
CII] 2326	0.003	0.14	0.18	0.002
CIII 977	0.225	9.95	11.5	0.150
CIII] 1909	0.023	0.85	0.92	0.014
CIV 1549	0.938	55.4	62.7	0.860
NIII] 1750	0.006	0.26	0.28	0.004
NIV] 1486	0.006	0.35	0.36	0.004
NV 1240	0.031	2.15	2.34	0.021
OI 1304	0.033	2.01	1.98	0.028
OI 8446	0.006	0.26	0.27	0.005
OIII] 1663	0.039	1.77	1.90	0.024
OIII 835	0.041	1.90	0.043	0.002
OIV] 1402	0.023	1.62	1.69	0.020
Mg II 2798	0.088	12.6	13.6	0.170
Silll 1207	0.088	4.46	4.70	0.068
Silll] 1892	0.113	3.37	3.72	0.061
SiIV 1397	0.300	17.5	18.9	0.230
Fell		0.77	0.99	0.015
Κα		0.38	0.42	0.003
sum	8.188	700.13	718.693	7.521
Lyα	8.54+08	1.17+09	1.08+09	1.08+09
l(total)	7.00+09	8.19+09	7.74+09	7.04+09

9 SAMPLE INPUT

In versions 90 and before this section listed all test cases and discussed the motivations for each. Now each file is self-descriptive, and should be consulted to use as examples and understand how the code is run. Column 1 of Table 17 lists the single-model test cases and the second column gives a brief description of the test.

These tests are included in the distribution, and are the input streams that are used to exercise **Cloudy** under both standard and extreme conditions.

Perl script to Run all models

Concept of assert commands

Perl script to check results

Table 17 The single-model standard test cases			
albedo.in measure rayleigh scattering of Lya			
amoeba.in find density and temperature from HST FOS spectra of [NII] in orion			
atlas.in Model of a Compact HII Region			
bigh.in bigh case B with largest possible H atom			
blister.in conditions similar to Orion nebula blister			
blr89.in final F+P 1989 BLR model table 3			
blr92.in standard blr cloud in Ferland et al. 1992			
brems.in generate continuum due to hot ism in high Z,z starburst			
casea.in case A			
caseblowt.in log density case B, T=5000, log n=2			
casebn2.in log density case B, T=5000, log n=2			
casebn8.in casebn8 high density case B			
casec.in case C			
comphi.in compton exchange in high temper limit			
complo.in compton exchange near low temperature limit			
compton.in Compton limit, test continuum partition			
conserv.in test that energy is conserved			
coolhii.in cool HII region model from Lexington Meeting			
coronal.in model of active region of solar corona			
cosmicray.in background cosmic ray ionization by suprathermal electrons only			
costar.in test costar continuum			
dqher.in (roughly) Ferland et al. DQ Her model			
eden.in Martin Gaskell's funny model			
fe2.in test feii			
feii.in thermal equilibrium of FeII in STE limit			
feiihin.in test feii			
feiihirad.in feii in case of high radiation density limit			
fileopac1.in part of test of stored file opacities			
fileopac2.in part of test using on-the-fly file opacities			
fluc.in Paris meeting Planetary nebula with density fluctuations			
grainlte.in check that grains equilibriate at correct temp in ste limit			
grains.in test all grain species temperature			
hcexc.in test collisional excitation only, very high density to force H to LTE			
hcion.in test collisional ionization only, no excitation, should be in lte			
hemis.in test hydrogen atom continuous emissivity, used for plot in hazy			
hheonlyotspp.in plane parallel conservation and hydrogenic emission for pure hydrogen			
hheonlyotssp.in pherical conservation and hydrogenic emission for hydrogen and helium			
hheonlyoutpp.in plane parallel conservation and hydrogenic emission for pure hydrogen			
hheonlyoutppff.in plane parallel filling factor for pure hydrogen			

hheonlyoutsp.in	spherical conservation and hydrogenic emission for hydrogen and helium
highn.in	high electron density approach to lte
honlyotsopen.in	test ots, inward fractions for pure hydrogen, open geo, filling factor
honlyotspp.in	plane parallel conservation and hydrogenic emission for pure hydrogen
honlyotssp.in	spherical conservation and hydrogenic emission for pure hydrogen
honlyoutopen.in	test inward fractions, open geo, filling factor, 2-photon emission
honlyoutpp.in	plane parallel conservation and hydrogenic emission for pure hydrogen
honlyoutsp.in	spherical conservation and hydrogenic emission for pure hydrogen
induc.in	constant temper black body limit from Ferland and Rees 1988
induchhe.in	half H, He gas with induced BF processes dominate
ism.in	interstellar cloud irradiated by ism background
kk.in	Kwan+Krolik Ap.J. 250, 478 BLR model
lalpha.in	Ly alpha forest cloud
laser1.in	test of H ionization in optically thin limit
laser2.in	test of H and HeI ionization in optically thin limit
laser3.in	test of H, HeI, and HeII ionization in optically thin limit
level2.in	test dominant level2 lines
liner.in	liner model
lowd0.in	test low density limit, this and lowdm6 should get same results
lowden.in	test optically thin model that extends to very low densities
lowdm6.in	low density test
lte.in	thermal equil black body limit from Ferland and Rees 1988
ltemetl.in	thermodynamic equilibrium with metals
map.in	map of heating vs cooling
newpdr.in	Tielens and Hollenbach pdr model, Table 2, paper b
nlr.in	Hagai's nlr model for Lexington Meeting
nova.in	dense nova shell
oldblr.in	Netzer and Ferland PASP 96, 593, Table 1
optimal.in	find density and temperature from HST FOS spectra of [NII] in orion
parishii.in	"New" Paris meeting HII region
parisnlr.in	paris meeting NLR model
parispn.in	parispn.in Meudon Planetary nebula
pdr.in	Tielens and Hollenbach pdr model, Table 2, paper b
pgrains.in	test all grain species temperature
pnots.in	Paris meeting Planetary nebula with ots
powell.in	find density and temperature from HST FOS spectra of [NII] in orion
primal.in	primordial abundances
qheat.in	grain quantum heating test
rauch.in	very hot PN model
rnfa.in	table 1 of Rees et al. ApJ 347, 648
rnfb.in	table 1 of Rees et al. ApJ 347, 648
sqrden.in	test with density falling as R ⁻ -2, and filling factor
strom.in	check pure hydrogen Stromgren sphere
subplex.in	find density and temperature from HST FOS spectra of [NII] in orion
test.in	this runs the standard, one command, test, which contains many asserts
	•
transpunch.in	first of transpunch/transread pair, punch continuum
transread.in	first of transpunch/transread pair, punch continuum
vbhum.in	test against Van Blerkom and Hummer, fig 4
werner.in	test run with Werner stellar atmosphere
wind.in	test of equations of motion in a wind
z3.in	redshift 1000 recombination epoch

10 REFERENCES

- Abbott, D. C. 1982, ApJ, 259, 282
- Adams, T. 1972, ApJ, 174, 439
- Aldrovandi, S., & Pequignot, D. 1972, A&A, 17, 88
- Aldrovandi, S., & Pequignot, D. 1974, Revista Brasileira de Fisica, 4, 491
- Ali, B., Blum, R. D., Bumgardner, T. E., Cranmer, S. R., Ferland, G. J., Haefner, R. I., & Tiede, G. P. 1991, PASP, 103, 1182
- Allen, C. W. 1976, Astrophysical Quantities, Third Edition (London: Athlone Press).
- Aller, L. H. 1984, in Physics of Thermal Gaseous Nebulae, (Dordrecht: Reidel).
- Aller, L. H., & Czyzak, S. J. 1983, ApJS, 51, 211.
- Arimoto, N., & Yoshii, Y. 1987, A&A, 173, 23
- Armour, Mary-Helen, Ballantyne, D.R., Ferland, G.J., Karr, J., & Martin, P.G., 1999, PASP 111, 1251-1257
- Arnaud, M., & Raymond, J. 1992, ApJ, 398, 394
- Arnaud, M., & Rothenflug, R. 1985, A&AS, 60, 425
- Avni, Y., & Tananbaum, H. 1986, ApJ, 305, 83
- Avni, Y., Worrall, D. M., & Morgan, W. A. ApJ, 1995, 454, 673
- Avrett, E. H., & Loeser, R. 1988, ApJ, 331, 211
- Bahcall, J.H., & Kozlovsky, B.-Z. 1969, ApJ, 155, 1077
- Bajtlik, S., Duncan, R. C., & Ostriker, J. P. 1988, ApJ, 327, 570
- Balbus, S. A., & McKee, C. F. 1982, ApJ, 252, 529
- Baldwin, J., Ferland, G. J., Martin, P. G., Corbin, M., Cota, S., Peterson, B. M., & Slettebak, A. 1991, ApJ, 374, 580
- Baldwin J. A., Ferland, G. J., Korista, K. T., Carswell, R., Hamann, F., Phillips, M., Verner, D., Wilkes, B., & Williams, R. E. 1996, ApJ, 461, 683
- Baldwin, J. A., Ferland, G. J., Korista K. T., & Verner, D. 1995, ApJ, 455, L119
- Baldwin, J.A., Verner, E.M., Verner, D.A., Ferland, G.J., Martin, P.G., Korista, K.T., & Rubin, R. H., 2000, ApJS, 129, 229-246
- Baldwin, J., Wampler, J., & Gaskell, C. M. 1989, ApJ, 338, 630
- Balick, B., Gammon, R. H., & Hjellming, R. 1974 PASP, 86, 616
- Ballantyne, D.R., Ferland, G.J., & Martin, P.G., 2000, ApJ 536, 773-777
- Bässgen, G., Bässgen, M., & Grewing, M. 1988, A&A, 200, 51
- Bates, D. R., Kingston, A. E., & McWhirter, R. W. P. 1962, Proc R Soc, A267, 297
- Bechtold, J., Weymann, R. J., Lin, Z., & Malkan, M. A. 1987, ApJ, 315, 180
- Bell, K. L., Kingston, A. E., & McIlveen, W. A. 1975, J. Phys. B 8, 358
- Berger, M. J., & Seltzer, S. M. 1965, NASA SP-3012
- Bergeron, J., & Collin-Souffrin, S. 1971, A&A, 14, 167
- Berrington, K., & Pelan, A. 1995, A&AS, 114, 367
- Bertoldi, F., & Draine, B. T. 1996, ApJ, 458, 222
- Bethe, H. 1930, Ann. Phys. 5, 325
- Bica, E. 1988, A&A, 195, 76
- Bieniek, R. J., & Dalgarno, A. 1979, ApJ, 228, 635
- Binette, L., Prieto, A., Szuszkiewicz, E., & Zheng, W. 1989, ApJ, 343, 135
- Black, J. H. 1978, ApJ, 222, 125
- Black, J. H. 1987, in *Interstellar Processes*, ed. D.J. Hollenbach & H.A. Thronson, (Dordrecht: Reidel), p 731
- Bohm, D., & Aller, L. H. 1947, ApJ, 105, 131
- Bonihala, J. R. M., Ferch, R., Salpeter, E. E., Slater, G., & Noerdlinger, P. 1979, ApJ, 233, 649
- Borkowski, K. J., & Harrington, J. P. 1991, ApJ, 379, 168
- Bottorff, M., Lamothe, J., Momjian E., Verner, E., Vinkovic, D. & Ferland, G. 1998 PASP, 110, 1040
- Bottorff, M. C., Ferland, & Gary J., 2000, MNRAS 316, 103-106
- Bottorff, Mark, Ferland, Gary, Baldwin, Jack, & Korista, Kirk, 2000, ApJ, 542, 644-654
- Boyd, R., & Ferland, G.J. 1987, ApJ, 318, L21

- Bowen, I. S. 1960, ApJ, 132, 1
- Bregman, J. D., Allamandola, L. J., Tielens, A. G. G. M., Geballe, T. R., & Witteborn, F. C. 1989, ApJ, 344, 791
- Broad, J. T., & Reinhardt, W. P. 1976, Phys Rev A 14, 2159
- Brocklehurst, M., 1970, MNRAS, 148, 417
- Brocklehurst, M., 1972, MNRAS, 157, 211
- Brown, R. L., & Mathews, W. G. 1970, ApJ, 160, 939
- Burgess, A. 1965, ApJ, 141, 1588
- Burgess, A., & Summers, H. P. 1969, ApJ, 157, 1007
- Burgess, A., & Summers, H. P. 1976, MNRAS, 174, 345
- Butler, S. E., Bender, C. F., & Dalgarno, A. 1979, ApJ, 230, L59
- Butler, S. E., & Dalgarno, A. 1979, ApJ, 234, 765
- Butler, S. E., Heil, T. G., & Dalgarno, A. 1980, ApJ, 241, 442
- Butler, S. E., & Dalgarno, A. 1980, ApJ, 241, 838
- Callaway, J. 1994, At Dat Nuc Dat Tab 57, 9
- Cameron, A.G.W. 1982, in *Essays in Nuclear Astrophysics*, ed CA Barnes, DD Clayton, & DN Schramm, (Cambridge: Cambridge Univ Press)
- Canfield, R. C., & Puetter, R. C. 1980, ApJ, 236, L7
- Cardelli, J. A. 1994, Science 264, 209
- Cardelli, J. A., et al. 1991, ApJ, 377, L57
- Carswell R. F. & Ferland, G. J. 1988, MNRAS, 235, 1121
- Castor, J. I. 1970, MNRAS, 149, 111
- Chaffee, F. H., & White, R. E. 1982, ApJS, 50, 169
- Chamerlain, J.W., 1956, ApJ, 124, 390
- Chan, E. S., Avrett, E. H., & Loeser, R. 1991, A&A, 247, 580
- Chapman, R. D., & Henry, R. J. W. 1971, ApJ, 168, 169
- Chidichimo, M. C. 1981, J. Phys. B., 14, 4149
- Clavel, J., & Santos-Lleo, M. 1990, A&A, 230, 3
- Clegg, R. E. S. 1987, MNRAS, 229, 31p
- Clegg, R. E. S., & Harrington, J. P. 1989, MNRAS, 239, 869
- Cohen, E. R., & Taylor, B. N. 1987, Rev Mod Phys 57, 1121
- Cota, S. A. 1987, Ph.D. Thesis, OSU
- Cota, S. A., & Ferland, G. J. 1988, ApJ, 326, 889
- Cowie, L. L., & Songaila, A. 1986, ARA&A 24, 499
- CrinkLaw, G., Federman, S. R., & Joseph, C. L. 1994, ApJ, 424, 748
- Crosas, M., & Weisheit, J.C. 1993, MNRAS, 262, 359
- Cruddace, R., Paresce, F., Bowyer, S., & Lampton, M. 1974, ApJ, 187, 497
- Dalgarno, A., & Kingston, A. E. 1963, Observatory, 83, 39
- Dalgarno, A., & McCray, R. A. 1973, ApJ, 181, 95
- Dalgarno, A., & Roberge, W. G. 1979, ApJ, 233, L25
- Dalgarno, A., Yan, Min, & Liu, Weihong 1999, ApJS, 125, 237
- Davidson, K. 1972, ApJ, 171, 213
- Davidson, K. 1975, ApJ, 195, 285
- Davidson, K. 1977, ApJ, 218, 20
- Davidson, K., & Netzer, H. 1979, Rep. Prog. in Physics 51, 715
- Davidson, K., & Fesen, R.A. 1985, ARA&A, 23, 119
- Desert, F.-X., Boulanger, F., & Puget, J. L. 1990, A&A, 237, 215
- Dove, J. E., Rush, A., Cribb, P., & Martin, P. G. 1987, ApJ, 318, 379
- Dove, J. E., & Mandy, M. E. 1986, ApJ, 311, L93
- Draine, B. T. 1978, ApJS, 36, 595
- Draine, B. T., & Lee, H. M. 1984, ApJ, 285, 89
- Draine, B. T., & Salpeter, E. E. 1979, ApJ, 231, 77
- Draine, B. T., & Sultin, B. 1987, ApJ, 320, 803
- Drake, S. A., & Ulrich, R.K. 1980, ApJS, 42, 351
- Elitzur, M. 1982, Rev. Mod. Phys 54, 1125

- Elitzur, M. 1984, ApJ, 280, 653
- Elitzur, M, 1992, Astronomical Masers, (Dordrecht: Kluwer)
- Elitzur, M., Ferland, G. J., Mathews, W. G., & Shields, G. 1983, ApJ, 272, L55
- Elitzur, M., & Ferland, G. J. 1986, ApJ, 305, 35
- Elvis, M. et al. 1994, ApJS, 95, 1
- Fabian, A. C., Pringle, J. E., & Rees M. J. 1976, MNRAS, 175, 43
- Federman, S. R., et al. 1993, ApJ, 413, L51
- Ferguson, J., Ferland, G. J., & A. K. Pradhan, 1995, ApJ, 438, L55
- Ferguson, J., & Ferland, G.J. 1997, ApJ, 479, 363
- Ferguson, J. W., Korista, K. T., Baldwin, J. A., & Ferland, G. J. 1997, ApJ, 487, 122
- Ferguson, Jason W., Korista, Kirk. T., and Ferland, Gary J., 1997, ApJS 110, 287-297.
- Ferland, G. J. 1977, ApJ, 212, L21
- Ferland, G. J. 1979, MNRAS, 188, 669
- Ferland, G. J. 1980a, MNRAS, 191, 243
- Ferland, G. J. 1980b, BAAS, 12, 853
- Ferland, G. J. 1980c, PASP, 92, 596
- Ferland, G. J. 1986, PASP, 98, 549
- Ferland, G. J. 1986, ApJ, 310, L67
- Ferland, G. J. 1992, ApJ, 389, L63
- Ferland, G. J. 1993, ApJS, 88, 49
- Ferland, G. J. 1999, PASP, 111, 1524
- Ferland, G. J. 1999a, in ASP 162, Quasars and Cosmology, ed G Ferland & J Baldwin
- Ferland, G. J., 1999b, ApJ 512 247-249
- Ferland, G. J., Baldwin J. A., Korista, K. T., Hamann, F., Carswell, R., Phillips, M., Wilkes, B., & Williams, R. E. 1996, ApJ, 461, 683
- Ferland, G., Binette, L., Contini, M., Harrington, J., Kallman, T., Netzer, H., Pequignot, D., Raymond, J., Rubin, R., Shields, G., Sutherland, R., & Viegas, S. 1995, in *The Analysis of Emission Lines*, Space Telescope Science institute Symposium Series, R. Williams & M. Livio, editors (Cambridge: Cambridge University Press)
- Ferland, G. J., & Elitzur, M. 1984, ApJ, 285, L11
- Ferland, G. J., Fabian, A. C., & Johnstone, R.M. 1994, MNRAS, 266, 399
- Ferland, G. J., Korista, K.T. & Peterson, B.M. 1990, ApJ, 363, L21
- Ferland, G. J., Korista, K.T., Verner, D. A., & Dalgarno, A. 1997, ApJ, 481, L115
- Ferland, G. J. Korista, K.T. Verner, D.A. Ferguson, J.W. Kingdon, J.B. Verner, & E.M. 1998, PASP, 110, 761
- Ferland, G. J., Lambert, D. L., Netzer, H., Hall, D. N. B., & Ridgway, S. T. 1979a, ApJ, 227, 489
- Ferland, G. J., Lambert, D. L., Slovak, M., Shields, G. A., & McCall, M. 1982, ApJ, 260, 794
- Ferland, G. J., & Mushotzky, R. F. 1982, ApJ, 262, 564
- Ferland, G. J., & Mushotzky, R. F. 1984, ApJ, 286, 42
- Ferland, G. J., & Netzer, H. 1979, ApJ, 229, 274
- Ferland, G. J., & Netzer, H. 1983, ApJ, 264, 105
- Ferland, G. J., Netzer, H., & Shields, G. A. 1979, ApJ, 232, 382
- Ferland, G. J., Peterson, B. M., Horne, K., Welsh, W. F., & Nahar, S. N. 1992, ApJ, 387, 95
- Ferland, G. J., & Persson, S. E. 1989, ApJ, 347, 656
- Ferland, G. J., & Rees, M. J. 1988, ApJ, 332, 141
- Ferland, G. J., & Shields, G. A. 1978, ApJ, 226, 172
- Ferland, G. J., & Shields, G. A. 1985, in *Astrophysics of Active Galaxies & Quasi-stellar Objects*, J.S. Miller, Ed.
- Ferland, G. J., & Truran, J. W. 1981, ApJ, 244, 1022
- Ferland, G. J., Williams, R. E., Lambert, D. L., Shields, G. A., Slovak, M., Gondhalekar, P. M., & Truran, J. W. 1984, ApJ, 281, 194
- Field, G. B. 1965, ApJ, 142, 431
- Francis, P. J. 1993, ApJ, 407, 519
- Gaetz, T. J., & Salpeter, E. E. 1983, ApJS, 52, 155
- Garstang, R.H. 1958, MNRAS, 118, 57

- Gavrila, M. 1967, Phys Rev 163, 147, also JILA Report #86, Sept 19, 1966
- Ginzburg, V. I., & Syrovatskii, S. I. 1964, The Origin of Cosmic Rays, (Oxford: Pergamon)
- Gould, R. S. 1978, ApJ, 219, 250
- Gredel, R., Lepp, S., & Dalgarno, A. 1987, ApJ, 323, L137
- Gredel, R., Lepp, S., Dalgarno, A., & Herbst, E. 1989, ApJ, 347, 289
- Greenhouse, M., et al. 1993, ApJS, 88, 23
- Grevesse, N., & Anders, E. 1989, *Cosmic Abundances of Matter*, AIP Conference Proceedings 183, p. 1, Ed. C. J. Waddington, (New York: AIP)
- Grevesse, N. & Noels, A. 1993 in *Origin & Evolution of the Elements*, ed. N. Prantzos, E. Vangioni-Flam, & M. Casse p. 15 (Cambridge: Cambridge Univ. Press)
- Guhathakurta, P., & Draine, B. T. 1989, ApJ, 345, 230
- Guilbert, P. W. 1986, MNRAS, 218, 171
- Guilbert, P., & Rees, M. J. 1988, MNRAS, 233, 475
- Habing, H. J. 1968, Bull. Astr. Inst. Netherlands 19, 421
- Halpern, J. P., & Grindlay, J. E. 1980, ApJ, 242, 1041
- Hamann, F., & Ferland, G. J. 1992, ApJ, 391, L53
- Hamann, F., & Ferland, G. J. 1993, ApJ, 418, 11
- Hamann, F., & Ferland, G. J. 1999, ARAA, 37, 487
- Harrington, J. P. 1969, ApJ, 156, 903
- Harrington, J. P. 1973, MNRAS, 162, 43
- Heitler, W. 1954, The Quantum Theory of Radiation (Oxford: Oxford University Press)
- Hjellming, R. M. 1966, ApJ, 143, 420
- Hollenbach, D., & McKee, C. F. 1979, ApJS, 41, 555
- Hollenbach, D., & McKee, C. F. 1989, ApJ, 342, 306
- Hollenbach, D.J., Takahashi, T., & Tielens, A.G.G.M., 1991, ApJ, 377, 192-209
- Hubbard, E. N., & Puetter, R. C. 1985, ApJ, 290, 394
- Hummer, D. G. 1962, MNRAS, 125, 21
- Hummer, D. G. 1968, MNRAS, 138, 73
- Hummer, D. G. 1988, ApJ, 327, 477
- Hummer, D. G, Berrington, K. A., Eissner, W., Pradhan, A. K., Saraph H. E., Tully, J. A. 1993, A&A, 279, 298
- Hummer, D. G., & Kunasz, 1980, ApJ, 236, 609
- Hummer, D. G., & Seaton, M. J. 1963, MNRAS, 125, 437
- Hummer, D. G., & Storey, P. J. 1987, MNRAS, 224, 801
- Hummer, D. G., & Storey, P. J. 1992, MNRAS, 254, 277
- Hutchings, J.B. 1976, ApJ, 205, 103
- Ikeuchi, S., & Ostriker, J. P. 1986, ApJ, 301, 522
- Jackson, J. D. 1975, Classical Electrodynamics (New York: Wiley)
- Kaler, J., & Jacoby, G. 1991, ApJ, 372, 215
- Janev, R. K., Langer, W. D., Post, D. E., & Evans, K. 1987, Elementary Processes in Hydrogen–Helium Plasmas (Berlin: Springer--Verlag)
- Jenkins, E. B. 1987, in *Interstellar Processes*, D. Hollenbach & H. Thronson, Eds, (Dordrecht: Reidel), p.533
- Johnson, L. C. 1972, ApJ, 174, 227
- Johnstone, R. M., Fabian, A. C., Edge, A. C., & Thomas, P. A. 1992, MNRAS, 255, 431
- Kaler, J., 1978, ApJ, 220, 887
- Kallman, T. R., & McCray, R. 1982, ApJS, 50, 263
- Karzas, W. J., & Latter, R. 1961, ApJS, 6, 167
- Kaastra, J. S., & Mewe, R. 1993, A&AS, 97, 443
- Kato, T. 1976, ApJS, 30, 397
- Kellerman, K. I. 1966, ApJ, 146, 621
- Khromov, G. S. 1989, Space Science Reviews 51, 339
- Kingdon, J. B., & Ferland, G. J. 1991, PASP, 103, 752
- Kingdon, J. B., & Ferland, G. J. 1993, ApJ, 403, 211
- Kingdon, J. B., & Ferland, G. J. 1995, ApJ, 450, 691

- Kingdon, J. B., & Ferland, G. J. 1996, ApJS, 106, 205
- Kingdon, J. B., Ferland, G. J., & Feibelman, W.A. 1995, ApJ, 439, 793
- Kingdon J.B., & Ferland, G.J., 1998, ApJ 506, 323-328
- Kingdon, J. B., & Ferland, G. J. 1998, ApJ, 516, L107-109
- Korista, K. T., Baldwin, J. A., & Ferland, G. J. 1998, ApJ, 507, 24
- Korista, K. T., & Ferland, G. J. 1989, ApJ, 343, 678
- Korista, K. T., & Ferland, G. J. 1998, ApJ, 495, 672
- Korista, K. T., Ferland, G. J., & Baldwin, J. 1997, ApJ, 487, 555
- Krolik, J., McKee, C. M., & Tarter, C.B. 1981, ApJ, 249, 422
- Kurucz, R. L. 1970, SAO Special Reports 309
- Kurucz, R. L. 1979, ApJS, 40, 1
- Kurucz, R. L. 1991, in Proceedings of the Workshop on Precision Photometry: Astrophysics of the Galaxy, ed.
 A. C. Davis Philip, A. R. Upgren, & K. A. James, (Schenectady: Davis), 27
- Kwan, J., & Krolik, J. 1981, ApJ, 250, 478
- Lambert, D. L., & Pagel, B. E. J. 1968, MNRAS, 141, 299
- La Franca, Franceshini, A., Cristiani, S., & Vio, R. 1995, A&A, 299, 19
- Lame N. J., & Ferland, G. J. 1991, ApJ, 367, 208
- Landini, M., & Monsignori Fossi, B. 1990, A&AS, 82, 229
- Landini, M., & Monsignori Fossi, B. 1991, A&AS, 91, 183
- Lanzafame, A., Tully, J. A., Berrington, K. A., Dufton, P. L., Byrne, P. B., & Burgess, A. 1993, MNRAS, 264, 402
- Laor, A., & Draine, B. T. 1993, ApJ, 402, 441
- Latter, W. B., & Black, J. H. 1991, ApJ, 372, 161
- Lea, S., & Holman, G. 1978, ApJ, 222, 29
- Lennon, D. J., & Burke, V. M. 1991, MNRAS, 251, 628
- Lenzuni, P., Chernoff, D. F., & Salpeter, E. E. 1991, ApJS, 76, 759
- Levich, E. V., & Sunyaev, R.A. 1970, Astrophysical Letters 7, 69
- Lepp, S., & Shull, J. M. 1983, ApJ, 270, 578
- Lightman, A. P., & White, T.R. 1988, ApJ, 335, 57
- Lites, B. W., & Mihalas, D. 1984, Solar Physics 93, 23
- Liu, X.-W., Storey, P. J., Barlow, M. J., & Clegg, R. E. S. 1995, MNRAS, 272, 369
- Longair, M. S. 1981, *High Energy Astrophysics*, (Cambridge: Cambridge University Press)
- Lotz, W. 1967, ApJS, 14, 207
- MacAlpine, G. M. 1971, ApJ, 175, 11
- Maguire, S. 1993, Writing Solid Code, Microsoft Press
- Mallik, D. C. V., & Peimbert, M. 1988, Rev Mexicana 16, 111
- Martin, P. G. 1979, *Cosmic Dust* (Oxford: Clarendon Press)
- Martin, P. G. 1988, ApJS, 66, 125
- Martin, P. G., & Ferland, G. J. 1980, ApJ, 235, L125
- Martin, P.G., & Rouleau, F., 1991, in Malina R.F., Bowyer S., eds, Extreme Ultraviolet Astronomy, Pergamon Press, Oxford, p. 341
- Martin, P. G., & Whittet, D. C. B. 1990, ApJ, 357, 113
- Masters, A. R., Pringle, J. E., Fabian, A. C., & Rees, M. J. 1977, MNRAS, 178, 501
- Mathews, W. G., Blumenthal, G. R., & Grandi, S. A. 1980, ApJ, 235, 971
- Mathews, W. G., & Ferland, G. J. 1987, ApJ, 323, 456
- Mathis, J. S. 1982, ApJ, 261, 195
- Mathis, J. S. 1985, ApJ, 291, 247
- Mathis, J. S., Rumpl, W., & Nordsieck, K. H. 1977, ApJ, 217, 425
- Mathis, J. S., & Wallenhorst, S. G. 1981, ApJ, 244, 483
- Matteucci, F., & Tornambe, A. 1987, A&A, 185, 51
- Matteucci, F., & Greggio, A. 1986, A&A, 154, 279
- Mazzotta, P., Mazzitelli, G., Colafrancesco, C., & Vittorio, 1998, A&AS 133, 403-409
- McKee, C. F. 1999, preprint, Astro-ph 9901370
- Mendoza, C. 1983, in *Planetary Nebulae*, IAU Sym 103, D. R. Flower, Ed., p 143, (Dordrecht: Reidel) Mewe, R. 1972, A&A, 20, 215

- Mihalas, D. 1972, Non-LTE Model Atmospheres for B & O Stars, NCAR-TN/STR-76
- Mihalas, D. 1978, Stellar Atmospheres, 2nd Edition (San Francisco: W.H. Freeman)
- Mihalszki, J. S., & Ferland, G. J. 1983, PASP, 95, 284
- Morrison, R., & McCammon, D. 1983, ApJ, 270, 119
- Morton, D. C., York, D. G., & Jenkins, E. B. 1988, ApJS, 68, 449
- Nahar, S. N., & Pradhan, A. K. 1992, ApJ, 397, 729
- Netzer, H. 1990, in *Active Galactic Nuclei, Saas-Fee Advanced Course* 20, Courvorsier, T.J.-L., & Mayor, M., (Springer-Verlag; Berlin)
- Netzer, H., Elitzur, M., & Ferland, G. J. 1985, ApJ, 299, 752
- Netzer, H., & Ferland, G. J. 1984, PASP, 96, 593
- Neufeld, D. A. 1989, Harvard Research Exam
- Neufeld, D. A., & Dalgarno, A. 1989, Phys Rev A, 35, 3142
- Novotny, Eva, 1973, Introduction to Stellar Atmospheres, (New York; Oxford University Press)
- Nussbaumer, H., & Storey, P. J. 1983, A&A, 126, 75
- Nussbaumer, H., & Storey, P. J. 1984, A&AS, 56, 293
- Nussbaumer, H., & Storey, P. J. 1986, A&AS, 64, 545
- Nussbaumer, H., & Storey, P. J. 1987, A&AS, 69, 123
- Oliveira, S., & Maciel, W. J. 1986, Ap&SS, 126, 211
- Oliva, E., Pasquali, A., & Reconditi, M. 1996, A&A, 305, 210
- Osterbrock, D. E. 1951, ApJ, 114, 469
- Osterbrock, D. E. 1989, Astrophysics of Gaseous Nebulae & Active Galactic Nuclei, (Mill Valley; University Science Press)
- Osterbrock, D. E., & Flather, E. 1959, ApJ, 129, 26
- Osterbrock, D. E., Tran, H. D., & Veilleux, S. 1992, ApJ, 389, 305
- Ostriker, J. P., & Ikeuchi, S. 1983, ApJ, 268, L63
- Pacholczyk, A. G. 1970, Radio Astrophysics (San Francisco: Freeman)
- Pagel, B. E. J. 1997, Nucleosynthesis and Chemical Evolution of Galaxies, (Cambridge: Cambridge University Press)
- Palla, F., Salpeter, E. E., & Stahler, S. W. 1983, ApJ, 271, 632
- Peebles, P. J. E. 1971, Physical Cosmology, (Princeton: Princeton U. Press)
- Peimbert, M. 1967, ApJ, 150, 825
- Pengelly, R. M. 1964, MNRAS, 127, 145
- Pequignot, D. 1986, Wordshop on Model Nebulae, (Paris: l'Observatoire de Paris) p363
- Pequignot, D., & Aldrovandi, S.M.V. 1986, A&A, 161, 169
- Pequignot, D., Stasinska, G., & Aldrovandi, S. M. V. 1978, A&A, 63, 313
- Pequignot, D., Petitjean, P., & Boisson, C. 1991, A&A, 251, 680
- Peterson, J. R., Aberth, W., Moseley, J., & Sheridan, J. 1971, Phys Rev A, 3, 1651
- Press W. H., Teukolsky, S.A., Vetterling, W. T., & Flannery, B. P. 1992, *Numerical Recipes*, (Cambridge; Cambridge University Press)
- Puetter, R. C. 1981, ApJ, 251, 446
- Puy, D., Alecian, G., Le Bourlot, J., Leorat, J., & Pineau des Forets, G. 1993, A&A, 267, 337
- Rauch, T. 1997 A&A, 320, 237
- Raymond, J. C., Cox, D. P., & Smith, B. W. 1976, ApJ, 204, 290
- Rees, M. J., Netzer, H., & Ferland, G. J. 1989, ApJ, 347, 640
- van Regemorter, H. 1962, ApJ, 136, 906
- Rephaeli, Y. 1987, MNRAS, 225, 851

Reilman, R. F., & Manson, S. T. 1979, ApJS, 40, 815, errata 46, 115; 62, 939

- Roberge, W. G., Jones, D., Lepp, S., & Dalgarno, A. 1991, ApJS, 77, 287
- Rossi, B. 1952, High-Energy Particles (New York; Prentice-Hall)
- Rouleau, F., & Martin, P.G. 1991, ApJ, 377, 526
- Rowan, T. 1990, Functional Stability Analysis of Numerical Algorithms, Ph.D. Thesis, Department of Computer Sciences, University of Texas at Austin
- Rubin, R. H. 1968, ApJ, 153, 671
- Rubin, R. H. 1983, ApJ, 274, 671

- Rubin, R. H. Martin, P. G. Dufour, R. J. Ferland, G. J Baldwin, J. A. Hester, J. J. & Walter, D. K. 1998, ApJ, 495, 891
- Rubin, R. H., Simpson, J. R., Haas, M. R., & Erickson, E. F. 1991, ApJ, 374, 564
- Rybicki, G. B., & Hummer, D. G. 1991, A&A, 245, 171
- Rybicki, G. B., & Hummer, D. G. 1992, A&A, 262, 209
- Rybicki, G. B., & Hummer, D. G. 1994, A&A, 290, 553
- Rybicki, G. B., & Lightman, A.P. 1979, Radiative Processes in Astrophysics (New York: Wiley)
- Sanders, D. B., et al. 1989, ApJ, 347, 29
- Saraph, H. E. 1970, J.Phys.B., 3, 952
- Savage, B. D., & Sembach, K. R. 1996, ARA&A, 34, 279
- Savin, Daniel Wolf, 2000, ApJ, 533, 106
- Savin, D. W.; Kahn, S. M.; Linkemann, J.; Saghiri, A. A.; Schmitt, M.; Grieser, M.; Repnow, R.; Schwalm, D.; Wolf, A.; Bartsch, T.; Brandau, C.; Hoffknecht, A.; Müller, A.; Schippers, S.; Chen, M. H.; Badnell, N. R., 1999, ApJS, 123, 687
- Sciortino, S., et al. 1990, ApJ, 361, 621
- Scott, J. S., Holman, G. D., Ionson, J. A., & Papadopoulos, K. 1980, ApJ, 239, 769
- Schaerer D., de Koter, A., Schmutz, W., & Maeder, A. 1996ab, A&A, 310, 837, & A&A, 312, 475
- Schaerer D., & de Koter A. 1997, A&A, 322, 592
- Schuster, A. 1905, ApJ, 21, 1
- Schutte, W. A., Tielens, A. G. G. M., & Allamandola, L. J. 1993, ApJ, 415, 397
- Schwarzschild, M. 1965, Structure & Evolution of the Stars, (New York: Dover)
- Seaton, M. J. 1959, MNRAS, 119, 81
- Seaton, M. J. 1959, MNRAS, 119, 90
- Seaton, M. J. 1987, J.Phys. B, 20, 6363
- Sellmaier, F. H., Yamamoto, T., Pauldrach, A. W. A., & Rubin, R. H. 1996, A&A, 305, L37
- Shields, G. A. 1976, ApJ, 204, 330
- Shine, R. A., & Linsky, J. L. 1974, Solar Physics 39, 49
- Shull, J. M. 1979, ApJ, 234, 761
- Shull, J. M., & Van Steenberg, M. E. 1982, ApJS, 48, 95
- Shull, J. M., & Van Steenberg, M. E. 1985, ApJ, 298, 268
- Sellgren, K., Tokunaga, A. T., & Nakada, Y. 1990, ApJ, 349, 120
- Sellmaier, F.H., Yamamoto, T., Pauldrach, A.W.A., Rubin, R.H 1996, A&A, 305, 37
- Sikora, M., Begelman, M. C., & Rudak, B. 1989, ApJ, 341, L33
- Simonyi, C. 1977, Meta-Programming: A Software Production Method, Thesis, Stanford University
- Snow, T. P., & Dodger, S. L. 1980, ApJ, 237, 708
- Snow, T. P., & York, D. G. 1981, ApJ, 247, L39
- Snow, T.P., & Witt, A. 1996, ApJ, 468, L65
- Spitzer, L. 1948, ApJ, 107, 6
- Spitzer, L. 1962, Physics of Fully Ionized Gasses, (New York: Interscience)
- Spitzer, L. 1978, Physical Processes in the Interstellar Medium, (New York: Wiley)
- Spitzer, L. 1985, ApJ, 290, L21
- Spitzer, L., & Tomasko, M. G. 1968, ApJ, 152, 971
- Stecher, T. P., & Williams, D. A. 1967, ApJ, 149, 29
- Stoy, R. H. 1933, MNRAS, 93, 588
- Storey P. J. 1981, MNRAS, 195, 27p
- Storey, P. J. 1994, A&A, 282, 999
- Storey, P. J., & Hummer, D. G. 1991, Comput. Phys. Commun. 66, 129
- Storey, P. J., & Hummer, D. G. 1995, MNRAS, 272, 41 (on the web at <u>http://adc.gsfc.nasa.gov/adc-cgi/cat.pl?/catalogs/6/6064/</u>)
- Swings, P., & Struve, O. 1940, ApJ, 91, 546
- Tarter, C. B., & McKee, C. F. 1973, ApJ, 186, L63
- Tarter, C.B., Tucker, W.H., & Salpeter, E.E., 1969, ApJ, 156, 943
- Tielens, A. G. G. M., & Hollenbach, D. 1985a, ApJ, 291, 722
- Tielens, A. G. G. M., & Hollenbach, D. 1985b, ApJ, 291, 746
- Tout, C. A., Pols, O. R., Eggleton, P. P. & Han, Z. 1996, MNRAS, 281, 257

- Turner, J., & Pounds, K. 1989, MNRAS, 240, 833
- Van Blerkom, D., & Hummer, D. G. 1967, MNRAS, 137, 353
- van Dishoeck, E.F., & Black, J.H., 1988, ApJ, 334, 771
- van Hoof, P. A. M. 1997, PhD Thesis, University of Groningen
- van Hoof, P.A.M., Beintema, D.A., Verner D.A., & Ferland, G.J., 2000a, A&A 354, L41-L44
- van Hoof, P.A.M., Van de Steene, G.C., Beintema, D.A., Martin, P.G., Pottasch, S.R., Ferland, G. J., 2000b, ApJ 532, 384-399
- van Regemorter, H. 1962, ApJ, 136, 906
- Vedel, H., Hellsten, U., & Sommer-Larsen, J. 1994, MNRAS, 271, 743
- Vernazza, J. E., Avrett, E. H., & Loeser, C. B. 1981, ApJS, 45, 635
- Verner, D. A., Yakovlev, D. G., Band, I. M., & Trzhaskovshaya, M. B. 1993, ADNDT, 55, 233
- Verner, D. A., & Yakovlev, 1995, A&AS, 109, 125
- Verner, D. A., & Ferland, G. J. 1996, ApJS, 103, 467
- Verner, D. A., Ferland, G. J., Korista, K., & Yakovlev D. G. 1996, ApJ, 465, 487
- Verner, D. A., Verner, K., & Ferland, G. J. 1996, ADNDT, 64, 1
- Verner, E.M. Verner, D.A. Korista, K.T. Ferguson , J.W. Hamann, F. & Ferland, G.J. 1999, ApJS 120, 101
- Voronov, G. S. 1997, ADNDT, 65, 1
- Voit, G. M. 1991, ApJ, 377, 1158
- Volk, K., and Kwok, S. 1988, ApJ, 331, 435
- Vriens, L., & Smeets, A. H. M. 1980, Phys Rev A, 22, 940
- Watson, W. D. 1972, ApJ, 176, 103
- Weisheit, J. C. 1974, ApJ, 190, 735
- Weisheit, J. C., & Collins, L. A. 1976, ApJ, 210, 299
- Weisheit, J. C., & Dalgarno, A. 1972, Astrophysical Letters, 12, 103
- Weisheit, J., Shields, G. A., & Tarter, C. B. 1981, ApJ, 245, 406
- Werner, K., & Heber, U. 1991, in *Stellar Atmospheres: Beyond Classical Models*, p 341, NATO ASI Series C, eds. L. Crivellari, I. Hubney, & D. G. Hummer, (Dordrect: Kluwer)
- White, R. E. 1986, ApJ, 307, 777
- Wiese, W.L., Fuhr, J.R., & Deters, T.M., 1996, J Phys Chem Ref Data, Monograph 7
- Wiese, W. L., Smith, M. W., & Glennon, B. M. 1966, NSRDS-NBS 4
- Wilkes, B. J., Ferland, G. J., Truran, J., & Hanes, D. 1981, MNRAS, 197, 1
- Wilkes, et al 1994, ApJS, 92, 53
- Wilkinson, D. T. 1987, in 13th Texas Symposium on Relativistic Astrophysics, M. P. Ulmer, ed., (Singapore: World Scientific), p209
- Williams, R. E. 1967, ApJ, 147, 556
- Williams, R. E. 1992, ApJ, 392, 99
- Wills, B., Netzer, H., & Wills, D. 1985, ApJ, 288, 94
- Winslow, A. M. 1975, Lawrence Livermore Lab. report UCID-16854
- Wishart, A. W. 1979, MNRAS, 187, 59p
- Wolfire, M. G., Tielens, A., & Hollenbach, D. 1990, ApJ, 358, 116
- Worral et al. 1987, ApJ, 313, 596
- Wyse, A. B. 1941, PASP, 53, 184
- York, D. G., Meneguzzi, M., & Snow, T. 1982, ApJ, 255, 524
- Xu, Y., & McCray, R. 1991, ApJ, 375, 190
- Zamorani, G., et al. 1981, ApJ, 245, 357
- Zheng, W., Kriss, G.A., Telfer, R.C., Grimes, JP. & Davidsen, A.F. 1997, ApJ, 475, 469
- Zuckerman, B. 1973, ApJ, 183, 863
- Zycki, P. T., Krolik, J. H., Zdziarski, A. A., & Kallman, T. R. 1994, 437, 597
- Zygelman, B., & Dalgarno, A. 1990, ApJ, 365, 239

- A absolute magnitude, 26 absolute magnitude command, 26 abundances absolute defined, 50 changing input order, 55 command, 51 default, 50 depletion, 57 fluctuations, 56 ionic, 262 overview, 50 printed, 601 reading from table, 56 scale defined, 50 set, 51 starburst, 54 variables used, 423 abundances command, 51 abundances starburst command, 54 acceleration. see radiative acceleration adding lines, 229 adding models together, 411, 412 adiabatic cooling. see expansion cooling age checking, 65 printing, 98 age command, 11, 65 agn command, 32 albedo punch cloud, 107 gas, 112 Alfvén velocity, 86 alpha code, 402 alpha ox printed, 602 setting, 31 aluminum, 267 apparent helium abundance, 616 arbitrary density law, 60 assign statement in VMS, 43 Atlas 1991 atmospheres, 45 atlas.in input script, 457 atomic data references, 418 atomic mass, 214 atomic weight, 214, 262 Auger effect, 258, 263 averaging over terms, 224 -B-

background cosmic, 34, 36, 37 cosmic ray, 84, 298 background command, 34 bangot, 410 Be-sequence cooling, 284

beta code, 402 Bethe-Block approximation, 298 blackbody definition, 206 evaluated, 206 blackbody command, 35 vary, 124 blister.in input script, 458 predictions, 444 blr.in input script, 459 Bohr radius, 222, 308 Boltzmann factor continuum, 204 line, 213 Born approximation, 222 bound-free opacity, 212 Bowen OI, 266 brems.in input script, 459 bremsstrahlung continuum, 36 cooling, 277 heating, 277 opacity, 212 bremsstrahlung command, 36 brightness temperature, 604 broken code, 419

-C-

calcium, 267 Cameron abundances, 52 carbon, 263 carbon fully molecular, stopped because of, 609 casea.in input script, 460 casebn2.in input script, 461 casebn8.in input script, 461 casec.in input script, 462 caunot, 410 cautions checks, 4 printed, 611 routine to print, 410 cdColm, 408 cdDriv, 407 cdEms, 408 cdGett, 412 cdInit, 406 cdIonf, 408 cdLine, 407 cdNoex, 407 cdNwcns, 409 cdOutp, 406 cdRead, 406 cdTalk, 407 cextra command, 82

changes to code, 418 charge transfer heavy element, 135, 260 punching rates, 114 statistical, 135 turning off, 133 chemical composition printed, 601 set, 51 Cloudy 84 vs 80, 436 90 vs 84, 434 adding lines, 229 changes to code, 418 date, 419 flowchart, 414 future improvements, 437 history, 431 known modes, 437 mailing list, 401 making a revision, 437 revision history, 431 running 1 model, 402 running a grid, 405 search phase, 423 setting up, 1 size, 431 source versions, 402 version numbers, 419 cloudy.ini file, 21, 132 co-adding many models, 411, 412 coding conventions, 416 broken code, 419 changes to code, 418 characters, 418 integers, 417 logical variables, 418 real numbers, 417 routine descriptions, 419 sanity checks, 418 strong typing, 417 test code, 419 collion.for input script, 464 collision rate, 222 detailed balance, 236 ionization, 259 collision strengths averaging, 224 defined, 221 g-bar, 222 column density, 420 effective defined, 90 printout, 616 ionized, 90 neutral, 90 total, 90 printed, 605 command comments, 20 continue option, 20 example, 21

format, 18 line length, 19 numerical input, 20 reading in, 414 temperature conventions, 21 comments input stream, 20 comparison calculations, 440 comphi.in input script, 465 compiling command, 128 compiling stellar atmospheres, 128 complo.in input script, 465 composition. see abundance Compton bound heating, 277 bound ionization, 238, 277 cooling calculated, 274 output, 606 energy exchange accuracy, 275 methods, 274 turning off, 133 heating calculated, 274 output, 606 ionization, 134 temperature, 275 output, 603 timescale output, 617 compton.in test input, 466 computer Cray, 401 Exemplar, 136 Sparcstation, 401 conserv.in input script, 466 constant density command, 59 constant gas pressure command, 59 constant pressure command, 59 constant temperature command, 82 continue command, 20 continuous heavy element opacity, 269 continuum agn, 32 arrays, 203 binning, 202, 260 changing, 129 Boltzmann factor, 204 cell width, 204 changing resolution, 203 described, 202 diffuse, 6, 204 printed, 98 punched, 105

diffuse printed, 621 emission, 240 energy bounds, 202 energy pointer, 203 fluoresence, 218 frequencies saved, 203 frequency array, 203 generation, 205 high energy limit, 136, 202 incident defined, 6 punched, 106 incident printed, 621 intensity specifying, 25 ionization edge pointers, 203 low energy limit, 202 luminosity specifying, 25 mesh defined, 202 normalization, 206 occupation number, 204, 232, 239, 603 opacity, 209 optical depth, 205 OTS, 204, 208 outward, 204 outward only, 209 outward punched, 107 PDR, 27 photon flux, 204 plasma frequency, 212 pointers, 203 printing, 98 range, 33, 202 reflected, 204 defined, 7 printed, 98 resolution, 129 shape commands, 33 specify, 2 summed, 205 surface brightness, 623 transmitted defined, 6 convergence tracing, 117 conversion factors, 207 coolcurve.for, 427 input script, 467 coolhii.in input script, 470 predictions, 441 cooling adiabatic, 277 balance, 84, 87 Be-sequence atoms, 284 bremsstrahlung, 277 collisional, 279 Compton, 116, 133, 134, 274, 277,606 cyclotron, 85 definition, 273 derivative, 285 error, 616 evaluation, 285 expansion, 277 extra, 82

free-free, 277, 278 grain, 73, 293, 294, 606 H-, 279 heavy elements, 280 history, 115 hydrodynamic, 277 hydrogen lines, 279 hydrogen recombination, 233 induced, 278 low energy continuum, 33, 34, 39 map, 84, 86, 97 n level atoms, 282 numerical derivatives, 138 plot, 97 printout, 98, 615 punch, 108 recombination, 233, 278 tests, 286 three level atoms, 281 two level atoms, 281 cooling flow continuum, 41 corners4.for input script, 469 coronal equilibrium command, 83 coronal.in input script, 468 cosmic ray, 82, 83, 86 background, 84, 298 command, 83 physics, 297 range, 297 cosmic ray command, 83, 298 cosmology distance, 307 covering factor command, 65 computational details, 424 defined, 8 geometric, 424 radiative transfer, 425 sphere option, 68 covering factor command, 65 Cray, 401 critical density, 223 cyclotron cooling, 85 cylinder command, 66 -Ddamping constant defined, 217 stored, 217, 226 density arbitrary law, 60 commands, 59 constant, 59 constant gas pressure, 59 constant total pressure, 59 cosmic ray, 83, 297 electron evaluated, 421 limiting, 91 output, 604 energy, 35 fluctuations, 61

globule law, 62

H-, 246

LTE, 245 H2 LTE, 245 hydrogen, 28, 59 LTE, 231 molecules, 245 output, 604 mass, 421 molecules, 245 particle, 421 power-law, 63, 64 range, 13 structure output, 613 structure saved, 422 table entered, 61 total particle, 303 wind law, 304 density per particle, 421 departure coefficients hydrogen, 231 printing, 98 depletion factors, 57 depth defined, 7 derivative cooling, 285 heating, 286 numerical heating, cooling, 138 destruction probability, 217 hydrogen, 232 line complete, 218 dielectronic recombination 3rd and 4th row elements, 130 Burgess, 130 Nussbaumer and Storey, 130 dielectronic recombination command, 130 diffuse fields defined, 6 evaluated, 204 punching source function, 115 test case, 474 transfer, 71 diffuse fields command, 71 diffuse OTS command, 71, 208 diffuse outward command, 71, 209 dilution factor, 36 dissociation energy H-, 245 H2, 245 H2+, 245 distance from redshift, 307 dlaw command, 60 Doppler width computed, 214 double optical depths command, 71 dqher.in input script, 470 predictions, 453 drive command, 131

drive escape probabilities, 131 drive fread, 131 drive gaunt factor command, 131 drive hyas command, 131 drive molecules command, 131 drive pointers command, 131 drive starburst command, 132 drive.for input script, 471 dumping line, 228 dust. see grain dyn, 302 -Eeden command, 132 eden.in input script, 472 Einstein coefficients, 215 electron adding extra, 132 density evaluated, 421 saved, 421 mean speed, 85 non-thermal secondaries, 305 secondary ionization, 305 supra-thermal, 305 temperature. see temperature element abundances stored, 262 chemical symbols, 262 ionization stage symbols, 262 names, 263 periodic table, 258 element table command, 56 elements command, 54 vary, 124 elements read command, 55 emission line list, 626 emission measure, 616 end of input stream, 19 end of line characters, 19 energy density gas, 302 energy density command, 27 equivalent width computing, 621 escape probability heavy element, 225 helium, 256 hydrogen, 232 line, 216 complete, 217 incomplete, 217 maser, 218 routines, 216 escape probability command, 71 excitation energy, 213 excitation temperature, 228 Exemplar, 136 expansion cooling, 277 extinguish command, 36

-Ff(nu) command, 27 fabden, 60 Fe II atom command, 72 punching intensities, 112 Fe II command, 72 filling factor, 66, 69, 304, 606, 616 defined, 8, 67 filling factor command, 66 vary, 124 fine structure lines, 225 fireball command, 37 floating point errors, 401, 429 flowcharts, 414 Cloudy, 414 ionte, 415 MainCl, 414 PIonte, 414 TauInc, 415 fluc.in input script, 472 fluctuations command, 56, 61 fluorescence continuum, 218 vields, 263 flux converting to luminosity, 623 force temperature command, 84 free-free cooling, 277 H-, 279 emission, 240 gaunt factor, 114, 212 heating, 277 H-, 279 opacity, 212 free-free cooling plasma frequency, 278 fudge factors command, 132

-G-

gas albedo punched, 107, 112 energy density, 302 equation of state, 302 heat content, 302 mass density, 421 opacity punching, 111 particle density, 421 pressure, 59, 303 gaunt factor drive command, 131 free-free, 114, 212 g-bar approximation, 222 geometry closed, 9, 68, 613 closed expanding, 221 closed static, 221 cylindrical, 7 definition, 7 details, 419 disk, 65 ionized from both sides, 71

matter-bounded, 9, 89, 91 open, 3, 8, 68, 76, 80, 221, 613 plane parallel, 7, 67, 611 plane parallel vs spherical, 34,65 printed, 611 radiation-bounded, 10, 89 specify, 2, 3 spherical, 4, 7, 68, 80, 611 thick shell, 7, 611 wind, 7, 69, 221, 617 globule command, 62 globule.in input script, 473 glossary routines, 692 symbols, 308 Go continuum, 28 gold code, 402 grain, 289 absorption efficiency factor, 290 agb, 74 area, 290 collisional charging, 292 drift velocity, 293 dust to gas ratio, 74 exist?, 295 extra species, 74 gas cooling, 294 gas heating, 294 gray, 74 heating and cooling, 293 HII Hegion, 74 ionic recombination, 134, 295 ISM, 73 opacity, 289 punching, 112 Orion, 74 PAH, 74 photoelectric emission, 290 planetary nebula, 74 potential, 292 quantum heating, 80 specifying, 72 temperature, 294 variables, 295 grains command, 72 grey1.opac, 74 grid0.for input script, 474

-H-

H II region abundances, 52 Habing radiation field, 27 Hartree, 207 Hazy printing, 438 hden command, 62 vary, 124 heating balance, 84, 87 bound Compton, 277 bremsstrahlung, 277 collisional ionization, 278 Compton, 116, 133, 134, 274, 277, 603, 606

continuum, 202 cosmic ray, 84, 297, 298 definition, 273 derivative, 286 evaluated, 285 extra, 82, 85, 607 fine structure lines, 225, 280 free-free, 134, 277, 278, 616 grain, 73, 290, 292, 293, 294, 606 H-, 279 heating array, 285 history, 115 hydrogen lines, 279 line, 286 low energy continuum, 33, 39 map, 84, 86, 97 neutron, 87 numerical derivatives, 138 photoelectric, 278, 613 plot, 97 print, 99, 116 printout, 615 punch, 108 quantum grain, 80 secondaries, 135 structure saved, 421 tests, 286 total, 605, 616 evaluated, 286 heavy element abundances stored, 262 adding lines, 229 atomic weights, 262 Auger ejection, 258, 263 charge transfer, 260 chemical symbols, 262 collisional ionization, 259 depletion, 57 FeII atom, 72 fluorescence yields, 263 ionization pointers, 260, 263 ionization potentials, 260 ionization stored, 262 labels for lines, 228 limits, 16 line pointers, 226 line transfer, 225 molecules, 253 names, 263 number of subshells, 261 opacity offsets, 260 overview, 257 photoionization cross section, 258 photoionization rates stored, 263 punching opacity, 112 punching pointers, 113 recombination dielectronic, 259 grain surface, 295 radiative, 259 reliability, 271 shell number, 263 TauLines subdirectory, 229 helium continuum pointers, 256

escape probability, 256 ion, 255 line pointer, 256 printout ionization, 606 lines, 614 punching, 108 radiative transfer, 256 recombination coefficients, 256 recombination efficiency punching, 114 singlets, 255 statistical weight, 240 triplets, 255 helium collisions command, 76 hemis.in input script, 474 hextra command, 85 high temperature approach command, 85 highn.in input script, 475 HII region abundances, 52 hiiregions.for input script, 476 hizlte.for input script, 477 hizqso.in input script, 478 Hungarian naming convention, 416 hydrodynamic cooling, 277 hydrogen 2s 2p collisions turned off, 133 balance equations, 236 Boltzmann factors, 231 collisional ionization, 236 collisional rate equations, 236 collisions turned off, 77 column density, 90 continuum pointer, 232 density, 28, 59, 62 LTE, 231 saved, 421 departure coefficient, 231 destruction probability, 232 emission, 240 escape probability, 232 H-, 246 H2, 250 HeH+, 250 induced recombination, 237 ionization energies, 231 ionization processes, 108 ionization solution, 236 level energies, 231 level populations punched, 108 levels command, 77 line intensities, 623 line pointer, 232 lowest temp command, 77 matrix inversion routines, 77 molecules, 245

number of levels, 77 oscillator strengths, 232 overview, 14 populations punched, 108 radiative rate equations, 237 recombination coefficients, 233, 238 elements defined, 236 recombination cooling, 233 recombination efficiency punching, 114 statistical weight, 231, 240 three body recombination, 236 top off, 78 transition probabilities, 232 driving, 131 turning off 2s 2p collisions, 133 turning off collisions, 77 hydrogen command, 76

negative ion, 246

-I-

illuminate command, 67 illuminated face, 7 induc.in input script, 478 induced emission probability, 215 induced recombination, 237 infinite loop. see unending loop init command, 21, 132 initialization command, 21, 132 initialization file, 21, 132 inner radius, 8 input line. see command intensity command, 25, 27 vary, 124 interpolate command, 38 interstellar radiation field, 41 ionic abundnaces, 262 ionization parameter command, 28 ionization potential density, 260 elements, 260 pointers, 260 ionte flowchart, 415 iron, 268 Fe II atom, 72 ISM abundances, 52 ism.in input script, 479 isobaric, 59 isochoric, 59 isoelectronic sequence, 263 iterate command, 78 iteration last, 423 variables, 422 _|_

Jeans length, 307, 617 Jeans mass, 307, 617

-Kkk.in input script, 480 predictions, 454 kmt.for, 427 kmt.in input script, 481 K-shell energy limit, 137 Kurucz 1991 atmospheres, 45 -Ll(nu) command, 28 labels for lines, 228 lalpha.in input script, 482 laser command, 39 laser1.in input script, 482 laser2.in input script, 483 laser3.in input script, 484 ldl.ir input script, 484 level 1 line, 225, 227 level 2 line, 225, 227 levels averaging, 224 line adding to line arrays, 229 array printing, 228 array punched, 109 asymmetries, 622 beaming, 622 Boltzmann factor, 213 closed expanding geometry, 221 closed static geometry, 221 continuum contrast, 622 continuum pumping contribution, 625 correction for stimulated emission, 213 data punched, 110 dumping, 228 energy pointer, 203 equivalent width, 621 escape probability, 216 excitation temperature, 228 fine structure heating, 225, 280 heating, 286 hydrogen escape probability, 232 infrared defined, 225 intensities with grains, 624 label generating, 228

punching, 110

open geometry, 221

optical depth arrays, 221,

optical depth, 213

level 1, 225, 227

level 2, 225, 227

list, 626

229

masing, 280

optical depth printing, 101 output, 634 outward, 204 pointer, 203 radiation pressure, 298 maximum printed, 616 printed, 605 stop, 608 redistribution function, 227 reflected, 204 source function, 216 spectroscopic designation, 228 surface brightness, 623 thermalization length, 223 trace, 116 transfer flowchart, 415 wavelength convention, 626 width, 214 wind geometry, 221 line cooling. see cooling liner.in input script, 485 linpack routines, 77 loop. see infinite loop LTE H- density, 245 H departure coefficient, 231 H level population, 231 H2 density, 245 lte.in input script, 485 ltemetl.in input script, 486 luminosity converting to flux, 623 distance, 307 sun, 308 luminosity command, 25, 29 vary, 124 -Mmachine environment, 401 magnesium, 266 magnetic field cyclotron cooling, 85 pressure, 86 magnetic field command, 85 magnitude absolute, 26 visual, 26 MainCl flowchart, 414 map number of steps, 138 output, 428 plot map command, 97 punch output, 111 map command, 86 map.in, 426 input script, 486 Martin, P.G. added lines, 97 print sort, 102 maser correction for stimulated emission, 213 escape probability, 218 mass

AMU, 308 electron, 308 Jeans', 308 proton, 308 sun, 309 mass density, 421 matchn2.in input script, 487 matrix inversion routines, 77 matter-bounded geometry, 9 mean speed electron, 85 metals command, 56 vary, 124 Milne relation, 233, 240 molecular weight, 421 molecules, 16, 243 drive command, 131 heavy element, 253 hydrogen, 245 printed, 607 Moore's Law, 431

-N-

naming convention, 416 negative line intensities, 615 negative mole abundance, stopped because of, 609 neon, 266 neutron heating, 87 neutrons command, 87 ngc5548.in input script, 487 nitrogen, 265 nlr.in input script, 487 predictions, 451 no 2p2s command, 133 no Auger effect command, 133 no charge transfer command, 133 no Compton effect command, 133 no feii pumping command, 133 no fine structure command, 133 no free free heating command, 134 no grain neutralization command, 134 no induced processes command, 134 no molecules command, 134 no on the spot command, 134 no photoionization command, 134 no radiation pressure command, 134 no recoil ionization command, 134 no scattering opacity command, 80 no secondary ionization command, 135 no Stark broadening command, 135

no three body recombination command, 135 no vary command, 119, 135 normalize command, 4, 95 noteot, 410 notes checks, 4 printed, 611 routine to print, 410 nova.in input script, 488 nuf(nu) command, 29 nul(nu) command, 30 numerical input. see command

-0-

observed quantities, 621 occupation number array, 204 continuum, 220, 239, 603 defined, 204 hydrogen, 232 pumping rate, 216 summed continuum, 205 OI fluorescense, 266 oldblr.in input script, 488 opacity, 209 absorption array, 205 arrays, 209 background line, 217 bound-free, 212 free-free, 212 heavy element continuous, 269 permited line, 215 pointers, 209 punching, 111 punching negative, 137 Rayleigh scattering, 211 scattering array, 205 stimulated emission, 213 optical depth arrays, 221 closed geometry, 221 commands, 70 continuum, 205, 212 convergence, 78 correction for stimulated emission, 213 double, 71 filling factor, 67 grain, 290 helium, 256 hydrogen, 232 incremented, 205 line, 213 line center, 213 line center vs mean, 214 mean vs line center, 214 open geometry, 221 output, 618 outward first estimate, 215 later updates, 215 test for definition, 215 printing, 101 sphere, 68

stimulated emission, 213 stopped because of, 609 stopping, 90, 92 updated, 205, 422 wind, 69, 221 optical to X-ray ratio, 31, 602 optim.in input script, 489 optimize amoeba method, 120 column density, 120 convergence criteria, 123 example, 119 increment, 120 intensity, 121 iterations, 121 lines, 121 luminosity, 121 no vary command, 119 Powell method, 122 Press et al codes, 119 punch, 122 range of variation, 122 subplex method, 123 tolerance, 123 trace flow, 123 trace starting at xx, 123 variables, 127 optimize column density command, 120 optimize increment command, 120 optimize intensity command, 121 optimize iterations command, 121 optimize lines command, 121 optimize luminosity command, 121 optimize Powell, 122 optimize punch command, 122 optimize range command, 122 optimize subplex command, 123 optimize tolerance command, 123 optimize trace flow, 123 optimize trace start command, 123 optimizing the spectrum, 118 Orion abundances, 52 orion.in input script, 489, 490 orionpdr.in input script, 491 oscillator strength, 213 absorption, 214 emission, 214 OTS fields described, 208 punching, 112 output apparent helium abundance, 616 cautions, 611 comments, 611 continuum, 620 emission lines, 612

header, 601 line, 634 map, 428 notes, 611 Peimbert, 617 redirection, 406 surprises, 611 warnings, 611 wavelength convention, 626 zone, 604 outward-only approximation, 209 oxygen, 265 -P-PAH grains, 74 Paris H II region, 442, 492

NLR, 493 planetary nebula, 493 parishii.in input script, 492 predictions, 442 parisnlr.in input script, 493 predictions, 449 parispn.in input script, 493 predictions, 446 parsec, 309 particle density, 303, 421 path setting, 138 PDR continuum, 27 pdr.in input script, 494 Peimbert temperature fluctuations, 617 periodic table, 258 phfit command, 138 phi(h) command, 30 vary, 124 photoerosion, 617 photoionization cross sections, 258 subshell, 112 version, 138 fluorescence yields, 263 rates evaluated, 210 rates punched, 109 rates stored, 263 turning off, 134 physical conditions punching, 114 PIonte flowchart, 414 Planck function. see blackbody computed, 115 plane parallel geometry, 7 planetary nebula abundances, 52 plasma frequency, 212 free-free cooling, 278 plot continuum, 96 map, 97 opacity, 96 printed, 612

publication quality, 95 range options, 96 plot command, 95 plot continuum command, 96 plot map command, 97 plot opacity command, 96 pnots.in input script, 494 pointers generating, 203 H continua, 232 H lines, 232 He continua, 256 He lines, 256 heavy element continua, 260 heavy element lines, 227 power law command, 39 vary, 124 power law continuum, 42 pressure compared, 86 constant, 59, 134 gas, 59 total, 59 convergence, 303 dynamic, 303 gas, 4, 59, 303, 606, 607 getting from code, 409 instability, 60, 70, 608 integrated continuum, 303 magnetic, 60, 86 printed, 607 punching, 114 radiation, 4, 68, 70, 78, 134, 303, 605, 606, 607, 616 ram, 303 total, 59, 302, 607 printed, 607 turbulent, 60, 81, 303 units, 302 variables and routines, 303 primal.in input script, 495 print ages command, 98 print all command, 97 print arrays command, 98 print continuum command, print coolants command, 98 print departure coefficients command, 98 print errors command, 99 print every command, 99 print faint command, 99 print heating command, 99 print last command, 99 print line all command, 100 print line collisions command, 100 print line heat command, 100 print line inward command, 100 print line optical depths command, 100 print line pump command, 100 print line sum command, 101 print off command, 101 print on command, 101

print only command, 101 print optical depths command, 101 print quiet command, 102 print short command, 102 print sort command, 102 print starting at command, 102 proper distance, 307 punch _dr_, 114 _raw continuum, 107 abundances, 104 charge transfer, 114 continuum, 104 continuum bins, 105 convergence, 107 cooling, 108 diffuse continuum, 105 emitted continuum, 106 file name, 103 gammas, 109 heating, 108 helium, 108 hydrogen conditions, 108 hydrogen ionization, 108 hydrogen populations, 108 incident continuum, 106 initial unit, 137 interactive continuum, 106 ionizaton structure, 108 ionizing continuum, 106 ip, 109 line contrast, 622 lines array, 109 contrast, 104, 138 cumulative, 109 data, 110 intensity, 110 labels, 110 structure, 110 map, 111 opacity, 111 OTS, 112 outward continuum, 107 overview, 113 PDR, 113 physical conditions, 114 pointers, 113 pressure, 114 recombination coefficients, 114 reflected continuum, 107 reserved, 103 results command, 113 using, 412 source function, 115 special, 115 tegrid, 115 TPredictor command, 115 transmitted continuum, 107 Verner, 112 punch command, 102 punch output, 102

Q(H) command, 30 vary, 124 gheat command, 80 quiet mode setting, 102, 407 -Rradiation pressure, 303 line, 298 maximum printed, 616 printed, 605 stop, 608 radiation-bounded geometry, 10 radiative acceleration computed, 301 printed, 607 punched, 114 wind, 605 radius defined, 7 inner, 8 saved, 422 sun, 309 radius command, 67 vary, 125 range option, 25 ratio command, 31 rauch.in input script, 495 Rayleigh scattering, 13, 68, 77, 211, 302, 619 reading commands with fillar, 414 reageo, 410 recombination arrays, 236 coefficients, 236 helium, 256 punching, 114 cooling, 233 efficiency, 236 punching, 114 grain surface, 295 hydrogenic, 233, 236 induced, 237 modifying dielectronic, 130 radiative rates, 259 three body, 135 redirecting output, 406 redshift distance, 307 reflector.in input script, 495 plotted, 7, 622 reliability, 16 revision history, 431 rnfa.in input script, 496 predictions, 455 rnfb.in input script, 497 predictions, 456 routine abscf, 213 AddOpac, 209 bangot, 410 beseq, 284

-0-

bnfun, 210 boltgn, 204, 414 broken, 419 caunot, 410 cdColm, 408 cdDriv, 407, 414 cdEms, 408 cdErrors, 410 cdGetPres, 409 cdGett, 113, 412 cdGetTe, 409 cdInit, 406, 414 cdIonf, 408 cdLine, 407 cdNoex, 407 cdNwcns, 409 cdOutp, 406 cdRead, 406 cdTalk, 407 cfit, 259 chIonLbl, 228 chLineLbl, 228 Cloudy, 414 coladd, 281 ColStrGBar, 223 conorm, 206 conpmp, 131 ConvIoniz, 415 coolr, 82, 277, 285 csphot, 209 descriptions, 419 dgeco, 77 dgesl, 77 diffem, 204 dmpary, 98 DoPunch, 102 drvary, 414 DumpLine, 228 eina, 215 EinstA, 131, 232 eovrlp, 217 esccom, 216 escinc, 216 escla, 216 escmase, 216, 218 esum, 421 fabden, 60 ffun, 205 ffun1, 205 fill, 129, 202 fillar, 414 flcsub, 62 fosc, 232 freeht, 277 fudge, 132 gamfun, 210 gamk, 210 GetPunch, 102 gffsub, 212 GrnVryDpth, 75 hcolst, 231 HCTIon, 260 HCTRecom, 260 HeTran, 256 highen, 274, 277 hjbar, 232 hmole, 245, 295 hrcf, 235

BiDiag, 272

htrans, 232, 238 HydroCool, 277 HydroPesc, 232 ionte, 414, 415 ipConSafe, 203 ipLinSafe, 203 ipoint, 203 ipShells, 263 level2, 281 level3, 281 LevelN, 282 LimitSh, 261 linpack, 77 MainCl, 414 MakeCharTran, 260 MakeRT, 225 MakeStatRT, 225 MakeWindRT, 225 matin1,77 metdif, 204 NextDR, 62, 420 nockon, 305 noteot, 410 oilevl. 266 opac0, 209 p8446, 266 phfit, 258 Plonte, 303, 414 Plankf, 115, 206 presur, 303, 304, 414 PrintElem, 424 PrtGamma, 210 PunCool, 108 PunHeat, 108 PutCS, 230 PutLine, 230 radinc, 205, 303 rdfile, 74 reageo, 410 rrfit, 259 SetCon, 136 SetPoint, 202, 263, 274 SumContinuum, 205 SumCool, 285, 286 SumHeat, 286 tauff, 278 TauInc, 415, 420 tauout, 215 TestCode, 419 TexcLine, 228 tfidle, 421, 422 TotalPressure, 303, 307, 421 update, 79, 205, 215, 422, 429 veclib, 77 velset, 214 warnot, 410 wgadd, 411 wginit, 411 wgline, 411 zonsrt, 419, 422 routine glossary, 692 routine names, 685 routine naming convention, 685 running Cloudy, 402 Rydberg unit, 206

-S-Saha equation arbitrary species, 243 ions, 244 molecular hydrogen, 245 secndary.in input script, 497 secondary ionization physics, 305 routines, 305 setting rate, 135 test case, 497 turning off, 135 variables, 305 set command, 135 charge transfer, 135 colimt, 135 csupra, 135 didz, 136 dr, 136 drmax, 136 drmin, 136 DstWght, 136 EdenError, 136 Exemplar, 136 flxfnt, 136 iPunDef, 137 kshell, 137 negopc, 137 nend, 137 nmaps, 138 numerical derivatives, 138 path, 138 phfit, 138 PunchLWidth, 138 test, 139 trace, 139 trim, 139 tsqden, 139 WeakHeatCool, 140 setting the path, 138 shell number, 263 shielded face, 7 size of code, 431 Sobolev approximation, 304 solar luminosity, 308 solar mass, 309 solar radius, 309 sound travel time, 616 source function punching continuum, 115 Sparcstation, 401 spectral index alpha ox, 31 incident continuum, 31, 602 transmitted continuum, 617 speed electron, 85 sphere approximations used, 68 static vs expanding, 9 sphere command, 3, 68 spherical geometry, 7 sphericity ratio, 420 sqrden.in input script, 498 stability thermal, 273, 604 starburst abundances, 54, 132

Stark broadening, 219 statistical weight electron, 231 H-, 245 H2, 245 H2+, 245 helium, 240 Ho, 245 hydrogen, 240 line, 227 nuclear ignored, 231 stellar atmospheres Atlas91, 45 compiling, 128 Kurucz, 44 Mihalas, 44 Werner, 48 stimulated emission optical depths corrected for, 213 stop cabon fully molecular, 609 code returned busted, 609 column density reached, 610 criteria discussed, 89 dr small rel to thick, 609 drad small, 609 highest Te reached, 610 internal error, 610 line ratio reached, 610 low electron fraction, 608 lowest EDEN reached, 608 lowest Te reached, 610 negative mole abundan, 609 nzone reached, 610 optical depth reached, 609 outer radius reached, 610 radiation pressure, 608 reason, 608 temperature out of bounds, 610 wind velocity < 0,608stop column density command, 90 vary, 125 stop eden command, 91 stop efrac command, 91 stop line command, 91 stop optical depth command, 92 stop temperature command, 93 stop thickness command, 93 vary, 125 stop zone command, 94 Stoy ratio, 101 strom.in input script, 498 strong typing, 417 structure variables, 421 style conventions for coding, 416 subroutine use as a, 404 sun luminosity, 308 mass, 309 radius, 309

supra-thermal ionization. See secondary ionization surface brightness computing, 623 surprises checks, 4 printed, 611 routine to print, 410 -Tt² analysis, 139 table agn, 40 akn 120, 41 Atlas, 45 cooling flow, 41 crab, 41 density law, 61 ISM radiation field, 41 Kurucz, 44 Mihalas, 44 power law, 42 read, 43 Rubin, 44 star Atlas, 45 Kurucz, 44 Mihalas, 44 Werner, 48 table command, 40 table stars command vary, 125 TauInc flowchart, 415 temperature blackbody, 35 brightness, 604 Compton, 275 output, 603 constant electron, 82 coronal equilibrium, 83 cosmic ray, 84 electron output, 604 energy density, 35 output, 603, 606 excitation, 228 failure, 84, 427 fireball, 34, 37 fluctuations, 617 force, 84 getting from code, 409 grain, 73, 294 output, 606 high approach, 85 history, 115 hydrogen low limit, 77 input format, 21 jumps, 428 last, 422 line excitation, 228 LTE limit, 286 Lyα excitation, 606 map, 86 map plot, 97 mean grain output, 618 multi-phase, 428 predictor punched, 115

proposed, 422 range, 13 saved, 422 stability, 428 stellar atmosphere, 45 stop command, 93 t2, 617 tolerance, 87 unstable, 604 variables, 422 terms averaging, 224 test code, 419 tests atlas.in, 457 blister.in, 458 blr.in, 459 brems.in, 459 casea.in, 460 casebn2.in, 461 casebn8.in, 461 casec.in, 462 collion.for, 464 comphi.in, 465 complo.in, 465 compton.in, 466 conserv.in, 466 coolcurve.for, 427, 467 coolhii.in, 470 corners4.for, 469 coronal.in, 468 dqher.in, 470 drive.for, 471 eden.in, 472 fluc.in, 472 globule.in, 473 grid0.for, 474 hemis.in, 474 highn.in, 475 hiiregions.in, 476 hizlte.for, 477 hizqso.in, 478 induc.in, 478 ism.in, 479 kk.in, 480 kmt.for, 427 kmt.in, 481 lalpha.in, 482 laser1.in, 482 laser2.in, 483 laser3.in, 484 ldl.in, 484 liner.in, 485 lte.in, 485 ltemetl.in, 486 map.in, 426, 486 matchn2.in, 487 ngc5548.in, 487 nlr.in, 487 nova.in, 488 oldblr.in, 488 optim.in, 489 orion.in, 489, 490 orionpdr.in, 491 parishii.in, 492 parisnlr.in, 493 parispn.in, 493 pdr.in, 494 pnots.in, 494

primal.in, 495 rauch.in, 495 reflector.in, 495, 622 rnfa.in, 496 rnfb.in, 497 secndary.in, 497 sqrden.in, 498 strom.in, 498 varyN.for, 499 varyNU.for, 500 vbhum.in, 501 werner.in, 502 wind.in, 502 thermal maps. see map creating, 97 thermal stability, 273, 604 thermalization length, 223 thick shell geometry, 7 three body recombination, 135 three level atoms, 281 time dependent command, 11, 87 timescale age command, 65 Compton, 617 photoerosion, 617 sound travel, 616 thermal, 617 title command, 115 tolerance command, 87 trace command, 116 trace convergence command, 117 trace output, 116 transition probability, 215 averaging, 224 driving hydrogenic, 131 turbulence, 82, 214, 607 setting, 81 velocity, 81 turbulence command, 81 two level atoms, 281 -Uunending loop. see loop Unix path, 138 -V-V filter, 26 van Regemorter approximation, 222 variable abnset, 424 alogete, 422 alogte, 422 amu, 262 AngleIllum, 67 anu, 203, 275 anuSave, 203 AtomicWeight, 214, 262 autocv, 79, 429

ch2pls, 420 character, 418 chargt, 272 chContLabel, 203 chDate, 419 chDffTrns, 71 chehp, 420 chElementSym, 262 chElNames, 263 chheat, 275 chi, 420 chii, 420 chIonStage, 262 chLineLabel, 203 chmin, 420 chOptRtn, 127 chVersion, 419 cmcool, 275 colden, 420 colh2, 420 colimt, 135, 609 collid, 306 condif, 204 cooling, 285 corind, 204 covgeo, 65, 424 covrt, 65, 425 create, 272 csigc, 274 csigh, 274 csupeff, 306 csupra, 135, 305 CTHion, 260 CTHrec, 260 ctot, 285 dampln, 217 darea, 296 densty, 307, 421 depabs, 205 deplon, 424 depsct, 205 depset, 423, 424 depth, 609 destroy, 272 didz, 136 diffus, 105, 204 dlaw, 61 dmetal, 57, 424 doppler, 214 drad, 419, 609 dReff, 420, 422 drnxt, 420 dsize, 296 dstab, 295 dstab1, 295 dstdft, 295 dstfac, 296 dston1, 295 dstpot, 295 dstq, 295 dstsc, 295 dstsc1, 295 DstWght, 136 dustp, 295 dVeff, 420, 422 dwork, 296 e2tau, 205 eden, 421 EdenError, 136, 421

edensqte, 421 EdenTrue, 421 ednstr, 421 eev, 295 efionz, 305 efrend, 91 egamry, 202 emm, 202 en1rvd, 207 endedn, 91 EnergyKshell, 137 eVdegK, 207 evRyd, 207 exctef, 305 exptau, 205 facexp, 205 ffheat, 277 floating, 417 flong, 62 flux, 107, 204, 275 FluxFaint, 136 flxfnt, 202 getpar, 412 gionrc, 295 grmetl, 57 halfte, 285 hbn, 231 hbrems, 277 hbul, 232 hcbolt, 231 HCharExcIon, 260 HCharExcRec, 260 HCionT, 232 hcont, 233 HCTMin, 135 hdamp, 217 HdeltaT, 232 hden, 15 hdest, 232 he1dmp, 217 he1lim, 256 he1rec, 256 he1stat, 240 he1tau, 256 he2dmp, 217 he2lim, 256 he2rec, 256 he2stat, 240 he2tau, 256 he3n, 255 heatef, 305 heating, 285 heatnt, 285 heatstr, 421 helax, 306 hemis, 232 HEnrRyd, 232 hesc, 232 hfrcin, 232 hgamnc, 237 hiistr, 421 HIonPot, 207 histr, 421 hlax, 306 hlbolt, 231 hlte, 231 HNIonRyd, 232 hrec, 236, 238 hreff, 134

hstat, 231, 240 htau, 232 htlim, 232 htnext, 232 htot, 286 in equations, 418 integers, 417 ipElement, 260 iphe11, 256 iphe2l, 256 ipHeavy, 261 iphl, 232 ipLnAovTOT, 227, 280 ipLnAul, 227 ipLnBolt, 227 ipLnColovTOT, 227 ipLnCont, 227 ipLnCool, 227, 280 ipLnCS, 227 ipLnCS1, 223, 226, 227 ipLnCS2, 226, 227 ipLnDamp, 227 ipLnDampRel, 226, 227 ipLnDesP, 226, 227 ipLnDTau, 227 ipLnEnrErg, 227 ipLnEnrWN, 227 ipLnEscP, 226, 227 ipLnGF, 227 ipLnGl, 227 ipLnGu, 227 ipLnHeat, 227, 281 ipLnInten, 227, 280 ipLnInwd, 226, 227 ipLnIonStg, 227 ipLnIpCont, 227 ipLnNelem, 227 ipLnNPhots, 227, 280 ipLnOpac, 227 ipLnOTS, 227, 280 ipLnPopl, 226, 227, 280 ipLnPopOpc, 226, 227, 280 ipLnPopu, 226, 227, 280 ipLnPump, 226, 227 ipLnRedis, 227 ipLnRyd, 227 ipLnTauCon, 226, 227 ipLnTauIn, 226, 227 ipLnTauTot, 226, 227 ipLnWlAng, 226, 227 ipLyHeavy, 261 ipSecIon, 305 iPunDef, 137 iter, 79, 414, 422 itermx, 79, 422 itoptm, 121 ItrDim, 93, 94, 422 KshellLimit, 137 ldDoPhoto, 415 LevTrace, 139 lgAbnSolar, 424 lgAutoIt, 79 lgDustOn, 295 lgElmtOn, 55, 423 lgFluor, 134 lgLastIt, 423 lgOpacOn, 423 lgOptimFlow, 123 lgPHFIT, 138

avdft, 295

bit32, 401

called, 414

cdsqte, 421

cextpw, 82

cextra, 82

boltzmann, 207

lgSearch, 423 lgTauOutOn, 215 lgTestOn, 139 lgVarOn, 127 limfal, 84, 427 limLevelN, 282 limpar, 119 limpun, 102 limspc, 23, 38 limTabD, 56, 61 lmhlvl, 77 logical, 418 mxstpl, 92 naming convention, 416 nAtlas, 130 ncell, 202 ndplot, 95 ndpts, 296 ndust, 295 nend, 137, 423 nflux, 202 nh, 232 nhe1, 256 nhe2, 256 nhe3lvl, 255 nhlvl, 77 nkrd, 19, 406 nmaps, 87, 111, 138 nobslm, 121 nparm, 127 npass, 422 npunlm, 109, 111 nRauch, 130 nsShells, 261 nta, 225 nterp, 39 ntff, 278 nupper, 202 nvarxt, 127 nvary, 127 nvfpnt, 127 nWerner, 130 nzdump, 98 nzlim, 421 nzone, 414, 422 occnum, 204 opac, 205, 209 ophe1f, 256 ophe2f, 256 ophf, 236 opsv, 209 otscon, 107, 204, 209 otslin, 107, 204, 209 otsmin, 134 outcon, 107, 204 outlin, 107, 204 pden, 303, 421 pdenstr, 422 perror, 303 pestrk, 232 pgas, 303 ph2lte, 245 phmlte, 245 PhotoRate, 263 phplte, 245 pinteg, 303 pirsq, 420 pnow, 303 PopLevls, 281, 284

prad, 303 PresInit, 303 presok, 303, 414 PunchLWidth, 104, 138, 622 r1r0sq, 420 radius, 419 radstr, 422 real, 417 RecomRate, 272 refcon, 106, 204 reflin, 204 rinner, 419 rmr0, 420 router, 420 sab, 295 scalem, 424 scatop, 205, 209 scmpla, 306 seccmp, 306 seche, 306 sechi, 306 secmet, 306 smetla, 306 solar, 423 SolarSave, 423 sqabs, 295 sqscat, 295 sse, 296 struc, 421 sublimat, 294 SummedCon, 205 SummedDif, 205 SummedOcc, 205 tauabs, 205 TauDmmy, 281 TauLines, 225 taumin, 221 tausct, 205 tautot, 205 te, 422 te1ryd, 207 tedust, 296 tehigh, 93 telogn, 422 telow, 93, 139 tend, 139 TeProp, 422 testr, 422 tfail, 414 tff, 278 tlast, 422 TLineARR, 229 TLineEXP, 229 tmn, 210 tsq1, 285 tsqden, 139 varang, 123, 127 vformt, 127 vincr, 120, 127 volstr, 422 vparm, 127 vtoler, 123 WavNKelv, 207 WavnNRvd, 207 WeakHeatCool, 140 widflx, 204 WindData, 225 wmole, 307, 421 x12, 135, 305

xIonFracs, 262, 424 yield, 263 variable naming convention, 416 varyN.for input script, 499 varyNU.for input script, 500 vbhum.in input script, 501 veclib routines, 77 velocity Alfvén, 86 Doppler, 81 electron, 85 mean speed, 85, 292 stored, 214 thermal, 81, 214 turbulent, 81 Verner Fe II atom, 72 version numbers, 419 VMS assign statement, 43 path, 138 Voigt function, 214 Volk, Kevin grain extra species, 74 gray grains, 74 PAH grains, 74 PN nuclei, 48 quantum grain heating, 80 stellar atmospheres, 45, 128 table density law, 61 -Wwarnings checks, 4 printed, 611 routine to print, 410 warnot, 410 wavelength output convention, 626 wavenumber, 222 Werner atmospheres, 48 werner.in input script, 502 wgadd, 411 wginit, 411 wgline, 411 wind, 304, 605 density vs radius, 304 line transfer, 304 wind command, 69 wind.in input script, 502 -X-X-ray to optical ratio, 31, 602 -Zzone attenuation, 209 defined, 4 limiting number, 18 output, 604 stopping, 94 thickness, 419 setting, 136 variables, 422